

PM 12 22 1 101

GOOD-WAY

5% SEVIN^(R) DUST

INSECTICIDE

Contains Carbaryl

FOR THE CONTROL OF CERTAIN INSECTS ON VEGETABLES
SMALL FRUITS, ORNAMENTALS AND PETS

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%

TOTAL 100%

Sevin^(R) is a registered trademark of Rhone-Poulenc Ag. Co.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT LB.


EPA REG. NO. 228-251

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584




(BACK PANEL)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly after handling. Wash hands and face before eating or



smoking. Do not contaminate food or feed. Avoid storage near food or feed products. Avoid contamination of feed, feeding troughs and watering receptacles. Food utensils such as teaspoons or tablespoons should not be used for food purposes after use in measuring pesticides. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Call a physician or poison control center immediately. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.
- IF IN EYES: Immediately flush eyes with plenty of water. Call a physician.
- IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas being treated. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **MAY KILL HONEY BEES IN SUBSTANTIAL NUMBERS.** Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.**

Apply Good-Way 5% Sevin Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7 day intervals. For best results, apply with a dust gun. Do not apply in excess.

SMALL FRUIT INSECT CONTROL

GRAPES: To control grape berry moth, Japanese beetles, June beetles, red-banded leaf roller, grape leaf folder, grape leafhopper and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. (Treat upper and lower surface of all foliage for complete coverage.) **Limitation** day of harvest. Tolerance 10 ppm.

STRAWBERRIES: To control meadow spittlebug, strawberry leaf roller and strawberry weevil. (Treat upper and lower surface of all foliage for complete coverage.) **Limitation** 1 day before harvest. Tolerance 10 ppm.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES and RASPBERRIES: To control Japanese beetles, leaf rollers, leafhoppers and raspberry aphids. Treat upper and lower surface of all foliage for complete coverage. Limitations 7 days before harvest. Tolerance 12 ppm.

PET USE

ON DOGS AND CATS: For ticks and fleas dust liberally over the entire animal, avoiding eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age. Do not use on pregnant dogs.

IN AND AROUND DOG AND CAT QUARTERS: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

GENERAL DIRECTIONS FOR USE

Apply Good-Way 5% Sevin Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results, apply with a dust gun. Do not apply in excess.

SMALL FRUIT INSECT CONTROL

GRAPES: To control grape berry moth, Japanese beetles, June beetles, red-banded leaf roller, grape leaf folder, grape leafhopper and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. (Treat upper and lower surface of all foliage for complete coverage.) Limitation 1 day before harvest. Tolerance 10 ppm.

STRAWBERRIES: To control meadow spittlebug, strawberry leaf roller and strawberry weevil. (Treat upper and lower surface of all foliage for complete coverage.) Limitation 1 day before harvest. Tolerance 10 ppm.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES and RASPBERRIES: To control Japanese beetles, leaf rollers, leafhoppers and raspberry aphids. Treat upper and lower surface of all foliage for complete coverage. Limitations 7 days before harvest. Tolerance 12 ppm.

PET USE

ON DOGS AND CATS: For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age. **IN AND AROUND DOG AND CAT QUARTERS:** For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

CROP	INSECTS CONTROLLED	DOSAGE	LIMITATIONS
Asparagus	On Spears: Asparagus beetle On Ferns And Brush: Asparagus beetle, Apache cicada	Treat upper and lower surface of all foliage for complete coverage.	1 day before harvest. Tolerance 10 ppm.
Okra	Corn earworm, stink bugs.		Day of harvest. Tolerance 10 ppm.
Peas*	Leafhoppers, Colorado potato beetle, Armyworms, grasshoppers.		3 days before harvest. Tolerance 10 ppm on peas, 100 ppm on forage.
Cauliflower*, Broccoli*, Kohlrabi*, Brussels sprouts*, Cabbage*	Flea beetle, harlequin bug, Corn earworms, armyworms, cabbage caterpillars		3 days before harvest. Tolerance 10 ppm.
Potato Eggplant	Colorado potato beetle, flea beetles, leaf hoppers		Day of harvest. Tolerance 10 ppm. (No tolerance on potatoes.)
Tomato	Tomato fruitworm, fall armyworm, European corn borer,		Day of harvest. Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Pepper	stink bugs, lace bugs and tarnished plant bugs		
Cucurbits*, Casaba Cucumber, Crenshaw, Cantaloupe, Honey Ball, Pumpkin Squash, Muskmelon, Honeydew, Watermelon.	Pickleworm, melonworm Cucumber beetles, leafhoppers, flea beetles, squash bugs		Day of harvest. Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Beans*	Mexican bean beetle Leafhoppers, bean leaf beetle, Western bean cut worm, flea beetles, Japanese beetle, velvet bean caterpillar Corn earworm, armyworms, stink bugs, tarnished plant bug In California: Lima bean pod borer, corn earworm, Lygus and stink bugs		Day of harvest. Tolerance 10 ppm on beans, 100 ppm on forage or hay.
Beet Crops*, Beets Radish, Carrots, Salsify, Turnips, Parsnips, Horseradish, Rutabaga	Flea beetles, leafhoppers, harlequin bug Six-spotted leafhopper (Aster yellows vector) Corn earworm, armyworms, cabbage caterpillars, tarnished plant bugs and stink bugs		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Do not apply 14 days before harvest on beets, salsify or turnips if tops are to be used for food purposes.			
Leafy Vegetables*, Kale Mustard Greens, Turnip Greens Dandelion, Endive, Swiss Chard, Beet Greens, Spinach, Parsley, Lettuce, Collards, Chinese Cabbage	Flea beetles, leafhoppers, harlequin bug Six-spotted leafhopper (Aster yellows vector) Corn earworm, armyworms, cabbage caterpillars, tarnished plant bugs and stink bugs		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsify tops, endive, Chinese cabbage; 12 ppm for other leafy greens.
Corn*	Corn earworm, corn rootworm adults, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers		Day of harvest. Tolerance 5 ppm in ears, Tolerance of 100 ppm on forage and fodder.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 or 3-day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

*When label directions for use are followed, forage, vines, hay and crop residues may be fed to meat or dairy animals.

ORNAMENTAL INSECT CONTROL: Good-Way 5% Sevin Dust can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Good-Way 5% Sevin Dust injures Boston ivy. Apply to the upper and lower leaf surface of all foliage for complete coverage to control such pests as:

apple aphid	mealy bugs	Japanese beetle	boxwood leaf miner	scale insects	plant bugs	willow leaf beetles	rose slug
bagworm	mimosa webworm	June beetles	elm leaf aphid	tent caterpillars	puss caterpillars	periodical cicada	leaf roller
birch leaf miner	oak leaf miners	lace bugs	elm leaf beetle	thorn bugs	psyllids	boxelder bug	
blister beetles	orange tortrix	leafhoppers	flea beetles	thrips (exposed)	rose aphid	gypsy moth	

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

STORAGE AND DISPOSAL

STORAGE: Store in tightly closed original container in a locked area away from children and domestic animals. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard in trash.

WARRANTY

Riverdale Chemical Company, the manufacturer, warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under normal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

