

SEP 30 2004

Jeannie Smith
Nufarm Americas, Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527

Dear Ms. Smith:

SUBJECT: Label Amendment and Reregistration Issues
Riverdale Team 1.33 Lawn Weed and Feed
Commercial and Residential Labels
EPA Registration No. 228-256
Your Submission Dated July 16, 2001

The labeling referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide act, as amended is acceptable with the following provisions:

For the Commercial labeling:

1. Revise the "First Aid" statement to read:

"IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor or treatment advice."

2. Under the "Hazards to Human and Domestic Animals", revise the statement of read:
"Harmful if absorbed through skin. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals."

3. a. Under "User Safety Recommendations" add: "Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

- b. "Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washable, use detergent and hot water. Keep and wash PPE separately from other laundry."

4. Under "Hazards to Humans and Domestic Animals" add: "Loaders, applicators and all other handlers must wear:

- Long sleeved shirt and long pants,
- Shoes and socks, and
- Chemical resistant gloves made of any waterproof material."

7505C
Moylan
9/30/04

On the Residential labeling:

1. Revise the "First Aid" statement to read:

"IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor or treatment advice."

2. Under the "Hazards to Human and Domestic Animals", revise the statement of read: "Harmful if absorbed through skin. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals."

This product cannot be reregistered until all of the active ingredients data requirements have been submitted and satisfied.

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,



Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

3/12

(Commercial)

RIVERDALE

TEAM® 1.33 LAWN

WEED AND FEED

A Dual Action Pre-Emergent Crabgrass Killer containing Benefin (Balan®) - Trifluralin and a Premium Turf Fertilizer.

For use on established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass for the control of Crabgrass and other listed Annual Weed Grasses on Home Lawns, Golf Courses and other Ornamental Lawns and Recreational Turf.

ACTIVE INGREDIENTS:

Benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine	0.89%
Trifluralin: (α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine)	0.44%
INERT INGREDIENTS:	98.67%
TOTAL	100.00%

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Sulfur Coated Urea	
X% Urea Nitrogen	
Available Phosphate (P_2O_5)	X%
Soluble Potash (K_2O)	X%

Primary Plant Foods derived from Urea, Sulfur Coated Urea, Ammonium Phosphate, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer Registrant) U.S. Patent No. 4,226,789

Team® and Balan® - Registered Trademarks of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT POUNDS

EPA REG. NO. 228-256

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED with COMMENTS In EPA Letter Dated:

SEP 30 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Revised 9/24/99 Added spreader settings via notification.

228-256

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with eyes, skin, or clothing. May cause skin sensitization reactions in certain individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is used after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until watering-in is complete and surface is dry.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.
- IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Riverdale Team® 1.33 Lawn Weed and Feed is a high quality fertilizer product that combines turf food to feed the lawn and a selective pre-emergence herbicide to control the below listed annual grasses.

Riverdale Team 1.33 Lawn Weed and Feed is a pre-emergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (poa annua), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. Riverdale Team 1.33 Lawn Weed and Feed controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds.

When Riverdale Team 1.33 Lawn Weed and Feed is applied as directed, the following established turfgrasses are tolerant:

- | | | |
|--------------------|--------------|--------------------|
| Kentucky bluegrass | Fescue | St. Augustinegrass |
| Perennial ryegrass | Zoysiagrass | Bahiagrass |
| Centipedegrass | Bermudagrass | |

Riverdale Team 1.33 Lawn Weed and Feed will not harm most nearby established ornamental trees and shrubs.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional)

Riverdale Team® 1.33 Lawn Weed and Feed (Commercial)

Page 3

P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

IMPORTANT PRECAUTIONS

Apply Riverdale Team 1.33 Lawn Weed and Feed to the turf prior to the germination of weed grasses since Riverdale Team 1.33 Lawn Weed and Feed will not control established weeds. Riverdale Team 1.33 Lawn Weed and Feed may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 28 pounds per 10,890 square feet (0.375 pounds Active Ingredient [A.I.] per 10,890 square feet). Riverdale Team 1.33 Lawn Weed and Feed should not be applied in the Spring to turfgrass planted the previous Fall. In Bermudagrass areas that have been over-seeded with Winter grasses, a Spring application of Riverdale Team 1.33 Lawn Weed and Feed may thin the overseeded grasses. Because of varying cultural practices, Riverdale Team 1.33 Lawn Weed and Feed should not be used on putting greens. Riverdale Team 1.33 Lawn Weed and Feed should not be used on dichondra. Riverdale Team 1.33 Lawn Weed and Feed should not be applied to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application.

Successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply Riverdale Team 1.33 Lawn Weed and Feed approximately one to two weeks prior to the anticipated germination of annual grass weed seeds, since established weeds will not be controlled. All Summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are watered-in soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Riverdale Team 1.33 Lawn Weed and Feed.

Spring germination Summer annual grasses - crabgrasses, foxtails and barnyardgrass.

A. Northern (Cool Season) Turfgrasses

For pre-emergence control, apply 28 to 37.5 pounds per 10,890 square feet (0.375 to 0.5 pounds A.I. per 10,890 square feet) of Riverdale Team 1.33 Lawn Weed and Feed in the Spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B. Southern (Warm Season) Turfgrasses

For pre-emergence control, apply 37.5 to 56.25 pounds (0.5 to 0.75 pounds A.I. per 10,890 square feet) of Riverdale Team 1.33 Lawn Weed and Feed in the late Winter or early Spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your spreader may need calibrating since proper application is based on the spreader's condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

**SPREADER SETTINGS
FOR SINGLE APPLICATION**

Spreader	50# Bag	Northern Lawns		Southern Lawns	
		Crabgrass etc.	Poa Annua	Crabgrass etc.	Poa Annua
SCOTT #35	19,450 Sq. Ft.	3	4½	-	-
	14,500 Sq. Ft.	4	5½	4	5½
	9,675 Sq. Ft.	-	-	6½	6
CYCLONE B-1	19,450 Sq. Ft.	2	3	-	-
	14,500 Sq. Ft.	3	4	3	4
	9,675 Sq. Ft.	-	-	4	4½

**FOR OPTIMUM PRE-EMERGENCE WEED CONTROL
APPLY AT THE APPROPRIATE TIME**

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE
 JAN. FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC.

1. _____ 2. _____ 3. _____
 4. _____ 5. _____ *6. _____

1. For Crabgrass etc. Northern Lawns. 4. For Crabgrass etc. Southern Lawns.
 2. For Goosegrass in Northern Lawns. 5. For Goosegrass in Southern Lawns.
 3. For Poa Annua in Northern Lawns. 6. For Poa Annua in Southern Lawns.

*Two applications spaced 8 to 10 weeks apart may be needed for season long control.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Riverdale Team 1.33 Lawn Weed and Feed Per 10,890 Square Feet	Pounds Per 1,000 Square Feet
28.00 pounds	2.5 pounds
37.50 pounds	3.4 pounds
56.25 pounds	5.1 pounds

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feed or foodstuffs. Do not stack more than two pallets high. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label

Riverdale Team® 1.33 Lawn Weed and Feed (Commercial)

Page 5

instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

WARRANTY: Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.
(PR938 042194/RV 092499)

[Editor's note: Subregistrant label may include a different (1) Fertilizer Analysis; (2) table of recommended spreader equipment and spreader settings for the fertilizer blend used; (3) Warranty Statement. Chlorine declaration is optional except where mandated by State fertilizer regulation.]

8/12

(Non-Commercial)

RIVERDALE

TEAM® 1.33 LAWN

WEED AND FEED

A Dual Action Pre-Emergent Crabgrass Killer containing Benefin (Balan®) - Trifluralin and a Premium Turf Fertilizer.

For use on established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass for the control of Crabgrass and other listed Annual Weed Grasses on Ornamental Lawns and Turf.

ACTIVE INGREDIENTS:

Benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine	0.89%
Trifluralin: (α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine)	0.44%
INERT INGREDIENTS:	98.67%
TOTAL	100.00%

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Sulfur Coated Urea	
X% Urea Nitrogen	
Available Phosphate (P_2O_5)	X%
Soluble Potash (K_2O)	X%

Primary Plant Foods derived from Urea, Sulfur Coated Urea, Ammonium Phosphate, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer Registrant) U.S. Patent No. 4,226,789

Team® and Balan® - Registered Trademarks of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT POUNDS

EPA REG. NO. 228-256

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

ACCEPTED
with COMMENTS
In EPA Letter Dated:

SEP 30 2004
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Revised 9/24/99 Added spreader settings via notifications.

228-256

Riverdale Team® 1.33 Lawn Weed and Feed (Non-Commercial)

Page 2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with eyes, skin, or clothing. May cause skin sensitization reactions in certain individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is used after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until watering-in is complete and surface is dry.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Riverdale Team® 1.33 Lawn Weed and Feed is a high quality fertilizer product that combines turf food to feed the lawn and a selective pre-emergence herbicide to control the below listed annual grasses.

Riverdale Team 1.33 Lawn Weed and Feed is a pre-emergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (poa annua), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. Riverdale Team 1.33 Lawn Weed and Feed controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds.

When Riverdale Team 1.33 Lawn Weed and Feed is applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass	Fescue	St. Augustinegrass
Perennial ryegrass	Zoysiagrass	Bahiagrass
Centipedegrass	Bermudagrass	

Riverdale Team 1.33 Lawn Weed and Feed will not harm most nearby established ornamental trees and shrubs.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

Riverdale Team® 1.33 Lawn Weed and Feed (Non-Commercial)

Page 3

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.
*Based on Inert fertilizer claimed.

IMPORTANT PRECAUTIONS

Apply Riverdale Team 1.33 Lawn Weed and Feed to the turf prior to the germination of weed grasses since Riverdale Team 1.33 Lawn Weed and Feed will not control established weeds. Riverdale Team 1.33 Lawn Weed and Feed may thin established annual bluegrass (*poa annua*) turfs and fine-leaved fescues at rates above 28 pounds per 10,890 square feet (0.375 pounds Active Ingredient [A.I.] per 10,890 square feet). Riverdale Team 1.33 Lawn Weed and Feed should not be applied in the Spring to turfgrass planted the previous Fall. In Bermudagrass areas that have been over-seeded with Winter grasses, a Spring application of Riverdale Team 1.33 Lawn Weed and Feed may thin the overseeded grasses. Because of varying cultural practices, Riverdale Team 1.33 Lawn Weed and Feed should not be used on putting greens. Riverdale Team 1.33 Lawn Weed and Feed should not be used on dichondra. Riverdale Team 1.33 Lawn Weed and Feed should not be applied to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipede grass until these turfs are well established.

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application.

Successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply Riverdale Team 1.33 Lawn Weed and Feed approximately one to two weeks prior to the anticipated germination of annual grass weed seeds, since established weeds will not be controlled. All Summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are watered-in soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Riverdale Team 1.33 Lawn Weed and Feed.

Spring germination Summer annual grasses - crabgrasses, foxtails and barnyardgrass.

A. Northern (Cool Season) Turfgrasses

For pre-emergence control, apply 28 to 37.5 pounds per 10,890 square feet (0.375 to 0.5 pounds A.I. per 10,890 square feet) of Riverdale Team 1.33 Lawn Weed and Feed in the Spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B. Southern (Warm Season) Turfgrasses

For pre-emergence control, apply 37.5 to 56.25 pounds (0.5 to 0.75 pounds A.I. per 10,890 square feet) of Riverdale Team 1.33 Lawn Weed and Feed in the late Winter or early Spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your spreader may need calibrating since proper application is based on the spreader's condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

14/12

**SPREADER SETTINGS
FOR SINGLE APPLICATION**

Spreader	50# Bag	Northern Lawns		Southern Lawns	
		Crabgrass etc.	Poa Annua	Crabgrass etc.	Poa Annua
SCOTT #35	19,450 Sq. Ft.	3	4½	—	—
	14,500 Sq. Ft.	4	5½	4	5½
	9,675 Sq. Ft.	—	—	6½	6
CYCLONE B-1	19,450 Sq. Ft.	2	3	—	—
	14,500 Sq. Ft.	3	4	3	4
	9,675 Sq. Ft.	—	—	4	4½

**FOR OPTIMUM PRE-EMERGENCE WEED CONTROL
APPLY AT THE APPROPRIATE TIME**

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE											
JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEPT.	OCT.	NOV.	DEC.
			1. _____	2. _____	3. _____						
		4. _____		5. _____						6. _____	

1. For Crabgrass etc. Northern Lawns. 4. For Crabgrass etc. Southern Lawns.
 2. For Goosegrass in Northern Lawns. 5. For Goosegrass in Southern Lawns.
 3. For Poa Annua in Northern Lawns. 6. For Poa Annua in Southern Lawns.

*Two applications spaced 8 to 10 weeks apart may be needed for season long control.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Riverdale Team 1.33
Lawn Weed and Feed
Per 10,890 Square Feet

Pounds Per
1,000 Square Feet

28.00 pounds
37.50 pounds
56.25 pounds

2.5 pounds
3.4 pounds
5.1 pounds

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a dry area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of other herbicides, insecticides, fungicides, fertilizers, or seeds.

PESTICIDE DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

WARRANTY: Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and other

12/12

Riverdale Team[®] 1.33 Lawn Weed and Feed (Non-Commercial)

Page 5

factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal ~~or~~ not reasonably foreseeable. The user assumes all risks of any such use.
(PR938 042194/RV 092499)

[Editor's note: Subregistrant label may include a different (1) Fertilizer Analysis; (2) table of recommended spreader equipment and spreader settings for the fertilizer blend used; (3) Warranty Statement. Chlorine declaration is optional except where mandated by State fertilizer regulation.]