

PM 23 228-260

TELEPHONE (800) 345-3330 1212
(708) 754-3330
FAX (708) 754-0314

Riverdale

C H E M I C A L C O M P A N Y

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3899

April 30, 1993

Ms. Joanne I. Miller
Product Manager (23)
Registration Division - H7505C
U.S. Environmental Protection Agency
Crystal Mall Bldg. #2
1921 Jefferson Davis Highway
Arlington, VA 22202

Dear Ms. Miller:


Subject: 2,4-D Exposure Reduction Program Label Revisions Due 3/1/93
Label Corrections

Following the submission of approximately sixty (60) labels, we conducted an audit and found, for more reasons than we care to go in to, it became necessary to correct the majority of the labels previously submitted.

We have made two copies of each corrected label. We respectfully request that they be entered in to our registration file as replacements. We apologize for any inconvenience this may cause the Agency.


Sincerely yours,

RIVERDALE CHEMICAL COMPANY


By Russell F. Sawyer
Regulatory Affairs Manager

RFS:ls

Enc.



2 of 2

RIVERDALE

SOLUTION^(R) WATER SOLUBLE

A High Concentrate Selective Weed Killer (Herbicide)

For Control of Many Broadleaf Weeds in both Non-Crop Areas and Certain Crop Areas, Lawns, Ponds, Ditchbanks, Pastures, and Rangelands. Also for Control of Trees by Injection.

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 96.9%

INERT INGREDIENTS: 3.1%

TOTAL 100.0%

Isomer Specific by AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid 80.5%

Solution is a Trademark of Riverdale Chemical Company

Patent Pending

Individual Soluble Bags Are Not To Be Offered For Sale

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS: 22½ lbs. (8 x 2 lb. 13 oz. packets)

EPA REG. NO. 228-260

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY


GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

CORROSIVE: Causes irreversible eye damage. May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin, or on clothing. Avoid inhaling vapor



(Back Panel cont.)

or spray mist. Remove saturated clothing as soon as possible and shower. Do not apply this product in such a manner as to, directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Please see additional Precautionary Statements in Directions For Use instructions.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention at once.
- IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention. Do not give anything by mouth to an unconscious person.
Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.
- IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. Use care to avoid spray contact or drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees, and ornamentals. Excessive amounts of this product in soil may temporarily inhibit seed germination and plant growth. Do not permit spray mist containing this product to drift onto them. Do not spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays to minimize drift. Spray drift can be lessened by (1) keeping the spray boom as low as possible, (2) applying at 20 pounds or less nozzle pressure using nozzles that produce a coarse spray pattern, and (3) applying when conditions such as wind, air stability and temperature inversions are not a factor. Do not use the same spray equipment for applying other materials to 2,4-D susceptible crops as injury may result. It is best to use a separate sprayer for application of insecticides and fungicides.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

When using on Pastures and Rangeland Grasses there is a (1) 7 day pre-grazing interval for dairy cattle; (2) 30 day preharvest interval for grass cut for hay; and (3) 3 day pre-slaughter interval for meat animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Warnings should state: "Do not enter treated areas unless wearing chemical resistant full body clothing including NIOSH approved respirator, goggles and gloves until sprays have dried." When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

DANGER: Area treated with 2,4-D amine salt on date of application. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, follow precautionary statements on label.

**ADDITIONAL PRECAUTIONARY STATEMENTS FOR AGRICULTURAL CROP USES (Including Sod Farms and Forestry workers engaged in the commercial production of wood fiber or timber products including forestry site preparation)
ALSO INDUSTRIAL/AQUATIC APPLICATIONS**

When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. For aerial applications in an enclosed cockpit and applicators applying this product from a tractor that has been a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. The rates of applications for pastures and rangelands are per acre per application site. Do not enter or allow worker entry into treated areas during the restricted-entry interval of 48 hours. There isn't any reentry restriction for Industrial/Aquatic applications, nor non-crop uses of pastures, rangelands and forests.

5 of 12

RIVERDALE Solution^(R) WATER SOLUBLE will kill or control the following weeds in addition to many other noxious plants susceptible to 2,4-D.

- | | | |
|---------------------|-----------------|--------------------|
| Alders | Goatsbeard | Furslane |
| American lotus | Goldenrod | Ragweeds |
| Arrowhead | Ground ivy | Rush |
| Artichoke | Gumweed | Russian thistle |
| Aster | Healall | Sagebrush |
| Austrian fieldcress | Hemp | St. Johnswort |
| Beggarticks | Henbit | Shepherdspurse |
| Biden | Hoary cress | Sicklepod |
| Bindweed | Honeysuckle | Smartweed |
| Bitterweeds | Horsetail | Sneezeweed |
| Bitter winter cress | Indigo | Southern wild rose |
| Blessed thistle | Indiana mallow | Sowthistle |
| Blue lettuce | Ironweed | Spanishneedles |
| Box elder | Jewelweed | Spatterdock |
| Broomweed | Jimsonweed | Stinging nettles |
| Buckhorn | Knotweed | Stinkweed |
| Bull thistle | Kochia | Sumac |
| Bulrush | Lambsquarters | Sunflower |
| Burdock | Locoweed | Sweet clover |
| Bur ragweed | Lupines | Tarweed |
| Buttercup | Mallow | Thistles |
| Canada thistle | Marshelder | Toadflax |
| Carpetweed | Mexican weed | Tumbleweed |
| Catnip | Morningglory | Velvetleaf |
| Chickweed | Musk thistle | Vervain |
| Chicory | Mustards | Vetch |
| Cockle | Nettles | Virginia creeper |
| Cocklebur | Nutgrass | Water hyacinth |
| Coffeebean | Orange hawkweed | Water lily |
| Coffeeweed | Parrotfeather | Water plantain |
| Common sowthistle | Parsnip | Water primrose |
| Creeping jenny | Pennycress | Watershield |
| Croton | Pennywort | Wild carrot |
| Curly indigo | Peppergrass | Wild garlic |
| Dandelion | Pepperweed | Wild lettuce |
| Dock | Pigweed | Wild onion |
| Dogbane | Plantains | Wild radish |
| Duckweed | Poison hemlock | Wild rape |
| Elderberry | Poison ivy | Wild strawberry |
| Flea bane (daisy) | Pokeweed | Wild sweet potato |
| Flaxweed | Poorjoe | Willow |
| Florida pusley | Povertyweed | Witchweed |
| Frenchweed | Prickly lettuce | Wormseed |
| Galinsoga | Primrose | Yellow rocket |
| | Puncture vine | |

Generally the lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply Solution during warm weather when weeds are young and growing actively. Use enough spray volume for uniform coverage by ground or air application. Unless otherwise recommended, suggested

application rates may be up to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. If only bands or rows are treated, leaving middles untreated, the dosage per crop per acre is reduce proportionately. Do not apply when temperature exceeds 90°F.

TO PREPARE THE SPRAY: Mix Solution Water Soluble only with water. Just add the recommended water to your spray tank then remove the water soluble packet from plastic outer bag, add the appropriate Solution into the spray tank, agitate for three to five minutes. Solution will make a stable water mixture. It is non-flammable. Do not handle water soluble bags or expose them to moisture since this may cause rupturing.

Individual water soluble bags are not to be offered for sale.

Notes: (1) Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds, but doing so may reduce selectivity to crops resulting in crop damage. (2) One 2 lb. 13 oz. packet of Riverdale Solution Water Soluble is equal to 5 pints of 2,4-D "4 lb." Amine. (3) Using a 2 lb. 13 oz. packet for a 5 acre coverage is equivalent to applying 2,4-D "4 lb." Amine at a rate of 1 pint per acre.

CORN (Field, Sweet and Popcorn): Pre-emergence (For annual grasses and broadleaf weeds) - Apply to soil anytime after planting but before corn emerges. Do not use on light sandy soil, or where soil moisture is low. Dissolve each 2 lb. 13 oz. water soluble packet in 4 to 30 gallons of water and apply over 1-1/8 to 2-1/4 acres. The amount of water is dependent on whether application is made with air or ground equipment.

Emergence - Dissolve each 2 lb. 13 oz. packet in 10 to 100 gallons of water per five acres. Apply just as corn plants are breaking ground.

Post-emergence (For broadleaf weeds) - Dissolve each 2 lb. 13 oz. packet in 10-100 gallons of water. This mixture will treat 5 to 10 acres. Apply when most weeds have germinated. Spray after corn emerges and until 8" tall. Use low rates on inbreds. Corn is susceptible to injury shortly after emergence and after unfolding of leaves. Do not spray during this period nor after first tassels appear. When corn is over 8" tall, use drop nozzle to keep spray off corn foliage. Spray must strike tops of weeds but should not drench corn plants. Do not apply from tasseling to dough stage. Injury to corn is most likely to occur if Solution Water Soluble is applied when corn is growing rapidly under high temperatures and high soil moisture conditions. In such conditions, use the low rate. For resistant weeds, use each 2 lb. 13 oz. packet to cover 2½ acres though corn injury may result. Do not use higher rates unless possible crop injury will be accepted. After application, delay cultivation for 8 to 10 days to allow the corn to overcome any temporary brittleness.

Preharvest: After the hard dough or denting stage, apply by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf and vines that interfere with harvesting. Dissolve the 2 lb. 13 oz. packet in 5 to 100 gallons of water to treat from 2½ to 5 acres.

WITH LIQUID NITROGEN SOLUTIONS: For late season control of young smartweed, cocklebur, annual morningglory and other annual broadleaf weeds less than 1 inch high. Field should be as clean as possible and corn 20 to 30 inches tall. The spray must be prepared by first adding the required amount of liquid nitrogen to spray tank (80 to 120 lbs. nitrogen per acre). Next dilute each 2 lb. 13 oz. packet of Riverdale Solution Water Soluble in 5 gallons of clean water. Start the agitator and (slowly) add the diluted Solution and water mixture. This combination of nitrogen and Solution will cover 5 acres. Spray immediately, maintaining continuous agitation until spray tank is empty. Direct the spray to lower 3" to 4" of corn stalk Use spray equipment designed

7212

to handle corrosive liquid nitrogen solutions. After spraying, remove any remaining solution and rinse rig thoroughly with water. Mix only one tank at a time. Do not spray during or immediately following cold weather.

SORGHUM (Milo): Post-emergence - Dissolve each 2 lb. 13 oz. packet in 10 to 100 gallons of water. Each dissolved 2 lb. 13 oz packet will cover from 5 to 7-1/2 acres. Apply when sorghum is 5" to 15" tall. Treat only after the sorghum is 5 inches high and preferable before it is 15 inches high. Spraying before 5" stage may inhibit root development. Do not treat during the boot, tasseling, or early dough stages. Reduce spray drift by keeping the boom and spray nozzle as low as possible. If crop is taller than 8 inches, use drop nozzle to keep the spray off the leaves. Temporary spray injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply Solution Water Soluble under these conditions, do not treat less than 7-1/2 acres per 2 lbs. 13 oz. packet.

NOTE: Corn & Sorghum Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

BARLEY, WHEAT, OATS AND RYE: Spring Post-emergence - In Spring grown grains, spray grain between full tillering and before the boot stage (usually 4" to 8" tall) and weeds are small, Dissolve and Solution Water Soluble 2 lb. 13 oz. packet in 5 to 100 gallons of water. For Spring grown barley, wheat and rye, apply this spray mixture on 3-1/2 to 10 acres. Oats are more sensitive to 2,4-D than other grains and should be sprayed in the Spring when well established and tillered and before jointing after crop has reached the dough stage. For oats, spray over 5 to 10 acres. In winter grains, dissolve each 2 lb. 13 oz. packet of Solution Water Soluble in 5 to 100 gallons of water. Apply mixture over 2 1/2 to 5 acres to control large weeds that will interfere with harvest or to suppress perennial weeds. Preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

SUGARCANE: Pre-emergence - Dissolve each 2 lb. 13 oz. packet in 10 to 50 gallons of water. Apply mixture over 1-1/8 acres as a blanket spray through lay-by, to aid in control of Johnsongrass seedlings and susceptible broadleaf weeds. **Post-emergence** - Dissolve each 2 lb. 13 oz. packet in 10 to 60 gallons of water. Apply over 1-2/3 to 2-1/2 acres when cane is 1' to 2' tall.

RICE: Dissolve each 2 lb. 13 oz. water soluble packet of Solution Water Soluble in 5 to 50 gallons of water. Apply over 1 1/4 to 1 1/2 acres to control curly indigo and other broadleaf weeds. Apply in the late tillering stage of rice development, at the time of first joint development (first to second green ring), usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch at early seeding, early panicle, boot, flowering or early heading growth stages.

NOTE: Some rice varieties under certain conditions can be injured by 2,4-D. Therefore before spraying, consult local Extension Service or University specialist for appropriate rates and timing of 2,4-D sprays.

ON FALLOW LAND: Dissolve each 2 lb. 13 oz. water soluble packet of Solution Water Soluble in 5 to 100 gallons of water. This mixture will cover 1 1/4 to 1 1/2 acres for control of annual broadleaf weeds. Use this mixture on 1 acre to control established perennial species, such as Canada thistle and field bindweed. Apply to weeds actively growing. Do not plant any crop for 3 months after treatment or until chemical has disappeared from soil.

GRASSES: In established pastures and rangelands, dissolve each 2 lb. 13 oz. water soluble packet in 5 to 100 gallons of water. This mixture will treat from 1-1/8 to 2-1/4 acres. Use the light rate on more easily injured grasses. Apply preferably when weeds are small and growing actively before bud stage. Fall or Spring is the best time to treat. Repeated treatments may be needed for less susceptible weeds. Treatments will kill or injure alfalfa, sweet clover and other legumes. White clover (including Ladino) may be injured by light application but recovers; repeated treatments will kill it. In some area dichondra, bentgrasses, carpet, buffalo, and St. Augustine grasses may be injured. Usually colonial bents are more tolerant than creeping types; velvets are most easily injured. Where bentgrass predominates, dissolve each 2 lb. 13 oz. water soluble packet in 10 to 100 gallons of water and spray this mixture over 10 acres. Repeat application after 3 weeks.

GRASS SEED CROPS: Dissolve each 2 lb. 13 oz. water soluble packet in 25 to 200 gallons of water. Spray mixture will treat 1-1/8 to 4-1/2 acres. Apply in Spring or Fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seeding grass only after the five leaf stage, using 4 1/2 acre rate to control small seeding weeds. After the grass is well established higher rates of up to one 2 lb. 13 oz. packet per 1-1/8 acres can be used to control hard-to-kill annual or perennial weeds. For best results, do not use on bentgrass unless grass injury can be tolerated.

CONTROL OF SOUTHERN WILD ROSE: On roadsides and fencerows, use 2 lb. 13 oz. packet of Solution Water Soluble plus 4-8 ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of one (2 lb. 13 oz.) packets of Solution Water Soluble per 1.13 acres per application per site.

ADDITIONAL PRECAUTIONARY STATEMENTS FOR TURF CONTROL

When using this product, wear long-sleeved shirt, long pants, socks, shoes, rubber gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection. In addition to the clothing and eye protection listed above, commercial mixers/loaders/applicators must wear chemical resistant in place of rubber gloves except when the product is applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

BROADLEAF WEED CONTROL IN NON-CROPLAND GRASS AREAS SUCH AS LAWNS, GOLF COURSES, CEMETERIES AND PARKS, AIRFIELDS, RIGHTS-OF-WAY, FENCEROWS, ROADSIDES, VACANT LOTS, DRAINAGE DITCHEBANKS AND SIMILAR PLACES: Use 1 to 2 packets (2 lb. 13 oz. each) of this product in 15 to 50 gallons of water per acre. On turf, apply a maximum of one (2 lb. 13 oz.) packets of this product per 1.13 acres per application per site. Treat when weeds are young and growing well. Do not use on dichondra or other herbaceous ground covers. Do not use on creeping grasses such as bent except for spot treatment nor on freshly seeded turf until grass is well established. Reseeding of lawns should be delayed following treatment. With Spring application, reseed in the Fall. With Fall application, reseed in Spring. Legumes are usually damaged or killed. Thoroughly wet weeds when applying this mixture. Bindweed, Whitetop, Perennial sow thistle, Blue

9 of 12

lettuce, Bur ragweed, Canada thistle and other noxious perennials somewhat resistant to 2,4-D will require repeated treatments to kill. To control small areas of woody plants, such as Willows, Honeysuckle, Virginia creeper, Alders and other susceptible to 2,4-D, use 1 to 2 water soluble, 2 lb. 13 oz. packets in 100 gallons water; spray to thoroughly wet plants when in full leaf. Re-treat as necessary for control of regrowth and seedlings. In general, it is better to cut tall woody growth and spray suckers when 2 to 4 feet high.

TREE INJECTIONS (Pine Release): To control hardwoods, such as Oaks, Hickory, Maple, Pecan, Elm, Sumac, Sweetgum and Hawthorn in forest and other non-crop areas, dissolve a 2 lb. 13 oz. packet of Solution Water Soluble in 1½ gallons of water and apply in a concentrate tree injector calibrated to apply 0.75 ml. per injection. Space injections 2" apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species such as Hickory, Dogwood, Red maple, Blue beech and Ash, make injections 1" to 1-1/2" apart, edge to edge. Treatment may be made at any time of the year. For best results, injections should be made during growing season, May 15 - October 15.

FOR AQUATIC WEEDS IN STILL LAKES, PONDS, DRAINAGE DITCHES AND MARSHES: Aerial Application - Use one 2 lb. 13 oz. packet of Solution Water Soluble in 5 to 15 gallons of water to cover one surface acre.

Boat Application - Use 2 lb. 13 oz. packet of Solution Water Soluble in 50 to 100 gallons of water per acre. Uniform coverage is essential. Avoid submerging plants after treatment. Application should be made when leaves are fully developed above water line and plants are actively growing.

Consult your State Game and Fish Department or Water Control Agency prior to application of this product for aquatic weed control

Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard, treat 1/3 to 1/2 of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatments along the shore and proceed outwards in bands to allow fish to move into untreated areas. Do not apply within 1500 feet of potable or irrigation water intakes.

ADDITIONAL PRECAUTIONARY STATEMENTS FOR BRUSH CONTROL

The maximum application rate for forestry site preparation is 1 gallon 6 ounces per acre per application per site. There isn't any restrictions on the annual maximum number of applications for brush control. No reentry restriction for non-agricultural uses.

WEEDS AND BRUSH ON IRRIGATION CANAL DITCHBANKS - SEVENTEEN WESTERN STATES: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington and Wyoming.

For control of annual and perennial broadleaf weeds, dissolve one 2 lb. 13 oz. packet in 30 to 200 gallons of water and apply mixture over 1½ to 3 acres. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder to control weeds, a repeat spray after 3 to 4 weeks using the same rates may be needed for maximum results. Apply no more than two treatments per season.

For woody brush and patches of perennial broadleaf weeds mix two 2 lbs. 13 oz. packets of Solution Water Soluble in 150 gallons of water. Wet foliage thoroughly using about 1 gallon of solution per square rod.

SPRAYING INSTRUCTION: Apply with low pressure (10 to 40 psi) power spray equipment mounted on truck, tractor or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is fairly calm; 5 mph or less. Do not use on small canals (less than 10 cfs) where water will be used for drinking purposes.

Boom spraying onto water surfaces must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than two-foot over-spray onto water with an average of less than one-foot over-spray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated area for at least 7 days after spraying. Water within banks should not be fished.

Local conditions may affect the uses of this chemical as shown on this label. Consult State Experiment Station or Extension Service Weed Specialist for specific recommendations for local weed problems and for information on possible lower dosages.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secure warehouse or storage building. This product should be stored in a cool, dry location. Do not store near seeds, fertilizers, insecticides, or fungicides. Container should be opened in a well ventilated area. All containers should be kept tightly sealed when not in use. Do not contaminate water, food or feedstuff by storage of pesticide disposal.


PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide spray mixtures or rinsate is a violation of Federal law. If container is damaged or if pesticide has spilled, contain all spillage. Clean up all spilled material with broom. Place in a closed, labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Dispose of empty outer plastic bags and shipping case in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable, the user assumes all risks of any such use.

(EPC: 02393)



11212

Solution^(R)
WATER SOLUBLE

A High Concentrate
Selective Weed Killer

KEEP OUT OF REACH OF CHILDREN

DANGER

See Product Label on Carton for
Additional Precautionary Statements
And Statement of Practical Treatment

EPA REG. NO. 228-260

EPA EST. NO. 228-IL-1

RIVERDALE CHEMICAL (LOGO)

