



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OCT 13 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Mary Beth Endres
Product Registration
Nufarm Americas, Inc.,
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Name "Dri-Clean Herbicide"
EPA Reg. No. 228-260
Application Dated September 23, 2009

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 09/23/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,
Rachel C. Hillman
Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2/18

(A)	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460 Application for Pesticide:	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number Please Attach An Original Form
-----	---	---	--

Section I

1. Company/Product Number 228-260	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Solution Water Soluble	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas Inc 150 Harvester Drive, Suite 200 Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain Below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - explain below

NOTIFICATION
OCT 13 2009

Explanation: Use additional page(s) if necessary. (For Section I and Section II).
 Notification for addition of Alternate Brand Name PR Notice 98-10.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No <i>*Certification must be submitted.</i>	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes," Unit Package Wt. No. Per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes," Package Wt. No. Per Container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
---	---	---	--

3. Location of Net Contents Information <input type="checkbox"/> Label or <input type="checkbox"/> Container	4. Size(s) of Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label or <input type="checkbox"/> On Labeling accompanying product
6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input type="checkbox"/> Paper Glued <input type="checkbox"/> Stenciled		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Mary Beth Endres	Title Registration Manager	Telephone No. (Include Area Code) 630-455-2000
--------------------------	-------------------------------	---

<p style="text-align: center;">Certification</p> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager	
4. Typed Name Mary Beth Endres	5. Date 09-23-2009	



3
18

Nufarm Americas Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527
Telephone: (630) 455.2000 Facsimile: (630) 455.2001
www.us.nufarm.com

Overnight Courier

September 23, 2009

Linda Arrington
Notification and Minor Formulations Team Leader
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

**Subject: Addition of Alternate Brand Name for
Solution Water Soluble, EPA Reg. No. 228-260**

Dear Ms. Arrington:

Nufarm Americas Inc. would like to add an alternate brand name for the above referenced product. We would like the **alternate brand name to be:**

Dri-Clean Herbicide


We feel that this label change is permitted and consistent with PR Notice 98-10 and the name is not false or misleading. One copy of labeling with the name change is attached.

Should you have any questions please contact me at the address and telephone number above or by email listed below.

Sincerely,

Mary Beth Endres
Registration Manager
Nufarm Americas Inc.
Email: maryb.endres@us.nufarm.com

Enclosure



4/18

Dri-Clean[®] Herbicide

NOTIFICATION

OCT 13 2009

**A HIGH CONCENTRATE, SELECTIVE WEED KILLER (HERBICIDE)
FOR CONTROL OF MANY BROADLEAF WEEDS IN NON-CROP AND CERTAIN CROP
AREAS, LAWNS, PONDS, DITCHBANKS, PASTURES, RANGELANDS, AND OTHER
LISTED SITES. ALSO FOR CONTROL OF TREES BY INJECTION.**

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 96.9%

OTHER INGREDIENTS: 3.1%

TOTAL: 100.0%

Isomer Specific by AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid 80.5%

**KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE BELOW FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

NOTE: Spanish language is optional


EPA REG. NO. 228-260
EPA EST. NO. 228-IL-1

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



Net Contents: 19 Oz. (538.6g) Packet

000228-00260.20090921b.EPA.ABN.Notification



5/18

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER / PELIGRO**

Corrosive. Causes irreversible eye damage. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Handlers using engineering controls must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks,
- In addition, mixers and loaders must wear chemical-resistant gloves and a chemical-resistant apron.

All other handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Chemical-resistant footwear plus socks,
- Goggles or faceshield,
- Chemical-resistant gloves,
- Chemical-resistant apron when applying as a spray to citrus or if exposed to the concentrate, and
- Chemical-resistant headgear if overhead exposure.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls Statements: Water-soluble packets when used correctly qualify as a closed loading system under the WPS. Mixers and loaders using water-soluble packets (1) must wear the PPE specified above for mixers and loaders and (2) must be provided, have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown a NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40CFR 170.240 (d) (6)].

USER SAFETY RECOMMENDATIONS	
Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.	
Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.	
Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTICE TO PHYSICIAN
Probable mucosal damage may contraindicate the use of gastric lavage.

6/18

ENVIRONMENTAL HAZARDS

For Terrestrial Uses: This pesticide may be hazardous to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in waste adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

For Aquatic Uses: Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

SPRAY DRIFT REDUCTION

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product."

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted-Entry Interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the Restricted-Entry Interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any water-proof material, shoes plus socks, protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

GENERAL INFORMATION

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition et al vs. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web site: <http://www.epa.gov/espp>.

WEEDS CONTROLLED

This product will kill or control the following weeds in addition to many other noxious plants susceptible to 2,4-D.

Alder	Dandelion	Musk thistle	Spatterdock
American lotus	Dock	Mustard	Stinging nettle
Arrowhead	Dogbane	Nettle	Stinkweed
Artichoke	Duckweed	Nutgrass	Sumac
Aster	Elderberry	Orange hawkweed	Sunflower
Austrian fieldcress	Flea bane (daisy)	Parrotfeather	Sweet clover
Beggartick	Flixweed	Parsnip	Tarweed
Biden	Florida pusley	Pennycress	Thistle
Bindweed	Frenchweed	Pennywort	Toadflax
Bitterweed	Galinsoga	Peppergrass	Tumbleweed
Bitter wintercress	Goatsbeard	Pepperweed	Velvetleaf
Blessed thistle	Goldenrod	Pigweed	Vervain
Blue lettuce	Ground ivy	Plantain	Vetch
Box elder	Gumweed	Poison hemlock	Virginia creeper
Broomweed	Healall	Poison ivy	Water hyacinth
Buckhorn	Hemp	Pokeweed	Water lily
Bull thistle	Henbit	Poorjoe	Water plantain
Bulrush	Hoary cress	Povertyweed	Water primrose
Burdock	Honeysuckle	Prickly lettuce	Watershield
Bur ragweed	Horsetail	Primrose	Wild carrot
Buttercup	Indigo	Puncture vine	Wild garlic
Canada thistle	Indiana mallow	Purslane	Wild lettuce
Carpetweed	Ironweed	Ragweed	Wild onion
Catnip	Jewelweed	Rush	Wild radish
Chickweed	Jimsonweed	Russian thistle	Wild rape
Chicory	Knotweed	Sagebrush	Wild strawberry
Cockle	Kochia	St. Johnswort	Wild sweet potato
Cocklebur	Lambsquarter	Shepherdspurse	Willow
Coffeebean	Locoweed	Sicklepod	Witchweed
Coffeeweed	Lupine	Smartweed	Wormseed
Common sowthistle	Mallow	Sneezeweed	Yellow rocket
Creeping jenny	Marshelder	Southern wild rose	
Croton	Mexican weed	Sowthistle	
Curly indigo	Morningglory	Spanishneedle	

Generally the lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply this product during warm weather when weeds are young and growing actively. Use enough spray volume for uniform coverage by ground or air application. Unless otherwise recommended, suggested application rates may be up to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. If only bands or rows are treated, leaving middles untreated, the dosage per crop per acre is reduced proportionately. Do not apply when temperature exceeds 90°F.

TO PREPARE THE SPRAY

Mix this product only with water. Just add the recommended water to your spray tank, then add the appropriate amount of product into the spray tank and agitate for three to five minutes. This product will make a stable water mixture. It is non-flammable.

NOTES: (1) Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds, but doing so may reduce selectivity to crops resulting in crop damage. (2) Every 2 packets (38 ounces) of this product is equal to 5 pints of 2,4-D "4 lb." Amine. (3) Using 2 packets (38 ounces) for a 5 acre coverage is equivalent to applying 2,4-D "4 lb." Amine at a rate of 1 pint per acre.

SELECTIVE WEEDING IN CROPS

WITH LIQUID NITROGEN SOLUTIONS

For late season control of young Smartweed, Cocklebur, Annual Morningglory and other annual broadleaf weeds less than 1 inch high. Field should be as clean as possible and corn 20 to 30 inches tall. The spray must be prepared by first adding the required amount of liquid nitrogen to spray tank (80 to 120 lbs. nitrogen per acre). Next dilute 2 packets (38 ounces) of this product in 5 gallons of clean water. Start the agitator and (slowly) add the diluted solution and water mixture. This combination of nitrogen and solution will cover 5 acres. Spray immediately, maintaining continuous agitation until spray tank is empty. Direct the spray to lower 3 to 4 inches of corn stalk. Use spray equipment designed to handle corrosive liquid nitrogen solutions. After spraying, remove any remaining solution and rinse rig thoroughly with water. Mix only one tank at a time. Do not spray during or immediately following cold weather.

CITRUS FLOOR

Broadleaf Control

Use 18 to 27 ounces of this product in 10-100 gallons of water to treat one acre. For band or spot treatment calculate rates according to the actual portion of acre treated. Apply as a directed spray onto weeds to point of runoff when weeds are young and actively growing. Repeat applications through the growing and dormant season as needed.

Ground application only using low silhouette or shielded sprayer to avoid contact with green plant tissue. Use with drift retardant to increase deposition and reduce drift. Make up to two applications through the dormant and growing seasons as needed. Do not harvest citrus within 40 days of application. Do not graze or feed cover crops from treated orchards to livestock.

CORN

(Field, Sweet and Popcorn)

Preemergence (For annual grasses and broadleaf weeds)

Apply to soil any time after planting but before corn emerges. Do not use on light sandy soil, or where soil moisture is low. Dissolve 2 packets (38 ounces) in 4 to 30 gallons of water and apply over 2 acres. The amount of water is dependent on whether application is made with air or ground equipment.

Emergence

Dissolve 2 packets (38 ounces) in 10 to 100 gallons of water per 5 acres. Apply just as corn plants are breaking ground.

Postemergence (For broadleaf weeds)

Dissolve 2 packets (38 ounces) in 10 to 100 gallons of water. This mixture will treat 5 to 10 acres. Apply when most weeds have germinated. Spray after corn emerges and until 8 inches tall. Use low rates on inbreds. Corn is susceptible to injury shortly after emergence and after unfolding of leaves. Do not spray during this period nor after first tassels appear. When corn is over 8 inches tall, use drop nozzle to keep spray off corn foliage. Spray must strike tops of weeds but should not drench corn plants. Do not apply from tasseling to dough stage. Injury to corn is most likely to occur if this product is applied when corn is growing rapidly under high temperatures and high soil moisture conditions. In such conditions, use the low rate. Do not use higher rates unless possible crop injury will be accepted. After application, delay cultivation for 8 to 10 days to allow the corn to overcome any temporary brittleness.

Preharvest

After the hard dough or denting stage, apply by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as Bindweed, Cocklebur, Dogbane, Jimsonweed, Ragweed, Sunflower, Velvetleaf and vines that interfere with harvesting. Dissolve 2 packets (38 ounces) in 5 to 100 gallons of water to treat from 2.5 to 5 acres.

USE PRECAUTIONS FOR FIELD AND POPCORN

Do not use treated crop as fodder for 7 days following application.

The preharvest interval (PHI) is 7 days.

Maximum of 3.5 pounds of product/acre per crop cycle.

Preplant or Preemergence

Limited to one preplant or preemergence application per crop cycle.

Maximum of 1.25 pounds of product /acre per application.

Postemergence

Limited to one postemergence application per crop cycle.

Maximum of 0.6 pound of product/acre per application.

Preharvest

Limited to one preharvest application per crop cycle.
Maximum of 1.87 pounds of product /acre per application.

USE PRECAUTIONS FOR SWEET CORN

Do not use treated crop as fodder for 7 days following application.
The preharvest interval (PHI) is 45 days.
Minimum of 21 days between applications.
Maximum of 1.87 pounds of product /acre per crop cycle.

Preplant or Preemergence

Limited to one preplant or preemergence application per crop cycle.
Maximum of 1.25 pounds of product /acre per application.

Postemergence

Limited to one postemergence application per crop cycle.
Maximum of 0.6 pound of product/acre per application.

**SORGHUM
(Milo)**

Postemergence

Dissolve 2 packets (38 ounces) in 10 to 100 gallons of water. This will cover from 5 to 7.5 acres. Apply when sorghum is 5 to 15 inches tall. Treat only after the sorghum is 5 inches high and preferably before it is 15 inches high. Spraying before the 5-inch stage may inhibit root development. Do not treat during the boot, tasseling, or early dough stages. Reduce spray drift by keeping the boom and spray nozzle as low as possible. If crop is taller than 8 inches, use drop nozzle to keep the spray off the leaves. Temporary spray injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply this product under these conditions, do not treat less than 7.5 acres per 2 packets (38 ounces).

NOTE: Corn & Sorghum Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed specialist for this information.

USE PRECAUTIONS FOR SORGHUM

The preharvest interval (PHI) is 30 days.
Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.
Limited to 1 application per crop cycle.
Maximum of 1.25 pounds of product/acre per application.

BARLEY, WHEAT, OATS AND RYE

Spring Postemergence

In spring grown grains, spray grain between full tillering and before the boot stage (usually 4 to 8 inches tall) when weeds are small. Dissolve 2 packets (38 ounces) of this product in 5 to 100 gallons of water. For spring grown barley, wheat and rye, apply this spray mixture on 3.5 to 10 acres. Oats are more sensitive to 2,4-D than other grains and should be sprayed in the spring when well established and tillered and before jointing after crop has reached the dough stage. For oats, spray over 5 to 10 acres. In winter grains, dissolve 2 packets (38 ounces) in 5 to 100 gallons of water. Apply mixture over 3.8 to 5 acres to control large weeds that will interfere with harvest or to suppress perennial weeds. Preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

USE PRECAUTIONS FOR SMALL GRAINS

The preharvest interval (PHI) is 14 days.

Postemergence

Limited to one postemergence application per crop cycle. Maximum of 1.56 pounds of product /acre per application.

Preharvest

Limited to one preharvest application per crop cycle.
Maximum of 0.63 pound of product /acre per application.
Limited to 2.2 pounds of product/acre per crop cycle.

SUGARCANE

Preemergence

Dissolve 2 packets (38 ounces) in 10 to 50 gallons of water. Apply mixture over 1-1/8 acres as a blanket spray through lay-by, to aid in control of Johnsongrass seedlings and susceptible broadleaf weeds.

Postemergence

Dissolve 2 packets (38 ounces) in 10 to 60 gallons of water. Apply over 1-2/3 to 2-1/2 acres when cane is 1 to 2 feet tall.

USE PRECAUTIONS FOR SUGARCANE

Do not harvest cane prior to crop maturity.
Do not apply more than 5 pounds of product / acre per crop cycle.

Preemergence

Limited to one application per crop cycle.
Maximum of 2.5 pounds of product/acre per application.

Postemergence

Limited to one application per crop cycle.
Maximum of 2.5 pounds of product/acre per application.

RICE*

Use 1 to 2 packets (19 to 38 ounces) of this product in 5 to 10 gallons of water. Apply over 1-1/4 to 1-1/2 acres to control Curly indigo and other broadleaf weeds. Apply in the late tillering stage of rice development, at the time of first joint development (first to second green ring), usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch at early seeding, early panicle, boot, flowering or early heading growth stages.

NOTE: Some rice varieties under certain conditions can be injured by 2,4-D. Therefore before spraying, consult local Extension

***Not for use in California**

USE PRECAUTIONS FOR RICE

The preharvest interval (PHI) is 60 days.
Maximum of 1.87 pounds of product /acre per crop cycle.

Preplant

Limited to one preplant application per crop cycle.
Maximum of 1.25 pounds of product/acre per preplant application.

Postemergence

Limited to one postemergence application per crop cycle.
Maximum of 1.87 pounds of product /acre per postemergence application.

ON FALLOW LAND

Dissolve 2 packets (38 ounces) of this product in 5 to 100 gallons of water. This mixture will cover 1-1/4 to 1-1/2 acres for control of annual broadleaf weeds. Use this mixture on 1 acre to control established perennial species, such as Canada thistle and Field bindweed. Apply to weeds actively growing. Do not plant any crop for 3 months after treatment or until chemical has disappeared from soil.

USE PRECAUTIONS ON FALLOW LAND

Limited to 2 applications per year.
Maximum of 2.5 pounds of product/acre per application.
Minimum of 30 days between applications Service or University specialist for appropriate rates and timing of 2,4-D sprays.

GRASSES

The rates of application are per application per site. In established pastures and rangelands, dissolve 2 packets (38 ounces) in 5 to 100 gallons of water. This mixture will treat from 1-1/8 to 2-1/4 acres. Use the light rate on more easily injured grasses. On rangeland apply a maximum of 2 packets (38 ounces) of this product per 1.13 acres per application per site. Apply preferably when weeds are small and growing actively before bud stage. Fall or spring is the best time to treat. Repeated treatments may be needed for less susceptible weeds. Treatments will kill or injure alfalfa, sweet clover and other legumes. White clover (including Ladino) may be injured by light application but recovers; repeated treatments will kill it. In some areas dichondra, bentgrasses, carpet, buffalo, and St. Augustine grasses may be injured. Usually colonial bents are more tolerant than creeping types; velvets are more easily injured. Where bentgrass predominates, dissolve 2 packets (38 ounces) in 10 to 100 gallons of water and spray this mixture over 10 acres.

USE PRECAUTIONS FOR GRASSES

The preharvest interval (PHI) is 7 days (cut forage for hay).

Postemergence

Limited to 2 applications per year.
Maximum of 2.5 pounds of product/acre per application.
Minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

GRASS SEED CROPS

Dissolve 2 packets (38 ounces) in 25 to 200 gallons of water. Spray mixture will treat 1-1/8 to 4-1/2 acres. Apply in spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seeding grass only after the five-leaf stage, using 4-1/2 acre rate to control small seeding weeds. After the grass is well established higher rates of up to 2 packets (38 ounces) per 1-1/8 acres can be used to control hard-to-kill annual or perennial weeds. For best results, do not use on bentgrass unless grass injury can be tolerated.

USE PRECAUTIONS FOR GRASS SEED CROPS

Do not apply more than 2 application per year.
The maximum application rate is 2.5 pounds of product per acre per application.
The minimum retreatment interval is 21 days.

11
18

APPLE AND PEAR ORCHARDS

Broadleaf Weeds

NON-BEARING TREES (well established - one year or older), and **BEARING TREES** before and after bloom - Use 27 ounces in 10 to 100 gallons of water per acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of an acre treated. Apply as a directed spray onto the weeds to point of runoff when weeds are young and actively growing (pre-bud to early bud stage).

USE PRECAUTIONS FOR POME FRUIT

The preharvest interval (PHI) is 14 days.
Do not cut orchard floor forage for hay within 7 days of application.

Postemergence

Limited to 2 applications per crop cycle.
Maximum of 2.5 pounds of product/acre per application.
Minimum of 75 days between applications.

GRAPE VINEYARDS

Established at least 3 years - Field Bindweed (Morning Glory), Canada Thistle and other 2,4-D susceptible broadleaf weeds. Dilute 18 to 27 ounces in 10 to 100 gallons of water to treat one acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of an acre treated. Apply when weeds are in the bud to early bloom stage and growing vigorously. Apply after shatter following bloom and before grape shoots reach the ground or during dormant season. Use a hooded boom and low pressure flooding nozzles to deliver coarse droplets.

USE PRECAUTIONS FOR GRAPES

Grapes are extremely sensitive to 2,4-D.
Use a direct application so no 2,4-D contacts grape leaves and young shoots or stems.
For use only in California.
The preharvest interval (PHI) is 100 days.
Limited to 1 application per crop cycle.
Maximum of 1.7 pounds of product/acre per application

STONE FRUIT, PISTACHIOS, AND NUT ORCHARDS

Broadleaf Weeds

Use 27 ounces in 10 to 100 gallons of water per acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of an acre treated.

Apply as a directed spray onto the weeds to point of runoff when weeds are young and actively growing (pre-bud to early bud stage). Make up to two applications through the dormant or growing seasons as needed. Do not harvest stone fruits within 40 days of application. Do not harvest nuts within 60 days of application. Do not graze or feed cover crops from treated orchards to livestock.

USE PRECAUTIONS FOR STONE FRUIT

The preharvest interval (PHI) is 40 days.
Do not cut orchard floor forage for hay within 7 days of application.

Postemergence

Limited to 2 applications per crop cycle.
Maximum of 2.5 pounds of product/acre per application.
Minimum of 75 days between applications.

USE PRECAUTIONS FOR TREE NUT

The preharvest interval (PHI) is 60 days.
Do not cut orchard floor forage for harvest within 7 days of application.

Postemergence

Limited to 2 applications per crop cycle.
Maximum of 2.5 pounds of product/acre per application.
Minimum of 30 days between applications.

USE PRECAUTIONS FOR FILBERTS

The preharvest interval (PHI) is 45 days.

Postemergence

Limited to 4 applications per year.
The maximum application rate is 1.2 pounds of product per 100 gallons of spray solution per application.
The minimum retreatment interval is 30 days.

PRECAUTIONS IN APPLYING 2,4-D IN ORCHARDS AND VINEYARDS

Apply only after irrigation and allow maximum time before the next irrigation. Do not apply around fruit trees or vines with hand gun. Use only on flat, fan-type nozzles and low pressures - 20 to 25 pounds. Use a lined boom applicator which can be calibrated and which will deposit the spray uniformly. Avoid contact with fruit, foliage, stems or lower limbs of trees or vines. Apply precisely and uniformly to prevent damage to the trees or vines and to obtain satisfactory weed control. Do not apply during windy periods or extremely high temperatures. Do not spray bare ground. To avoid leaching, do not apply to dry soils. Apply when soil is moist and do not irrigate for 5 to 7 days after application.

FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS IN SOYBEANS (PREPLANT ONLY)

General Information: This product is a herbicide that provides control of many emerged susceptible annual and perennial broadleaf weeds. This product may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. This product should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below. Do not use any tillage operations between application of this product and planting soybeans.

Mixing Instructions: Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

Applications Procedures: Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

APPLICATION TIMING AND USE RATES		
2,4-D Formulation Used	Maximum Rate (per acre)	When to Apply (Days prior to planting soybeans)
Dri-Clean Herbicide	At a rate of 0.5 lb. a.e./acre <u>Amount</u> <u>Acreage</u> 2 packets (38 ounces) 4.50	NOT LESS THAN 15 DAYS Limited to 2 applications per crop cycle
	At a rate of 1.0 lb. a.e./acre <u>Amount</u> <u>Acreage</u> 2 packets (38 ounces) 2.25	NOT LESS THAN 30 DAYS Limited to 1 application per crop cycle

WEEDS CONTROLLED

- | | | |
|-----------------------------|------------------------|-------------------------|
| Alfafa* | Garlic-wild* | Ragweed-common |
| Bindweed* | Horseweed or marestail | Ragweed-giant |
| Bullnettle | Ironweed | Shepherdspurse |
| Bittercross-smallflowered | Lambsquarters-common | Smartweed-Pennsylvania* |
| Buttercup-smallflowered | Lettuce-prickly | Sowthistle-annual |
| Carolina geranium | Morningglory-annual | Speedwell |
| Cinquefoil-common and rough | Moustaill | Thistle-Canada* |
| Clover-red* | Mustard-wild* | Thistle-bull |
| Cocklebur-common | Pennycress-field | Velvetleaf |
| Dandelion* | Peppergrass* | Vetch-hairy* |
| Evening primrose-cutleaf | Purslane-common | Virginia copperleaf |

*These species are only partially controlled.

For best weed control at time of treatment, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage. The response of individual weed species to this product is variable. Consult your local county or State Agricultural Extension Service or crop consultant for advice.

APPLICATION RESTRICTIONS AND PRECAUTIONS: Important Notice - Unacceptable injury to soybeans planted in fields treated with this product may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide applications until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Do not use on low organic sandy soils (<1.0%).

Do not apply this product when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants.

The maximum rate per crop cycle is 1.2 pounds of product per acre.

Livestock Grazing Restriction: Do not feed hay, forage or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

In fields treated with this product, plant soybean seed as deep as practical or at least 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is completely covered.

Do not apply this product prior to planting soybeans, if you are not prepared to accept the results of soybean injury, including possible loss of stand and yield.

Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D use.

ORNAMENTAL TURF

(Golf Courses, Cemeteries, Parks, Sports Fields, Turfgrass, Lawns and other Grass Areas)

Use 2 packets (38 ounces) of this product per acre in 15 to 50 gallons of water. On turf, apply a maximum of 30 ounces of this product per acre per application per site. Treat when weeds are young and growing well. Do not use on dichondra or other herbaceous ground covers. Do not use on creeping grasses such as bent except for spot treatment nor on freshly seeded turf until grass is well established. Reseeding of lawns should be delayed following treatment. With spring application, reseed in the fall. With fall application, reseed in spring. Legumes are usually damaged or killed. Thoroughly wet weeds when applying this mixture. Bindweed, Whitetop, Perennial sowthistle, Blue lettuce, Bur ragweed, Canada thistle and other noxious perennials somewhat resistant to 2,4-D will require repeated treatments to kill.

USE PRECAUTIONS FOR ORNAMENTAL TURF

Postemergence

Do not apply more than 2 broadcast applications per year.

The maximum application rate is 1.8 pounds of product per acre per application.

The maximum seasonal rate is 3.6 pounds of product per acre, excluding spot treatments.

GENERAL WEED CONTROL IN NON-CROPLAND AREAS

(Fencerows, Hedgerows, Roadsides, Ditches, Rights-of-Way, Utility Power Lines, Railroads, Airports, and Industrial Sites)

Use 2 packets (38 ounces) of this product per acre in 15 to 50 gallons of water. Treat when weeds are young and growing well. Do not use on dichondra or other herbaceous ground covers. Do not use on creeping grasses such as bent except for spot treatment nor on freshly seeded turf until grass is well established. Reseeding of lawns should be delayed following treatment. With spring application, reseed in the fall. With fall application, reseed in spring. Legumes are usually damaged or killed. Thoroughly wet weeds when applying this mixture. Bindweed, Whitetop, Perennial sowthistle, Blue lettuce, Bur ragweed, Canada thistle and other noxious perennials somewhat resistant to 2,4-D will require repeated treatments to kill. To control small areas of woody plants, such as Willows, Honeysuckle, Virginia creeper, Alders and others susceptible to 2,4-D, use 4 packets (76 ounces) per acre in adequate water (100 gallons); spray to thoroughly wet plants when in full leaf. Re-treat as necessary for control of regrowth and seedlings. In general, it is better to cut tall woody growth and spray suckers when 2 to 4 feet high.

SPOT TREATMENT IN NON-CROP AREAS: To control broadleaf weeds in small areas with a hand sprayer, use two packets (38 ounces) of this product in 50 gallons of water, fill sprayer and apply to thoroughly wet all foliage.

CONTROL OF SOUTHERN WILD ROSE: On roadsides and fencerows use 4 packets (76 ounces) of this product plus 4 to 8 ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required.

USE PRECAUTIONS FOR NON-CROPLAND

Postemergence (annual and perennial weeds)

Limited to 2 applications per year.

Maximum of 2.5 pounds of product per acre per application.

Minimum of 30 days between applications.

Postemergence (woody plants)

Limited to 1 application per year.

Maximum of 5 pounds of product per acre per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

TREE INJECTIONS IN FORESTS AND OTHER NON-CROP AREAS

(Pine Release)

To control hardwoods, such as Oaks, Hickory, Maple, Pecan Elm, Sumac, Sweetgum and Hawthorn in forest and other non-crop areas, dissolve 2 packets (38 ounces) of this product in 1-1/2 gallons of water and apply in a concentrate tree injector calibrated to apply 0.75 ml per injection. Space injections 2 inches apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species such as Hickory, Dogwood, Red maple, Blue beech and Ash, make injections 1 to 1-1/2 inches apart, edge to edge. Treatment may be made at anytime of the year. For best results, injections should be made during growing season, May 15 to October 15. Limited to one injection application per year, maximum of 2 ml of 4.0 lbs ae formulation per injection site.

AQUATIC APPLICATIONS

WEEDS AND BRUSH ON IRRIGATION CANAL DITCHBANKS - SEVENTEEN WESTERN STATES: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington and Wyoming. For control of annual and perennial broadleaf weeds, dissolve two packets (38 ounces) of this product per acre in enough water to wet all parts of the brush foliage, stem and bark. This may require 30 to 200 gallons of water per acre. Treat when weeds are young and actively growing before the bud or early bloom stage. For woody brush and patches of perennial broadleaf weeds, mix 4 packets (76 ounces) of this product in 150 gallons of water. Wet foliage thoroughly using about 1 gallon of solution per square rod.

SPRAYING INSTRUCTIONS: Apply with low pressure (10 to 40 psi) power spray equipment mounted on truck, tractor or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is fairly calm; 5 mph or less. Do not use on small canals (less than 10 cfs) where water will be used for drinking purposes.

Boom spraying onto water surfaces must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than two-foot over-spray onto water with an average of less than one-foot over-spray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated areas for at least 7 days after spraying. Water within banks should not be fished.

Local conditions may affect the uses of this chemical as shown on this label. Consult State Experiment Station or Extension Service weed specialist for specific recommendations for local weed problems and for information on possible lower dosages.

DITCHBANK APPLICATION

Postemergence

Limited to 2 applications per season.

Maximum of 2.5 pounds of product per acre per application.

Minimum of 30 days between applications.

Spot treatment permitted.

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance. Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

$$\text{Average Width (ft.)} \times \text{Average Depth (ft.)} \times \text{Average Velocity (ft. per sec.)} = \text{CFS}$$

For ditchbank weeds

Do not allow boom spray to be directed onto water surface.

Do not spray across stream to opposite bank.

For shoreline weeds

Allow no more than 2 foot overspray onto water

FOR AQUATIC WEEDS SUCH AS WATER HYACINTH IN WATERS THAT ARE QUIESCENT OR SLOW MOVING (LAKES, PONDS, RESERVOIRS, CANALS, RIVERS, BAYOUS, STREAMS, DRAINAGE DITCHES, MARSHES, ETC.)

Aerial Application - Use 2 packets (38 ounces) of this product in 5 to 15 gallons of water to cover 1.2 surface acres. Use drift control spray equipment or thickening agents mixed into the spray solution. Apply through standard boom systems with a minimum of 5 gallons of spray mix per acre.

Surface Application - Use 2 to 4 packets (38 to 76 ounces) of this product in 50 to 100 gallons of water per 1 acre. Uniform coverage is essential. Avoid submerging plants after treatment. Application should be made when leaves are fully developed above water line and plants are actively growing. Use power sprayers operated with a boom or spray gun mounted on a boat, tractor or truck. Thorough wetting of foliage is essential for maximum control. Special precautions such as the use of low pressure, large nozzles and thickening agents should be taken to avoid spray drift in areas of sensitive crops. For DIRECTA-SPRAY™ operation, use this product with 1 pint of drift control agent in 50 to 100 gallons of water. For other applications, follow the drift control agent label for mixing directions. Consult your State Game and Fish Department or Water Control Agency prior to application of this product for aquatic weed control. Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard, treat 1/3 to 1/2 of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatments along the shore and proceed outward in bands to allow fish to move into untreated areas. For large bodies of weed infested waters, leave buffer strips of at least 100 feet wide and delay treatment of these strips for 4 to 5 weeks or until the dead vegetation has decomposed. Repeat as necessary to kill regrowth and plants missed in previous application.

FLOATING AND EMERGENT WEEDS

Limitations and Restrictions for Floating and Emergent Weeds

- Maximum of 5 pounds of product per surface acre per application.
- Limited to 2 applications per season.
- Minimum of 21 days between applications.
- Spot treatments are permitted.
- Apply to emergent aquatic weeds in ponds, lakes, reservoirs, marshes, bayous, drainage ditches, non-irrigation canals, rivers, and streams that are quiescent or slow moving.
- Coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for aquatic applications.

Water Use for Floating and Emergent Weeds

1. Water for irrigation or sprays:

- A. If treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as pastures, turf, or cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at anytime after the 2,4-D aquatic application.
- B. Due to potential phytotoxicity considerations, the following restrictions are applicable:
If treated water is intended to be used to irrigate or mix sprays for plants grown in commercial nurseries and greenhouses; and other plants or crops that are not labeled for direct treatment with 2,4-D, the water must not be used unless one of the following restrictions has been observed:

- i. A setback distance from functional water intake(s) of greater than or equal to 600 feet was used for the application, or,
- ii. A waiting period of 7 days from the time of application has elapsed, or,
- iii. An approved assay indicates that the 2,4-D concentration is 100 ppb (0.1 ppm) or less at the water intake. Wait at least 3 days after application before initial sampling at water intake.

2. Drinking water (potable water):

A. Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits.

The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in the water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb at the time of consumption.

B. For floating and emergent weed applications, the drinking water setback distance from functioning potable water intakes is greater than or equal to 600 feet.

C. If no setback distance of greater than or equal to 600 feet is used for application, applicators or the authorizing organization must provide a drinking water notification prior to a 2,4-D application to the party responsible for public water supply or to individual private water uses. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water.

The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

Example:

Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting must include the day and time of application. Posting may be removed if analysis of a sample collected at the intake 3 or more days following application shows that the concentration in the water is less than 70 ppb (100 ppb for irrigation or sprays), or after 7 days following application, whichever occurs first.

Text of notification:

Wait 7 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water, irrigation, or sprays, unless water at functioning drinking water intakes is tested at least 3 days after application and is demonstrated by assay to contain not more than 70 ppb 2,4-D (100 ppb for irrigation or sprays). Application Date: _____ Time: _____

D. Following each application of this product, treated water must not be used for drinking water unless one of the following restrictions has been observed:

- i. A setback distance from functional water intake(s) of greater than or equal to 600 feet was used for the application, or,
- ii. A waiting period of at least 7 days from the time of application has elapsed, or,
- iii. An approved assay indicates that the 2,4-D concentration is 70 ppb (0.07 ppm) or less at the water intake. Sampling for drinking water analysis should occur no sooner than 3 days after 2,4-D application. Analysis of samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

E. Note: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

F. Drinking water setback distances do not apply to terrestrial applications of 2,4-D adjacent to water bodies with potable water intakes.

SUBMERSED WEEDS

Limitations and Restrictions for Submersed Weeds

- Maximum of 13.4 pounds of product per acre-foot per application.
- Limited to 2 applications per season.
- Apply to aquatic weeds in ponds, lakes, reservoirs, marshes, bayous, drainage ditches, non-irrigation canals, rivers, and streams that are quiescent or slow moving.
- Do not apply within 21 days of previous application.
- When treating moving bodies of water, applications must be made while traveling upstream to prevent concentration of 2,4-D downstream from the application.
- Coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for such use.

Directions for Use Associated with the Specific Use Pattern.

Table 1. Amount of 2,4-D to Apply for a Target Subsurface Concentration			
Surface Area (Acre)	Average Depth (Feet)	For Typical Conditions 2 ppm 2,4-D ae/acre-foot (Pounds)	For Difficult Conditions* 4 ppm 2,4-D ae/acre-foot (Pounds)
1	1	5.4	10.8
	2	10.8	21.6
	3	16.2	32.4
	4	21.6	43.2
	5	27.0	54.0

* Examples include spot treatment of pioneer colonies of Eurasian Water Milfoil and certain difficult to control aquatic species.

Water Use for Submersed Weeds

1. Water for irrigation or sprays:

- A. If treated water is intended to be used only for crops or non-crop areas that are labeled for direct treatment with 2,4-D such as pastures, turf, or cereal grains, the treated water may be used to irrigate and/or mix sprays for these sites at anytime after the 2,4-D aquatic application.
- B. Due to potential phytotoxicity and/or residue considerations, the following restrictions are applicable:
If treated water is intended to be used to irrigate or mix sprays for unlabeled crops, non-crop areas or other plants not labeled for direct treatment with 2,4-D, the water must not be used unless one of the following restrictions has been observed:
 - i. A setback distance described in the Drinking Water Setback Table was used for the application, or,
 - ii. A waiting period of 21 days from the time of application has elapsed, or,
 - iii. An approved assay indicates that the 2,4-D concentration is 100 ppb (0.1 ppm) or less at the water intake. See Table 3 for the waiting period after application but before taking the initial sampling at water intake.

2. Drinking water (potable water):

- A. Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits.
The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of 2,4-D in the water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that 2,4-D concentrations in potable water do not exceed 70 ppb at the time of consumption.
- B. For submersed weed applications, the drinking water setback distances from functioning potable water intakes are provided in Table 2. Drinking Water Setback Distance (below).
- C. If no setback distance from the Drinking Water Setback Table (Table 2) is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut-off all potable water intakes prior to a 2,4-D application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water.
The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under state or local law or as a condition of a permit.

Example:

Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application. Posting may be removed if analysis of a sample collected at the intake no sooner than stated in Table 3 (below) shows that the concentration in the water is less than 70 ppb (100 ppb for irrigation or sprays), or after 21 days following application, whichever occurs first.

Text of notification:

Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water, irrigation, or sprays, unless water at functioning drinking water intakes is tested no sooner than (insert days from Table 3) and is demonstrated by assay to contain not more than 70 ppb 2,4-D (100 ppb for irrigation or sprays).
Application Date: _____ Time: _____

D. Following each application of this product, treated water must not be used for drinking water unless one of the following restrictions has been observed:

- i. A setback distance described in the Drinking Water Setback Distance Table was used for the application, or,
- ii. A waiting period of at least 21 days from the time of application has elapsed, or,
- iii. An approved assay indicates that the 2,4-D concentration is 70 ppb (0.07 ppm) or less at the water intake. Sampling for drinking water analysis should occur no sooner than stated in Table 3. Analysis of samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

E. Note: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

F. Drinking water setback distances do not apply to terrestrial applications of 2,4-D adjacent to water bodies with potable water intakes.

3. Except as stated above, there are no restrictions on using water from treated areas for swimming, fishing, watering livestock or domestic purposes.

Table 2. Drinking Water Setback Distance for Submersed Weed Applications			
Application Rate and Minimum Setback Distance (feet) From Functioning Potable Water Intake			
1 ppm*	2 ppm*	3 ppm*	4 ppm*
600	1200	1800	2400
* ppm acid equivalent target water concentration			

Table 3. Sampling for Drinking Water Analysis After 2,4-D Application for Submersed Weed Applications			
Minimum Days After Application Before Initial Water Sampling at the Functioning Potable Water Intake			
1 ppm*	2 ppm*	3 ppm*	4 ppm*
5	10	10	14
* ppm acid equivalent target water concentration			

STORAGE AND DISPOSAL

Do not contaminate water, feed or foodstuff by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secure warehouse or storage building. This product should be stored in a cool, dry location. Do not store near seeds, fertilizers, insecticides, or fungicides. Container should be opened in a well-ventilated area. All containers should be kept tightly sealed when not in use. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If container is damaged or if pesticide has spilled, contain all spillage. Clean up all spilled material with broom. Place in a closed, labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. After removal of all PVA packets, dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

18
18

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV092109N)

Dri-Clean is a registered trademark of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.

OPTIONAL FRONT PANEL TEXT

FOR CONTROL OF MANY BROADLEAF WEEDS IN GRAPE VINEYARDS, APPLE, PEAR, STONE FRUIT AND NUT ORCHARDS. ALSO FOR NON-CROP USE IN ORNAMENTAL TURF, AND NON-CROPLAND AREAS.