

SEP 29 2003

Russell F. Sawyer  
Nufarm Americas, Inc.  
Riverdale - A Nufarm Company  
1333 Burr Ridge Parkway, Suite 125A  
Burr Ridge, IL 60527-0866


Dear Mr. Sawyer:

**SUBJECT: Riverdale Tri-aPower Selective Herbicide  
Label Amendment  
EPA Registration No. 228-262  
Your Application Dated July 9, 2003**

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

The amended label supersedes all previously accepted ones. A stamped copy is enclosed for your records. Please submit one copy of your printed label before you release the product for shipment.

Sincerely yours,



Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

RIVERDALE

TRI-POWER SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL  
IN ORNAMENTAL LAWNS AND TURF GRASSES  
ALSO FOR WOODY PLANTS, ROADSIDES, AND SIMILAR NON-CROP AREAS

CONTROLS (or KILLS): Dandelion, Chickweed, Black medic, Knotweed, Plantain,  
Oxalis, Clover, Cocklebur, Thistle and many other species of  
broadleaf weeds; some of which are listed on this label.

CONTAINS MCPA, MECOPROP-p AND DICAMBA

GET THE OPTICAL ADVANTAGE

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	40.42%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	7.99%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	3.97%
INERT INGREDIENTS:	47.62%
TOTAL	100.00%

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	33.00%	3.1 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	6.60%	0.6 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	3.30%	0.3 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

Riverdale, Tri-Power and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company.

For Use By Professional Turf Maintenance Personnel,  
Landscaping or Commercial Applicators Only

**KEEP OUT OF REACH OF CHILDREN**

**DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que  
se la explique a usted en detalle.

**SEE INSIDE BOOKLET FOR FIRST AID  
AND ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA REG. NO. 228-262      NET CONTENTS      GALS.      EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS, 60527-0866

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

SEP 29 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-262

NOTE: Spanish language is optional

Revised 1/31/03 Changed First Aid Statement per PR 2001-1 via Amendment.  
7/9/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

(Inside Booklet)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER - PELIGRO**

Corrosive. This concentrate causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear face shield or goggles when mixing and loading this product. After product has been diluted in accordance with the directions for use on turf below, goggles, face shield or safety glasses are not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID STATEMENT**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eyelids open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**ENVIRONMENTAL HAZARDS**

Drift of runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT.** USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Tri-Power is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas; also for Woody Plants, Roadsides and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

**USE PRECAUTIONS**

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control (or kill) of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control (or kill) weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

Tri-Power can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of Tri-Power.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop's Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding the Tri-Power).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

**WEEDS CONTROLLED (OR KILLED) BY TRI-POWER**

Bedstraw, Bindweed, Black Medic, Buckhorn, Burdock, Buttercup, Canada thistle, Carpetweed, Chickweed, Chicory, Clover, Cocklebur, Dandelion, Dock, Dog fennel, English daisy, Florida pusley, Frenchweed, Goldenrod, Ground ivy, Hawkweed, Healall, Heartleaf, drymary, Henbit, Jimsonweed, Knotweed, Kochia, Lambsquarter, Lespedeza, Mallow, Morningglory, Mustard, Oxalis, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison ivy, Poison oak, Purslane, Ragweed, Red clover, Red sorrel, Sheep sorrel, Shepherdspurse, Smartweed, Speedwell, Spurge, Spurweed, Stinging nettle, Stitchwort, Thistle, Toadflax, Veronica, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wood sorrel, Yarrow and many other broadleaf weeds.

**ORNAMENTAL LAWNS AND TURFS**

**Cool Season Grasses:** Apply Tri-Power Selective Herbicide at the rate of 2.5 to 4.1 pints in 20 to 240 gallons of spray solution per acre (0.9 to 1.5 fluid ounces in 0.5 to 4 gallons of water per 1,000 square feet) to control (or kill) weeds growing in ornamental lawn turf planted to Kentucky bluegrass, perennial ryegrass, tall fescue or fine fescue.

**Warm Season Grasses:** Apply Tri-Power Selective Herbicide at the rate of 2 to 3 pints in 20 to 240 gallons of spray solution per acre (0.7 to 1.1 fluid ounces in 0.5 to 4 gallons of water per 1,000 square feet) to control (or kill) weeds growing in Bermuda grass or zoysia grass. Applications to other grass species including bahiagrass, kikuyugrass, buffalograss, centipedegrass, carpetgrass or St. Augustinegrass should be avoided unless injury can be tolerated. Avoid applications to warm season grasses emerging from Winter dormancy unless a slight delay in Spring green-up can be tolerated.

**Bentgrass Turf (Other Than Golf Course Greens and Tees):** On closely mowed bentgrass, apply Tri-Power at a maximum rate of 2-1/2 pints per acre (0.9 fluid ounces in 3 gallons of water per 1,000 square feet) in May or mid-August through September when weeds are actively growing. Exercise care when applying to avoid overdosing bentgrass, or turf injury may result. Slight turf yellowing should disappear after about 1 week.

**Bentgrass Turf (Putting and Bowling Greens):** On closely mowed bentgrass, apply this product at a maximum rate of 1 fluid ounce in 5 gallons of water per 1,500 square feet - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

**NOTE**

Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

**Herbi™ Controlled Droplet Applicator:** For Cool Season Grasses - Add 1-2/3 to 2-1/2 pints of Tri-Power to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution. For Listed Warm Season Grasses - Apply 1-1/4 to 1-2/3 pints of Tri-Power, then add water to make total of 5 pints of mixture. While walking at approximately 1 pace (3 feet) per second spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of Tri-Power must be applied when grass is stressed from heat, drought, etc. Herbi™ is a trademark of North American Micron.

**Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators:** For Cool Season Grasses - Use Tri-Power at the rate of 2-1/2 to 3-1/2 pints per acre (0.9 to 1.25 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). For Listed Warm Season Grasses - Use 2 to 3 pints of Tri-Power per acre (0.7 to 1.1 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

**NOTE:** For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

**OTHER NON-CROP AREAS**

**Roadsides (Including Aprons and Guard Rails), Rights-of-Way, and Other Similar Non-Crop Areas:** For the control of broadleaf weeds, mix at a rate of 1/4 to 3/4 gallon of Tri-Power per 50 to 300 gallons of water. This mixture will cover 43,500 square feet. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate of 3 ounces of Tri-Power per gallon of water and spray to thoroughly wet all foliage.

**For Control of Woody Plants:** Apply to both stems and foliage any time from the time foliage

