

PM23

228-263

TELEPHONE (800) 345-3330
(708) 754-3330
FAX (708) 754-0314

log 5

Riverdale

C H E M I C A L C O M P A N Y

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

February 22, 1993

CERTIFICATION


I, being an authorized representative of Riverdale Chemical Company certify that all containers of Riverdale Super Green Weed and Feed, EPA Reg. No. 228-263 produced by June 15, 1994, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify this company by January 1, 1995, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Sincerely,

RIVERDALE CHEMICAL COMPANY

James K. Champion
James K. Champion
President

JKC:ls



CPDA Chemical Producers
and Distributors
Association

PLCAA
Pesticide Labeling
Code of Best Practices

2 of 5

(Non-Commercial Label)

RIVERDALE

SUPER GREEN WEED AND FEED
X-X-X

Premium Lawn Care

This product is designed for the control of dandelion and many other broadleaf weeds. See back panel for a complete list of weeds controlled by this product.

TRIPLE ACTION

Feeds Kills Dandelions and other Broadleaf weeds Contains Iron for Color

Treats (Covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENT:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid* 0.86%

INERT INGREDIENTS: 99.14%
TOTAL 100.00%

*Isomer Specific AOAC Method, Equivalent to 2,4-Dichlorophenoxyacetic Acid .. 0.57%

GUARANTEED ANALYSIS:°

Total Nitrogen (N) XZ
XZ Ammoniacal Nitrogen
XZ Urea Nitrogen
Available Phosphoric Acid (P₂O₅) XZ
Soluble Potash (K₂O) XZ
Primary Plant Foods derived from Di-Ammonium Phosphate,
Urea, Triple Super Phosphate, and Muriate of Potash
Iron (Fe) XZ
Secondary Plant Food derived from Iron Sulfate
Chlorine Maximum XZ

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT - 20 LBS.

EPA REG. NO. 228-263

EPA EST. NO. 228-IL-1

Revised A/O 1/14/93

MANUFACTURED BY
 RIVERDALE CHEMICAL COMPANY
 GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dusts have settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.
 IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
 IF ON SKIN: Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

NOTE: Do not apply on Lippia, St. Augustine, Bentgrass or Dichondra. DO NOT USE this product where desirable clovers are present.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns, such as:

- | | | | |
|-----------|------------|---------------|-----------|
| Bindweed | Fanweed | Lambsquarters | Pigweed |
| Buckhorn | Goldenrod | Mallow | Plantain |
| Burdock | Ground ivy | Morningglory | Purslane |
| Chickweed | Healall | Mustard | Ragweed |
| Chicory | Henbit | Nettles | Smartweed |
| Cocklebur | Jewelweed | Oxalis | Speedwell |

Creeping Jenny	Jimsonweed	Pennywort	Thistle
Dandelion	Kochia	Peppergrass	Yarrow

and many other weeds susceptible to 2,4-D

WHEN TO APPLY: For best results apply when weeds are young and actively growing in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not over-apply. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

HOW TO APPLY: This 20 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at suggested rate of 4 lbs. per 1000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS: ONE PASS APPLICATION:°	MAKE	SETTING
	Cyclone	4
	Sears	5
	Scott's	5
	Jackson	5
	Greenview	5

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE. Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (ERP 061594)

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

505
505