

COPY



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP 16 1993

Russell F. Sawyer
RIVERDALE CHEMICAL CO.
425 West 194th Street
Glenwood, IL 60425

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SUBJECT: Label Amendment Submission of 08/17/93 in Response to PR Notice 93-7
EPA Reg. No. 228-264
RIVERDALE TRIPLET SELECTIVE HERBICIDE


Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully **REMOVES THE PRODUCT FROM THE SCOPE** of PR Notice 93-7 once the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements, the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

Sincerely,


James Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Enclosures



Recycled/Recyclable
Printed with Soy/Caro
contains at least 50%

198

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Russell F Sawyer
RIVERDALE CHEMICAL CO
425 WEST 194TH ST.
GLENWOOD IL 60425

Subject: Label Amendment Submission of 08/17/93
in Compliance with the PR Notice 93-7
EPA Reg Nr. 228-264
RIVERDALE TRIPLET SELECTIVE HERBICIDE

Modify the exclusionary statement to conform to the EPA-
approved language specified in PRN 93-11.

ACCEPTED
with COMMENTS
In EPA Letter Dated

FEB 16 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-264

RIVERDALE
TRIPLET(R)
SELECTIVE HERBICIDE

Selective Broadleaf Weed Control For Turfgrasses
To Control Dandelion, Clover, Henbit, Plantains, Wild Onion, and
Many Other Broadleaf Weeds

Contains 2,4-D, Mecoprop and Dicamba

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	30.56%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	16.34%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	2.77%

INERT INGREDIENTS:	50.33%
TOTAL	100.00%

Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	25.38%, 2.44 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	13.50%, 1.30 lbs./gal.
***3,6-Dichloro-o-anisic Acid	2.30%, 0.22 lbs./gal.

Triplet(R) Is A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

**SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT**

NET CONTENTS

EPA REG. NO. 228-264

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive, causes irreversible eye damage. Do not get in eyes, or on

NOTE: Spanish language is optional

Revised A/O 4/20/93

X
3

4078

skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed or inhaled. Wear face shield or goggles and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. In addition to the clothing and eye protection listed above, commercial mixers/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention at once.
- IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention.
- IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of

5058

the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Triplet^(R) is for use on Lawns, Turf (excluding pastures, sod farms and turf grown to product seed) and Golf Fairways, Aprons, Tees and Roughs, Parks, Cemeteries and similar non-crop areas.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Triplet Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either state or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

- | | | |
|----------------|---------------|--------------------|
| Bedstraw | Dock | Morningglory |
| Black medic | Ground ivy | Peppergrass |
| Buckhorn | Healall | Pigweed |
| Burdock | Henbit | Plantain |
| Chicory | Knotweed | Poison ivy |
| Chickweed | Lambsquarters | Poison oak |
| Clover | Lespedeza | Purslane |
| Dandelion | Mallow | Ragweed |
| Sheep sorrel | Spurge | Wild lettuce |
| Shepherdspurse | Wild carrot | Wild onion |
| Speedwell | Wild garlic | Yarrow |

and other broadleaf weeds.

X 4

ORNAMENTAL LAWNS & TURF

APPLICATIONS:

RATES:

Cool Season Grasses - Use this product at the rate of 3-4 pints per acre (1.1 to 1.5 fl. oz. per 1000 sq. ft.).

For Normal Application - Use 40-80 gallons of water per acre. (1-2 gallons water per 1,000 square feet)

For Professional Lawn Maintenance - Use 1-5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses - Such as St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses - Apply Triplet at a maximum rate of 2-2½ pints in 40-200 gallons of water per acre. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass:

On closely mowed bentgrass (putting and bowling greens) apply this product at a maximum rate of one fluid ounce in 5 gal. water per 1,500 square ft. - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

HerbiTM Controlled Droplet Applicator: For cool season grasses - add 2-3 pints of Triplet to the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 sq. ft. (approx. ¾ acre). Do not overlap spray patterns.

For listed warm season grasses - add 1.5 to 2 pints of Triplet to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Other Controlled Droplet Applicators: (CDA), atomizers, and Spinning disk Applicators:

For cool season grasses - Use Triplet at the rate of 3-4 pints per acre (1.1 to 1.5 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1-4 water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use 2 to 2½ pints of Triplet per acre (.75 to .9 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1-4 gal. of water per acre is normal for this type equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume Equipment

Use as little as 5 gallons water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

HerbiTM is a trademark of North American Micron

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers also disposable by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.



WARRANTY

Riverdale Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(ERP 102393)(PR937&8 042194)



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 16 1994

Russell F. Sawyer
RIVERDALE CHEMICAL CO.
425 West 194th Street
Glenwood, IL 60425

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SUBJECT: Label Amendment Submission of 08/17/93 in Response to PR Notice 93-7
EPA Reg. No. 228-264
RIVERDALE TRIPLET SELECTIVE HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully **REMOVES THE PRODUCT FROM THE SCOPE** of PR Notice 93-7 once the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements, the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

BEST AVAILABLE COPY

Sincerely,

James Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Enclosures



Recycled/Recyclable
Printed with Soy/Canola inks
contains at least 50% recycled fiber

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Russell F Sawyer
RIVERDALE CHEMICAL CO
425 WEST 194TH ST.
GLENWOOD IL 60425

Subject: Label Amendment Submission of 08/17/93
in Compliance with the PR Notice 93-7
EPA Reg Nr. 228-264
RIVERDALE TRIPLET SELECTIVE HERBICIDE

Modify the exclusionary statement to conform to the EPA-
approved language specified in PRN 93-11.

ACCEPTED
with COMMENTS
In EPA Letter Dated

Feb 10 1974

RIVERDALE
TRIPLET(R)
SELECTIVE HERBICIDE

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-264

Selective Broadleaf Weed Control For Turfgrasses
To Control Dandelion, Clover, Henbit, Plantains, Wild Onion, and
Many Other Broadleaf Weeds

Contains 2,4-D, Mecoprop and Dicamba

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	30.56%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	16.34%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	2.77%

INERT INGREDIENTS:	50.33%
TOTAL	100.00%

Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	25.38%, 2.44 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	13.50%, 1.30 lbs./gal.
***3,6-Dichloro-o-anisic Acid	2.30%, 0.22 lbs./gal.

Triplet(R) Is A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que
se la explique a Usted en detalle.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS

EPA REG. NO. 228-264

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive, causes irreversible eye damage. Do not get in eyes, or on



skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed or inhaled. Wear face shield or goggles and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. In addition to the clothing and eye protection listed above, commercial mixers/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention at once.
- IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.
- IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of

BEST AVAILABLE COPY

the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ADDITIONAL PRECAUTIONS FOR SOD FARM APPLICATIONS

When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield or goggles and chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands, face and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. There isn't any restriction on the annual maximum number of broadcast applications for sod farms. Do not enter or allow worker entry into treated areas during the restricted-entry interval of 48 hours.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Triplet Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either state or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Bedstraw	Dock	Morningglory
Black medic	Ground ivy	Peppergrass
Buckhorn	Healall	Pigweed
Burdock	Henbit	Plantain
Chicory	Knotweed	Poison ivy
Chickweed	Lambsquarters	Poison oak
Clover	Lespedeza	Purslane
Dandelion	Mallow	Ragweed
Sheep sorrel	Spurge	Wild lettuce
Shepherdspurse	Wild carrot	Wild onion
Speedwell	Wild garlic	Yarrow

and other broadleaf weeds.

ORNAMENTAL LAWNS & TURF**APPLICATIONS:**

Cool Season Grasses - Use this product at the rate of 3-4 pints per acre (1.1 to 1.5 fl. oz. per 1000 sq. ft.).

RATES:

For Normal Application - Use 40-80 gallons of water per acre.
(1-2 gallons water per 1,000 square feet)

For Professional Lawn Maintenance - Use 1-5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses - Such as St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses - Apply Triplet at a maximum rate of 2-2½ pints in 40-200 gallons of water per acre. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass:

On closely mowed bentgrass (putting and bowling greens) apply this product at a maximum rate of one fluid ounce in 5 gal. water per 1,500 square ft. - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If

hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

HerbiTM Controlled Droplet Applicator: For cool season grasses - add 2-3 pints of Triplet to the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 sq. ft. (approx. 3/4 acre). Do not overlap spray patterns.

For listed warm season grasses - add 1.5 to 2 pints of Triplet to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Other Controlled Droplet Applicators: (C.A.), atomizers, and Spinning disk Applicators: For cool season grasses - Use Triplet at the rate of 3-4 pints per acre (1.1 to 1.5 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1-4 water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use 2 to 2½ pints of Triplet per acre (.75 to .9 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1-4 gal. of water per acre is normal for this type equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume Equipment Use as little as 5 gallons water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

HerbiTM is a trademark of North American Micron

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers also disposable by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(ERP 102393) (PR938 042194)