



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OCT 2.5 2005

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Theodore D. Head Nufarm Americas Inc. 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527

Dear Mr. Head:

Subject: Supplemental Labeling and Maximum Use Rates for 2,4-D Riverdale Tripplett Selective Herbicide EPA Registration Number 228-264 Application Submission dated September 13, 2005

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, is acceptable provided that the following revision is made:

- 1. Delete "Forestry" from "Maximum Use Rates for Forestry and Non-Crop Sites", since this product is not labeled for forestry uses.
- 2. Modify the Maximum Use Rates box to add a statement similar to the following: The rates on this supplemental labeling supersede the application rates on container label and/or label booklet for Non-Crop Sites.

One copy of the label stamped "Accepted with Comments" is enclosed for your records. Please submit one copy of the final printed supplemental label before the product is released for shipment. If you have any questions, please contact Tracy White by phone at (703) 308-0042 or via email at <u>white.tracy@epa.gov.</u>

Sincerely,

oanne D. Miller

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Enclosure

SUPPLEMENTAL LABELING READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR TRIPPLET SELECTIVE HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. ACCEPTED bel" as used in this supplemental labeling refers to the label booklet for this product and this supplement. with COMMENTS In EPA Letter Dated: OCI 2 5 2075 **Riverdale Tripplett** Under the Federal Insecticide, Fungicide, and Rodenticide Act Nufarm as amended, for the pesticide **Selective Herbicide** registered under EPA Reg. No. EPA Reg. No. 228-264 28-264 Maximum Use Rates For Forestry and Non-Crop Sites

DIRECTIONS FOR USE

{

ť

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the label booklet for this product must be followed.

Also refer to additional "GENERAL INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Product should not be used in or near greenhouses.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

NON-TURF AREAS

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1.5 to 3.5 quarts of Triplet per acre in enough water to wet all parts of the foliage. This may require 50 to 300

gallons of water per acre. This mixture will cover 43,560 square feet. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of Triplet per gallon of water and spray to thoroughly wet all foliage. For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to ao dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

Woody Brush: Add 1 to 1-1/2 gallons of Triplet per acre to wet all parts of the brush foliage, stem and bark. This may require up to 200 to 600 gallons of water per 43,560 sqaure feet depending upon the height and thickness of the brush. Mix thoroughly before spraying

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions for application.

* Maximum Seasonal Application Rate Rate to non-crop and forestry sites is 4 pounds 2,4-D Acid equivelant per acre per application site.

NOTICE

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" in the label booklet. If terms are not acceptable, return unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under the "WARRANTY LIMITATIONS AND DISCLAIMER".

Manufactured By Nufarm Americas Inc. Burr Ridge, IL 60521

Ţ.

09/05

(

