

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

TERM OF ISSUANCE

Conditional

NAME OF PESTICIDE PRODUCT

Riverdale Sweet Sixteen Weed  
and Feed with CL/DAP

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

┌ Riverdale Chemical Company ┐  
425 West 194th Street  
Glenwood, IL 60425-1584

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/Cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase, "EPA Registration No. 228-269" to your label before you release the product for shipment.
3. One year storage stability data at ambient warehouse conditions must be submitted to EPA within 15 months of the date on this registration notice.
4. An acute inhalation toxicity test must be submitted to EPA within 9 months of the date on this registration notice.
5. Revise the Precautionary Statement to include; "Harmful if absorbed through skin. Causes moderate eye injury. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling".

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

6. Revise the statement of practical treatment to include; "If in eyes, Flush with plenty of water. Get medical attention if irritation persists. If on skin, Wash with plenty of soap and water. Get medical attention."

7. On the label, under "Environmental Hazards", delete "or wetlands (swamps, bogs, marshes, and potholes)." and replace this with, "to areas where surface water is present or to intertidal areas below the mean high water mark."

Submit three (3) copies of your final printed labeling, with changes referred to above, before you release the product for shipment. Refer to the A-79 previously mailed to you for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(c). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

**BEST AVAILABLE COPY**

CONCURRENCES

SYMBOL	H7505C							
SURNAME	Robbins							
DATE	7/24/92							

RIVERDALE

SWEET SIXTEEN  
WEED AND FEED WITH CL/DAP™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	1.01%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.40%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.09%

INERT INGREDIENTS: .....	98.50%
TOTAL .....	100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid .....	0.82%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid .....	0.33%
***3,6-Dichloro-o-Anisic Acid .....	0.08%

GUARANTEED ANALYSIS:°

Total Nitrogen (N) .....	XZ
XZ Ammoniacal Nitrogen	
XZ Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	XZ
Soluble Potash (K <sub>2</sub> O) .....	XZ

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum .....	ACCEPTED with COMMENTS in EPA Letter Dated	XZ
-----------------------------	--	----

The urea nitrogen in this product has been coated to provide XZ coated slow release urea nitrogen.

JUL '28 1992

†CL/DAP™ - Trademark of Riverdale Chemical Company

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act as amended, this pesticide is registered with EPA Reg. No. 228-269

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 16 LBS.

EPA REG. NO. 228-

EPA EST. NO. 228-IL-1

BEST AVAILABLE COPY

## MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY  
GLENWOOD, ILLINOIS 60425-1584

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wear long pants, long-sleeved shirt and shoes when applying this product. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Do not allow persons and pets on treated areas until dusts have settled.

## STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.  
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.  
IF ON SKIN: Wash skin with soap and water.

## ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH CL/DAP is a stable mixture of three herbicides. It is for use on Lawns and Ornamental Turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns such as:

Beggarstick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbert-leaved saltbush, Hawkweed, Heallall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarters, Mallow, Marsh elder, Mexican poppy, Morningglory, Mustard, Nettle, Oxalis, Pennycress, Pennywort, Peppergass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak.

**BEST AVAILABLE COPY**

Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanish needle, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild patunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow charlock, Yellow daisy, Yellow rocket.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. For best results avoid: (1) application if rainfall is expected within 48 hours following treatment; (2) cutting grass 1 to 2 days before and after application. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 16 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .8 lb. per 250 sq. ft. Clean spreader thoroughly after use.

#### SPREADER SETTINGS

##### ONE PASS APPLICATION:

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

**BEST AVAILABLE COPY**

### STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

**PRODUCT DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RS 030191)

The following note will not be part of the actual label copy.

**NOTE:** Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

BEST AVAILABLE COPY