



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Russell F. Sawyer
Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866

AUG 20 2003

Dear Mr. Sawyer:

Subject: Revised Label
Riverdale Tri-Power Weed and Feed
EPA Registration Number 228-270 ✓
Your Submission Dated June 20, 2003

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- On the "Residential" labeling delete "all". The phrase "It is for use on all Residential Turf" is contradicted by the statement "DO NOT USE on Dichondra and Carpet grass".

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

If you have any questions concerning this letter, please contact Mr. James Stone at 703-305-7391.

Sincerely yours,

A handwritten signature in black ink that reads "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

2712

(Commercial Label)

ACCEPTED
with COMMENTS
in EPA Letter Dated
AUG 20 2003

RIVERDALE®
TRI-POWER®
WEED AND FEED

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

GET THE OPTICAL ADVANTAGE

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid	0.808%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.157%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)**	0.079%
INERT INGREDIENTS:	98.956%
TOTAL:	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.660%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.130%
***3,6-Dichloro-o-Anisic Acid	0.066%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-P.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

Riverdale, Tri-Power and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-270

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 2/28/03 Changed First Aid Statement per PR 2001-1, also redefined Ornamental Turf per EPA's request via Amendment.
6/20/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if absorbed through skin. Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour the washwaters on the ground; spray or drain over a large area away from well, other water sources and desirable plants.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRI-POWER® WEED AND FEED is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggartick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dogfennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod,

Ground ivy, Halbert-leaved saltbush, Hawkweed, Healall, Heartleaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarters, Mallow, Marsh elder, Mexican poppy, Morningglory, Mustard, Nettles, Oxalis, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanishneedle, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow charlock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 2 to 3 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound (or 18 pound) bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds (3.6 pounds for 18# bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader of 250 square feet and adjust to apply at the rate of 1 pound (.9 lb. for 18# bag) per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	5
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert Fertilizer claimed.

5712

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feed or foodstuffs. Do not stack more than two pallets high. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticide. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 062003)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Tri-Power® A Premium Broadleaf Herbicide

The following note will not be part of the actual label copy.

NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

ACCEPTED
with COMMENTS
In EPA Letter Dated
AUG 20 2003

6712

(Residential Label)

RIVERDALE®
TRI-POWER®
WEED AND FEED

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-270

**THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.**

Weed contacted by this product gradually die and disappear from the lawn.
Treats (or covers) 5,000 Sq. Ft. (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid	0.808%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.157%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)	0.079%

INERT INGREDIENTS:

TOTAL	98.956%
	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.660%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.130%
***3,6-Dichloro-o-Anisic Acid	0.066%

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-P

GUARANTEED ANALYSIS:¹

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur-Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum ²	X%
The urea nitrogen in this product has been coated to provide a coated slow release urea nitrogen.	

Fertilizer guaranteed by (Name of fertilizer registrant)

Riverdale, Tri-Power and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA REG. NO. 228-270

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 2/28/03 Changed First Aid Statement per PR 2001-1 and Disposal Statement per PR 2001-6, also redefined Ornamental Turf per EPA's request via Amendment.
6/20/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if absorbed through skin. Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRI-POWER® WEED AND FEED is a stable mixture of three herbicides. It is for use on all Residential Turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggartick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dogfennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbert-leaved saltbush, Hawkweed, Healall, Heartleaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter, Mallow, Marshelder, Mexican poppy, Morningglory, Mustard, Nettle, Oxalis, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanishneedle, Speedwell, Spurge, Spurweed, Stinging nettle, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow char-lock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or

8 7 12

early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound (or 18 pound) bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds (3.6 pounds for 18# bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce settings for slower speed; raise settings for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound (.9 lb. for 18# bag) per 250 square feet.

SPREADER SETTINGS
ONE PASS APPLICATION:¹

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	3
Sears Broadcast	3
Sears	5
Scotts Calibrated	5

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert Fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- A 50' x 50' lawn produces enough oxygen for a family of 4 every day.
- Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 062003)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Tri-Power® A Premium Broadleaf Herbicide

The following note will not be part of the actual label copy.

NOTE: ¹Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. ²Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation. ³(Optional) Do not apply during time periods when the maximum daily temperatures are in excess of 90°F.

(Additional Optional Fertilizer and Advertising Information)

Lawn Fertilizer
Plus Weed Control
Effectively Controls Dandelions And Other Broadleaf Weeds

(ILLUSTRATION)

- * Exclusive DeepGreen® Feeding System Keeps Lawns Green While Controlling Weeds, Whatever The Weather--Hot Or Cold, Wet Or Dry.
 - * Contains Time-Release Nitrogen For Longer Feeding And Consistently Green Lawns.
 - * Feeds Up To Three Times Longer Than Ordinary Soluble Lawn Fertilizer.
 - * 100% Satisfaction Guaranteed.
- * Toll-Free Helpline: 1-888-XXXXXX-1

10 8 12

EXPERIENCE THE (SELLER'S NAME) ADVANTAGE™

Only (Seller's Name) has the extraordinary DeepGreen® nutrient delivery system containing an exclusive polymer-coated time-release nitrogen source, assuring you the predictable, continuous feeding ordinary soluble lawn fertilizers can't deliver. So you'll enjoy the advantage of a consistently beautiful lawn week after week, whether it's hot or cold, wet or dry.

(Illustration)
(Seller's) DeepGreen® System Feeds Your Lawn Continuously When Ordinary Soluble Fertilizers Can't.

SUGGESTED SPREADER SETTINGS

One Pass Application

(Illustration) **Drop Spreaders:** First, mow lawn then apply 2 border strips along each end of the lawn. Apply fertilizer along the length of the lawn overlapping slightly between passes.

Drop Spreader

Setting

- K-Gro, Sears Drop 5
- Republic E-Z Drop 5
- Scotts Drop -- All Models 6
- Ortho Drop 5

(Illustration) **Rotary Spreaders:** First, mow lawn then apply fertilizer in a circular pattern, working from the outer edge towards the center of the lawn. Overlap slightly to insure uniform application and complete coverage.

Rotary Spreader

Setting

- Cyclone 4
- Republic E-Z Rotary 3
- Scotts Easy Green Rotary 26
- Scotts Speedy Green Rotary 4 1/2
- Ortho Rotary 6000 3

The above suggested spreader settings are approximate. In general, spreaders should be calibrated to apply 1 lb. of this fertilizer product per 275 square feet of lawn.

Be sure to close spreader openings when loading, turning or when stopped.

(SELLER'S NAME) 100% GUARANTEE OF SATISFACTION

If not totally satisfied with the results of this product, simply send proof of purchase and a brief description of your problem to (Seller name & address) for a prompt refund.

(SELLER'S NAME) HELPLINE

Questions concerning lawn care or fertilizer application? Just call toll free 1-888-XXXXXX-1

To avoid staining patios, sidewalks, swimming pools and other surfaces, remove any spilled product from these areas immediately by sweeping, vacuuming or blowing.

Made in the U.S.



TRIPLE ACTION WEED CONTROL FORMULA

X•X•X

Smart Solutions for a Beautiful Lawn™**LONGER LASTING NUTRITION**

This product is a premium grade herbicide and turf food combination which controls a wide spectrum of broadleaf weeds while providing both fast and long feeding action

DEEP GREEN COLOR

Provides nutrition (X% nitrogen, X% phosphate and X% potassium) for a deep, rich green lawn

TRIPLE-ACTION WEED CONTROL

Weeds contacted by this product gradually die and disappear from the lawn

DELIVERS MORE NUTRITION WHEN NEEDED MOST

X% nitrogen includes temperature-release nutrients that release when the lawn is growing and needs it most

(Logo)

**AMERICAN
HORTICULTURAL
SOCIETY**

NuLife proudly supports
AHS National

THIN, LOW DENSITY GRASS

Low density makes your lawn look uncared for, invites weed infestation, and makes your lawn susceptible to damage and disease. It is also an indicator of poor organic activity, which may indicate a lack of organic material.

WEAK UNDERGROWTH

Healthy roots are the key to strong, drought-resistant lawns. Root systems need phosphorous and potassium, and are aided by the organic materials and beneficial bacterial action.

POOR SUB-LAYER AND DEPLETED SOIL

Lawns need a rich sublayer of decomposing organic material to continually improve the soil and release the full range of nutrition through beneficial bacterial action. A proper layer of decomposing material helps retain moisture and provides a constantly cycling organic system.

**BEFORE
(Illustrations)****WEED INFESTATION**

Widespread weed infestation indicates a lack of sufficient plant density to naturally choke off new weeds.

POOR COLOR

Low nitrogen, low iron, acid soil or disease cause yellowing or poor color. Low usable nitrogen is the most typical cause. The lawn needs a consistent supply of nitrogen to develop and maintain good color.

**AFTER
(Illustrations)****UNIFORM DENSITY AND WEED FREE**

NuLife Weed & Feed keeps the lawn weed free in two ways. First, special formula Tri-Power® weed control has three active ingredients that work together to kill most types of broadleaf weeds. Secondly, NuLife Weed & Feed provides a steady supply of slow-release nitrogen to help the lawn become vigorous and dense, which prevents new weeds from germinating and taking hold.

DEEP GREEN COLOR

NuLife Weed & Feed provides nutrition (24% nitrogen, 6% phosphorous, 12% potassium) for a deep, rich green lawn. Nitrogen is the most important factor in lawn color. Augment with soil sweeteners if the soil becomes acidic.

THICK, LUSH, DENSE LAWN

NuLife Weed & Feed provides three sources of nitrogen: fast-release nitrogen for immediate greening; slow release for longer time between feeding, and temperature-sensitive sources that provide nitrogen when your lawn is growing and needs it most.

DURABLE ROOT SYSTEMS AND UNDERGROWTH

NuLife Weed & Feed contains phosphorous, potassium and 5% natural organic based materials to strengthen your lawn's root system, resist drought, and promote the vigor of your lawn

The NuLife "Greenhouse of Ideas" Program for Vigorous Lawns

NuLife is constantly developing and testing new ideas to make your gardening more successful and satisfying. At NuLife, we blend organic and non-organic materials to give you the best of both worlds, and to help your garden and the environment.

All NuLife formulations have been carefully developed for effectiveness and monitored for highest quality. They represent the best of the horticultural sciences and contain advanced or proprietary formulations. Good gardening! To you from NuLife.

NuLife Guarantee of Quality: NuLife guarantees your satisfaction or your money back. If you are dissatisfied for any reason, return the bag or proof of purchase to the manufacturer for a refund limited to the amount of purchase.

