EPA REGISTRATION NO. US ENVIRONMENTAL PROTECTION AGENCY 228-276 OFFICE OF PESTICIDES PROGRAMS TERM OF ISSUANCE REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 Conditional NAME OF PESTICIDE PRODUCT [X] REGISTRATION NOTICE OF PESTICIDE: FI REREGISTRATION (Under the Federal Insecticide, Fungicide, Riverdale CL/DAP Liquid Weed and Feed and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) Riverdale Chemical Company 425 West 194th Street Glenwood, IL 60425-1584 NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basic of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the libeling accepted in connection with this Registration/Reregistration is returned herewith. Registration is n no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you: required for Submit and/or cite all data registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the labeling changes listed below before you release the product for shipment: Add the phrase "EPA Registration No. 228-276." Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. ATTACHMENT IS APPLICABLE SIGNATURE OF APPROVING OFFICIAL DATE EPA Form 8570-6 (Rev. 5-76) PREVIOUS EDITION MAY BE USED UNTIL SUDDI WIR PHANAME

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

A stamped copy is enclosed for your records.

Joanne I. Miller Product Manager (23) Fungicide-Herbicide Branch Registration Division (H7505C)

Enclosures

CONCURRENCES			
EXIMOL) 47505C			
EJAHAME DE KENNY			
DATE \$ 10/1/92			
ESA Same 4000 44 (4500)		OFFICIAL FILE COPY	

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

RIVERDALE

CL/DAPTM LIQUID WEED AND FEED

Mit Jane

READY-TO-USE

ACCEPTED
with COMMENTS
to EPA Letter Dated:

20CT 1992

Under the Federal Insecticide, Pungicide, and Redentice reflect as amended, for the pesticide registers) under EP. Reg. To.

FOR USE WITH A HOSE-END SPRAYER ON GRASS AND ORNAMENTAL TURF NO MIXING

Just Connect Hose, Turn On Water, and Spray

TREATS (OR COVERS) 5,000 TO 7,500 SQ. FT. (Read directions for different grasses.)

CONTAINS MCPA, MECOPROP, DICAMBA AND A X-X-X FERTILIZER

ACTIVE INGREDIENTS: + Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*			
Isomer Specific AOAC Method, Equivalent to: *2-Methyl-4-Chlorophenoxyacetic Acid **2-(2-Methyl-4-Chlorophenoxy)propionic Acid **3,6-Dichloro-o-Anisic Acid			
GUARANTEED ANALYSIS: ° Total Nitrogen (N)	x%		
+CL/DAP IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY	Patent Pending.		

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK AND SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS 1 QT. (32 FL. OZ.)

NET WEIGHT '2,3 LBS.

• • •

EPA Reg. No. 228-ETA

EPA Est: No. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, IL 60425-1584

Revised A/O 8/27/92

1

(Left Side Panel)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear goggles, long pants, long-sleeved shirt, shoes and gloves when handling or applying this product. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow persons and pets on treated areas until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED:

Get medical attention.

IF IN EYES:

Immediately flush eyes with plenty of water. If irritation

develops get medical attention.

IF ON SKIN:

Wash with plenty of soap and water. Get medical attention.

IF INHALED:

Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsisten: with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Spray anytime during the growing season when weeds are actively growing (except as noted for S-Southern or Warm Season Grasses).

5,000 Sq. Ft. Coverage for (N-Northern or Coo¹ Season Grasses) such as Blue Grass, Fescue, and Rye grasses.

7,500 Sq. Ft. Coverage for (S-Southern or Warm Season Grasses) common Bermuda, Zoysia and Bent grasses. Some temporary injury may occur with these grasses. Applications to other grass species including bahiagrass, kikuyugrass, buffalograss, centipedegrass, carpetgrass or St. Augustinegrass should he avoided unless injury can be tolerated. In Spring, do not apply until grass has reached full green-up. In Fall, exercise care when grass growth is going into dormancy. When applying to Bent grass, height of grass should be 1-1/2" or more. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet

(optional P_2O and K_2O levels) and CL/DAP^{TM} to control listed weeds. Please contact your local extension office for questions regarding a regular fertilizer program. *Based on Inert fertilizer claimed.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been moved 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed or plant stolons for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 85°F. Shake well before using

WEEDS CONTROLLED BY CL/DAPTM LIQUID WEED AND FEED

Bindweed, Black Medic, Buckhorn, Burdock, Buttercup, Canada thistle, Carpetweed, Chickweed, Chicory, Clover, Cocklebur, Dandelion, Dock, Dog fennel, English daisy, Florida pusley, Frenchweed, Goldenrod, Ground ivy, Hawkweed, Healall, Heartloaf drymary, Henbit, Jimsonweed, Knotweed, Kochia, Lambsquarters, Mallow, Morningglory, Mustards, Oxalis, Pennywort, Pepperweed, Pigweed, Plantain, Poison ivy, Poison oak, Purslane, Ragweed, Red clover, Red sorrel, Sheep sorrel, Shepherdspurse, Smartweed, Speedwell, Spurge, Spurweed, Stinging nettle, Stitchwort, Thistles, Toadflax, Veronica, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wood sorrel, Yarrow and many other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3-4 weeks.

(HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray. Such sprayers are Ortho's Orthomatic Lawn Spray, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The following are adjustable/dial sprayers:

MANUFACTURER	<u>SETTING</u>
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	• • 4
Gilmour #486	12

SPECIAL NOTE: some adjustable sprayers do not hold 32 oz. The above settings are based on using 32 oz. per 5,000 or 7,500 sq. ft. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 5,000 sq. ft. on Cool Season Grasses or 7,500 sq. ft. for application on S-Southern or Warm Leason Grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this

Page 4

(

product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level. Point in direction away from face and body. Walk back and forth at a steady pace. Overlap slightly for uniform coverage.

(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and this product will mix automatically at the proper ratio. Walk back and forth at steady pace. Overlap slightly for uniform coverage.

(Left Side Panel)

COVERAGE MEASURING GUIDE (Marks on left for 1 qt. disposable bottle. Not to scale.)

1000 Sq. Ft. 1500

2000 Sq. Ft. 3000

3000 Sq. Ft. 4500

4000 Sq. Ft. 6000

5000 Sq. Ft. 7500 Cool Warm Season Season

. . . .

(Right Side Panel)

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near feed or foodstuffs. PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in strict accordance with directions and instructions for use. (RS 030191)

The following not will not be part of the actual label copy.

*Note: (Guaranteed (fertilizer) Analysis will vary among various commercial fertilizer manufacturers.