

NOTICE OF PESTICIDE: ☒ REGISTRATION
☐ REREISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE

Conditional

NAME OF PESTICIDE PRODUCT

Riverdale Sweet Sixteen Weed and Feed
with Dissolve

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Riverdale Chemical Company
425 West 194th Street
Glenwood, IL 60425-1584

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the labeling changes listed below before you release the product for shipment:

a. Add the phrase "EPA Registration No. 228-280."

b. Add the appropriate EPA Establishment Number to the Homeowner version of the label.

3. Submit an acceptable acute inhalation study within eight (8) months of the date of this Notice.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

☐ ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

2 of 10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosures

CONCURRENCES							
SYMBOL	H7505C						
SURNAME	D. KENNY						
DATE	10/1/92						

RIVERDALE
SWEET SIXTEEN WEED AND FEED
WITH DISSOLVE™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS
A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.70Z
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.36Z
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	0.36Z
INERT INGREDIENTS:	98.58Z
TOTAL	100.00Z

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.58Z
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.30Z
***2-(2,4-Dichlorophenoxy)propionic Acid	0.30Z

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	XZ
XZ Ammoniacal Nitrogen	
XZ Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	XZ
Soluble Potash (K ₂ O)	XZ

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated
Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum XZ

The urea nitrogen in this product has been coated to provide
XZ coated slow release urea nitrogen.

†DISSOLVE™ - Trademark of Riverdale Chemical Company Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

1 OCT 1992
NET WEIGHT 16 LBS.

EPA REG. NO. 228-EIN

Revised A/O 8/24/92

Write the Federal Government
Chemicals, and
228-750

BEST AVAILABLE COPY

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt and shoes when applying this product. Wash thoroughly with soap and water after handling. Do not allow persons or pets on treated areas until dusts have settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
IF ON SKIN: Wash skin with soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH DISSOLVE™ is a stable mixture of three herbicides. It is for use on Lawns and Ornamental Turf.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns; such as:

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bident, Bindweed, Bitterweed, Plecter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Puckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean,

5 4 10

Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .8 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE

SETTING

Central Calibrated
Cyclone Broadcast
Jackson
Sears Broadcast
Sears
Scotts Calibrated

5
3-3/4
5
2
5
4

POST APPROVED COPY

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RS 030191)

The following note will not be part of the actual label copy.

^oNOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

7 of 10

RIVERDALI
SWEET SIXTEEN WEED AND FEED
WITH DISSOLVE™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS
A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.70%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.36%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	0.36%

INERT INGREDIENTS:	98.58%
TOTAL	100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.58%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.30%
***2-(2,4-Dichlorophenoxy)propionic Acid	0.30%

ACCEPTED
with COMMENTS
in EPA Letter Dated:
1001 1992

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	XZ
XZ Ammoniacal Nitrogen	
XZ Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	XZ
Soluble Potash (K ₂ O)	XZ

Under XZ Label Instructions
Apply to the product
Signature of the Pesticide
EPA Reg. No. 228-IL-1

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated
Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum XZ

The urea nitrogen in this product has been coated to provide
XZ coated slow release urea nitrogen.

†DISSOLVE™ - Trademark of Riverdale Chemical Company Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 16 LBS.

EPA REG. NO. 228-EIN

EPA EST. NO. 228-IL-1

Revised 8/24/92



8 of 10

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, shoes and chemical resistant gloves when loading and applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. Wash thoroughly with soap and water after handling. Do not re-enter or permit workers to re-enter treated areas until dusts have settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
IF ON SKIN: Wash skin with soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH DISSOLVETM is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf (not pastures), Parks, Sod Farms, and Golf Fairways, Aprons, Tees and Roughs.

a 10

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns; such as:

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bide, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1000 square feet. Settings are based on an average

BEST AVAILABLE COPY

walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .8 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS

ONE PASS APPLICATION:°	<u>MAKE</u>	<u>SETTING</u>
	Central Calibrated	5
	Cyclone Broadcast	3-3/4
	Jackson	5
	Sears Broadcast	3
	Sears	5
	Scotts Calibrated	4

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RS 030191)

The following note will not be part of the actual label copy.

°NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

BEST AVAILABLE COPY