


228-280

07/09/2003

1/20



(Residential Label)

RIVERDALE®

SWEET SIXTEEN WEED AND FEED

WITH DISSOLVE®

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.
 Treats (or Covers) 5,000 Sq. Ft. (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.700%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	0.182%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	0.179%
INERT INGREDIENTS:	98.939%
TOTAL	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.58%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.15%
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	0.15%

‡CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-P AND DICHLORPROP-p.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	

Fertilizer guaranteed by (Name of fertilizer registrant)

Riverdale, Dissolve and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company Patent No. 5,266,553

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-280 NET WEIGHT LBS. EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 2/14/03 Changed First Aid Statement per PR 2001-1 and Disposal Statement per PR 2001-6. Also deleted California use limitation via Amendment.
 6/13/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

(Back Panel)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
- IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH DISSOLVE® is a stable mixture of three herbicides. It is for use on all Residential Turf.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bidden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw,

Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Nutgrass, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 pounds for 16 lb. bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .7 pound (.8 lb. for 16 lb. bag) per 250 square feet.

4/20

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed, or other pesticides.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 061303)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

5/20

The following note will not be part of the actual label copy.

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

- Editor's notes:
1. Subregistrant Warranty Statements previously accepted by U.S. EPA for use on their own registered products are not included on this label.
 2. Subregistrant's trademark name acknowledgments are not included on this label.

Additional Fertilizer and Marketing Claims

For Thick, Green, Weed-Free Lawns

Extended Greening From Controlled Release Nitrogen

For best year round results, this product should be used in conjunction with (product name) Lawn Food.

Refer to the table and picture below to determine the size of your lawn. Refer to "Path A: when using a broadcast spreader and "Path B" when using a drop spreader.

_____	Length (Ft.)	Illustration
X _____	Width (Ft)	
= _____	Total Sq. Ft.	of
- _____	Non-Lawn Areas	
= _____	Total Sq. Ft.	Property
	Coverage Needed	

Illustration

of

Path A

Illustration

of

Path B

Contains Dissolve® A Premium Broadleaf Herbicide

Additional Marketing Statements

- Feeds desirable grasses
- Controls Dandelions and more than 250 other Broadleaf Weeds
- Controlled release Nitrogen feeds grass longer

Pledge of Satisfaction

We have made every effort in the formulation of this product to insure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn and turf specialists will make every effort to assure your success.

For best results when used on your lawn, this product should be used in conjunction with (product name) Lawn Food.

If, for any reason, you are not completely satisfied with this product, your money will be refunded. Send proof of purchase to the address listed with explanation of your problem.

Additional Marketing and Fertilizer Statements

CONTROLS OVER 60 TYPES OF UGLY WEEDS COMPLETELY - ROOTS AND ALL!

- Feeds your lawn with Superturf to produce a deep green, vigorous, competitive turf.
- Apply anytime during growing season Spring - Summer - Fall.
- Another Quality Product from (Company Name)!

4 - APPLICATION ANNUAL LAWN CARE PROGRAM			
(ILLUSTRATIONS)			
Apply 1st in my Annual Lawn Care Program in the Spring. Controls Crabgrass while feeding your lawn.	Apply 2nd in my Annual Lawn Care Program Spring through Fall. Controls Broadleaf Weeds when they appear while feeding your lawn.	Apply 3rd in my Annual Lawn Care Program in Summer. Controls insects while feeding your lawn.	Apply 4th in my Annual Lawn Care Program in Fall. Provides nutrients for a greener, thicker lawn.

ADDITIONAL FERTILIZER AND MARKETING HYPE

SUGGESTED SPREADING APPLICATIONS

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. This bag covers approximately 5,000 square feet (100 ft. x 50 ft.). Close spreader hopper and adjust the spreader setting using the Spreader Settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Close the spreader hopper whenever spreader stops or turns. For one pass application rate.

ILLUSTRATION	<p>Broadcast Spreader Settings</p> <p>For best results, apply in a circular path starting on the outside and working toward the center (see Broadcast Spreader Path diagram). A little overlap of product will insure complete coverage.</p>	ILLUSTRATION
--------------	---	--------------

ILLUSTRATION	<p>Drop Spread Settings</p> <p>First apply 2 border strips along the edge of each end of the lawn (see Drop Spreader Path diagram). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.</p>	ILLUSTRATION
--------------	---	--------------

Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

Controls over 250 broadleaf weeds, including: Dandelion, Plaintain, Chickweed, Knotweed, Henbit and Spurge.

Contains controlled-release nitrogen for extended feeding.

7/20

Nitrogen	Phosphorus	Potassium
<p>Essential for plant growth and a thick green lawn. (OR) Nitrogen is essential for giving your lawn lush green color and dense foliage. (OR) For a Dense Green Lawn</p>	<p>To promote root growth and grass multiplication. (OR) Phosphorus helps your grass develop and maintain vigorous root growth. (OR) Improves Root Growth</p>	<p>To help grass resist the elements and stay strong. (OR) (Also referred to as Potash). This nutrient is vital for vigorous growth while helping your lawn cope with drought and other extreme weather conditions. (OR) Vital for growth & drought resistance</p>

8/20

(SELLER'S) ANNUAL LAWN PROGRAM

For best results this product should be used in conjunction with (Seller's) Crabgrass Preventer Plus Lawn Food, (Seller's) Lawn Food or (Seller's) Fall Lawn Food.

HOW IT WORKS:

Start anytime! No guesswork, just follow these steps for a thick, green, weed-free lawn. It's easy...simply apply the FRANK'S™ product that corresponds to the season.

	•PREVENTS Crabgrass & other unsightly weeds	apply early spring	apply mid-late spring	•CONTROLS Dandelions & more than 140 other weeds
	•ENCOURAGES quick spring greenup	SELLERS CRABGRASS Preventer Plus Lawn Food	SELLERS WEED & FEED	•THICKENS & greens your lawn with extended feeding Nitrogen
(A)	•GREENS your lawn with extended feeding Nitrogen	apply fall	(A) apply summer	•GREENS with Iron & Sulfur (A)
	•THICKENS fall lawns & encourages quick spring greenup	SELLERS FALL LAWN FOOD	SELLERS LAWN FOOD	•THICKENS & promotes a lush green lawn
	•NOURISHES your lawn with extended feeding Nitrogen			•NOURISHES your lawn with extended feeding Nitrogen
	•GREENS with Iron & Sulfur			•GREENS with Iron & Sulfur

(A) BAG ILLUSTRATIONS

DO NOT HAND APPLY.

SPECIAL CARE NOTICE: This product contains the micronutrient Iron to help prevent Iron deficiency. Iron may stain concrete surfaces such as sidewalks, patios and swimming pools.

Products should not be applied on water dampened concrete and should be removed promptly after application by sweeping or vacuuming.

CONTROLS ALL COMMON WEED TYPES

Where beautiful gardens begin™

- Controls Dandelions & more than 250 other weeds
- Thickens & greens your lawn
- Greens with Iron & Sulfur

Contains Poly-S® controlled release nitrogen for extended greening (For best results, use with other lawn care products in the [Seller's Name] Fertilizer Program.)

Contains the 3 way power of Dissolve® for Broadleaf, Creeping and Woody weeds.

9/20

Controls over 110 common lawn weeds while feeding your lawn with a premium blend of fertilizer nutrients that provide quick and extended greening.

BROADCAST SPREADER SETTINGS

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

DROP SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

(Attach Sam Walton info)

Sam Walton believed in the ability of American workers to produce the finest products in the world.

That's why we're searching the USA to create something special. Innovative products, made to our own higher standards, that we're proud to call Sam's American Choice™. We believe these products, available at Wal-Mart and Wal-Mart Supercenters, offer better value than the leading national brands at the always low Wal-Mart prices. We think you'll agree, or we'll refund your money, no questions asked!

"Better products at the always low Wal-Mart prices..."

Always

Sam m. Walton

We guarantee it.

Sam Walton

COMPETITIVE EDGE SCHOLARSHIP FUND

In the next 10 years, the United States will have the jobs Americans need, but Americans may not have the skills these jobs demand. To have a competitive edge in the future, American must have a well-educated, trained work force. To help achieve this goal, Wal-Mart and our vendor partner will each donate 1% of the cost of this product to our Competitive Edge Scholarship Program.

GreenThumb Fertilizer Program

The following represents the suggested application timing of the products listed in the GreenThumb Program. Abnormal climatic conditions can affect appropriate timing.

U.S. Climate Zones:



Bluegrass, Ryegrass, Fine & Tall Fescues

Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagrass, Zoysiagrass

Bermudagrass, Zoysiagrass, Centipedegrass, Bahiagrass, St. Augustinegrass

GreenThumb Premium Crabgrass Preventer + Lawn Fertilizer

- Promotes a lush, green lawn while controlling crabgrass and most other annual weed grasses.
- Contains the pre-emergent herbicide, Team*

- Apply mid March - April
- Apply early March - mid April
- Apply mid February - late March

GreenThumb Premium Weed & Feed Lawn Fertilizer

- Controls over 140 broadleaf weeds, including Dandelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.
- Contains the premium broadleaf herbicide Dissolve®

- Apply mid April - late June
- Apply late March - May
- Apply mid March - mid May

GreenThumb Premium Lawn Fertilizer

- Specialty formulated to promote even, lush growth of your grass.
- Contains slow release nitrogen to extend feeding of the lawn.

- Apply June - August
- Apply mid May - early September
- Apply mid April - September

GreenThumb Premium Summerizer Lawn Fertilizer

- Special non-burning formula, when used as directed, for light summer application.
- Contains slow release nitrogen to extend feeding of the lawn.

- Apply June - August
- Apply mid May - early September
- Apply mid April - September

GreenThumb Premium Winterizer Lawn Fertilizer

- Formulated for fall application to prepare your turf for winter.
- Contains slow release nitrogen to extend feeding of the lawn.

- Apply September - mid October
- Apply September - late October
- Apply September - mid November

Team is a reg. TM of DowElanco; Dissolve - Riverdale Chemical Co., Patent No. 6,256,653

11/20

ALTERNATE ARRANGEMENT OF LISTED WEEDS

Alder	Dogbane	Mustard, Tumble
Artichoke	Dogfennel	Mustard, White
Aster	Elderberry	Mustard, Wild
Aster, Wild	Fiddleneck	Nettle
Bedstraw	Fieldcress, Austrian	Nettle, Bull
Bedstraw, Pineywoods	Filaree, Redstem	Nettle, Burning
Beggartick	Flax, False	Nettle, Dead
Betony, Florida	Flea bane (daisy)	Nettle, Stinging
Biden	Fleabane, Hairy	Nettle, Tall
Bindweed	Fleabane, Rough	Nutgrass
Bindweed, Field	Flixweed	Oak, Poison
Bindweed, Hedge	Florida pusley	Onion, Wild
Bitter wintercress	Four-O'-Clock, Wild	Oxalis
Bittercress, Hairy	Frenchweed	Pansy, Field
Bitterweed	Galinsoga	Pantain, Blackseed
Black-eyed Susan	Galinsoga, Smallflower	Parsley-piert
Bloodflower, Milkweed	Garlic, Wild	Parsnip
Boxelder	Geranium, Carolina	Parsnip, Wild
Brassbuttons	Geranium, Wild	Pearlwort
Bristly Oxtongue	Goathead	Pennycress
Broomweed	Goatsbeard	Pennywort
Buckhorn	Goldenrod	Peppergrass
Buckwheat, Wild	Ground-Ivy	Pepperweed
Bullthistle, Wavyleaf	Groundsel	Pepperweed, Virginia
Bur ragweed	Gumweed	Pigweed
Burdock	Hawkweed	Pigweed, Prostrate
Burweed	Hawkweed, Orange	Pigweed, Redroot
Buttercup	Healall	Pigweed, Russian
Buttercup, Bulbous	Heartleaf drymary	Pigweed, Smooth
Carpetweed	Heathaster	Pigweed, Tumble
Carrot, Wild	Hemlock, Poison	Scarlet Pimpernel
Catchweed Bedstraw	Hemp	Plantain
Catnip	Henbit	Plantain, Blackseed
Catsear	Hoary cress	Plantain, Bracted
Catsear, Spotted	Honeysuckle	Plantain, Broadleaf
Chamomile, Corn	Horsenettle	Plantain, Buckhorn
Chickweed	Horsetail	Plantain, Hoary
Chickweed, Common	Ironweed	Plantain, Narrowleaf
Chickweed, Mousear	Ivy, Poison	Plantain, Slender
Chicory	Jewelweed	Plantain, Woolly
Cinquefoil	Jimsonweed	Pokeweed
Cinquefoil, Rough	Knapweed, Spotted	Poorjoe
Clematis, Western	Knawel	Povertyweed
Clover	Knotweed	Prairie
Clover, Annual Yellow Sweet	Knotweed, Prostrate	Prickly Sida
Clover, Crimson	Kochia	Primrose
Clover, Hop	Lambsquarter	Puncture vine
Clover, Red	Lespedeza	Purslane
Clover, Strawberry	Lettuce, Blue	Radish, Wild
Clover, Sweet	Lettuce, Prickly	Ragweed
Clover, White	Lettuce, Wild	Rape, Wild
Cockle	Locoweed	Rose, Southern Wild
Cocklebur	Lupine	Rush
Cocklebur, Spiny	Mallow	Salsify, Western
Coffeebean	Mallow, Indiana	Shepherdspurse
Coffeeweed	Mallow, Venice	Smartweed
Coreopsis, Plains	Marigold, Roundleafed	Sneezeweed
Creeping jenny	Marigold, Wild	Sorrel, Red
Croton	Marshelder	Sorrel, Sheep
Croton, Woolly	Matchweed	Sowthistle
Cudweed	Medic, Black	Sowthistle, Common
Curly indigo	Mexicanweed	Spanishneedle
Daisy, English	Morningglory	Spatterdock
Daisy, Oxeye	Morningglory, Woolly	Speedwell
Dandelion	Mugwort	Spiny
Dandelion, Fall	Mullein, Common	Spiny Amaranth
Dandelion, False	Mustard	Spurge
Dock	Mustard, Black	Spurge, Prostrate
Dock, Broadleaf	Mustard, Garlic	Spurge, Spotted
Dock, Curly	Mustard, Hedge	Spurweed
Dock, Smooth	Mustard, Tansy	St. Johnswort

12/20

Stinkweed
Stitchwort
Strawberry, Wild
Sumac
Sunflower
Sunflower, False
Sweet Potato, Wild
Tansy Ragwort
Tanweed
Tarweed
Thistle
Thistle, Blessed
Thistle, Bull
Thistle, Canada
Thistle, Musk

Thistle, Russian
Thistle, Scotch
Toadflax
Trailing Crownvetch
Trefoil, Tick
Tumbleweed
Velvet leaf
Veronica
Vervain
Vervain, Blue
Vervain, Hoary
Vervain, Prostrate
Vervain, Tall
Vetch
Vetch, Bird

Vetch, Milk
Vetch, Narrowleaf
Vetch, Wild
Virginia Buttonweed
Virginia Creeper
Willow
Witchweed
Woodsorrel
Wormseed
Yarrow
Yellow rocket
Yellowflower Pepperweed
And other broadleaf
weeds

13/20

(Commercial Label)

RIVERDALE®

SWEET SIXTEEN WEED AND FEED

WITH DISSOLVE®

ACCEPTED

JUL 9 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 228-280

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.
Treats (or Covers) 5,000 Sq. Ft. (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.700%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.182%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†	0.179%
INERT INGREDIENTS:	98.939%
TOTAL	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.58%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.15%
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	0.15%

†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

Riverdale, Dissolve and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company Patent No. 5,266,553

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-280

NET WEIGHT LBS.

ERA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 2/14/03 Changed First Aid Statement per PR 2001-1 via Amendment.
6/13/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

(Back Panel)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks, shoes and gloves when loading and applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
- IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

SWEET SIXTEEN WEED AND FEED WITH DISSOLVE® is a stable mixture of three herbicides. It is for use on Ornamental Turf such as Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St.

15/20

Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Nutgrass, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennygrass, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

16/20

HOW TO APPLY: This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 pounds for 16 lb. bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .7 pound (.8 lb. for 16 lb. bag) per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:°

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

17/20

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 061303)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS:

The following note will not be part of the actual label copy.

NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Additional Fertilizer and Marketing Claims

For Thick, Green, Weed-Free Lawns

Extended Greening From Controlled Release Nitrogen

For best year round results, this product should be used in conjunction with (product name) Lawn Food.

Refer to the table and picture below to determine the size of your lawn. Refer to "Path A: when using a broadcast spreader and "Path B" when using a drop spreader.

_____	Length (Ft.)	Illustration
X _____	Width (Ft)	
= _____	Total Sq. Ft.	of
- _____	Non-Lawn Areas	
= _____	Total Sq. Ft.	Property
	Coverage Needed	

Illustration	Illustration
of	of
Path A	Path B

Contains Dissolve® A Premium Broadleaf Herbicide

Additional Marketing Statements

- Feeds desirable grasses
- Controls Dandelions and more than 250 other Broadleaf Weeds
- Controlled release Nitrogen feeds grass longer

18/20

Pledge of Satisfaction

We have made every effort in the formulation of this product to insure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn and turf specialists will make every effort to assure your success.

For best results when used on your lawn, this product should be used in conjunction with (product name) Lawn Food.

If, for any reason, you are not completely satisfied with this product, your money will be refunded. Send proof of purchase to the address listed with explanation of your problem.

ADDITIONAL FERTILIZER AND MARKETING HYPE

SUGGESTED SPREADING APPLICATIONS

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. This bag covers approximately 5,000 square feet (100 ft. x 50 ft.). Close spreader hopper and adjust the spreader setting using the Spreader Settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Close the spreader hopper whenever spreader stops or turns. For one pass application rate.

ILLUSTRATION	<p>Broadcast Spreader Settings</p> <p>For best results, apply in a circular path starting on the outside and working toward the center (see Broadcast Spreader Path diagram). A little overlap of product will insure complete coverage.</p>	ILLUSTRATION
--------------	---	--------------

ILLUSTRATION	<p>Drop Spread Settings</p> <p>First apply 2 border strips along the edge of each end of the lawn (see Drop Spreader Path diagram). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.</p>	ILLUSTRATION
--------------	---	--------------

Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

Controls over 250 broadleaf weeds, including: Dandelion, Plaintain, Chickweed, Knotweed, Henbit and Spurge.

Contains controlled-release nitrogen for extended feeding.

19/20

<p>Nitrogen</p> <p>Essential for plant growth and a thick green lawn.</p> <p>(OR)</p> <p>NITROGEN</p> <p>For a dense green lawn</p>	<p>Phosphorus</p> <p>To promote root growth and grass multiplication.</p> <p>(OR)</p> <p>PHOSPHATE</p> <p>Improves root growth</p>	<p>Potassium</p> <p>To help grass resist the elements and stay strong.</p> <p>(OR)</p> <p>POTASH</p> <p>Vital for growth & drought resistance</p>
---	--	---

FOR THE BEST LAWN IN TOWN

ALTERNATE ARRANGEMENT OF LISTED WEEDS

- | | | |
|---|---|---|
| <ul style="list-style-type: none"> Alder Artichoke Aster Aster, Wild Bedstraw Bedstraw, Pineywoods Beggartick Betony, Florida Biden Bindweed Bindweed, Field Bindweed, Hedge Bitter wintercress Bittercress, Hairy Bitterweed Black-eyed Susan Bloodflower, Milkweed Boxelder Brassbuttons Bristly Oxtongue Broomweed Buckhorn Buckwheat, Wild Bullthistle, Wavyleaf Bur ragweed Burdock Burweed Buttercup Buttercup, Bulbous Carpetweed Carrot, Wild Catchweed Bedstraw Catnip Catsear Catsear, Spotted Chamomile, Corn Chickweed Chickweed, Common Chickweed, Mousear Chicory Cinquefoil Cinquefoil, Rough Clematis, Western Clover Clover, Annual Yellow Sweet Clover, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White | <ul style="list-style-type: none"> Cockle Cocklebur Cocklebur, Spiny Coffeebean Coffeeweed Coreopsis, Plains Creeping jenny Croton Croton, Woolly Cudweed Curly indigo Daisy, English Daisy, Oxeye Dandelion Dandelion, Fall Dandelion, False Dock Dock, Broadleaf Dock, Curly Dock, Smooth Dogbane Dogfennel Elderberry Fiddleneck Fieldcress, Austrian Filaree, Redstem Flax, False Flea bane (daisy) Fleabane, Hairy Fleabane, Rough Flixweed Florida pusley Four-O'-Clock, Wild Frenchweed Galinsoga Galinsoga, Smallflower Garlic, Wild Geranium, Carolina Geranium, Wild Goathead Goatsbeard Goldenrod Ground-Ivy Groundsel Gumweed Hawkweed Hawkweed, Orange Healall Heartleaf drymary Heathaster Hemlock, Poison | <ul style="list-style-type: none"> Hemp Henbit Hoary cress Honeysuckle Horsenettle Horsetail Ironweed Ivy, Poison Jewelweed Jimsonweed Knapweed, Spotted Knawel Knotweed Knotweed, Prostrate Kochia Lambsquarter Lespedeza Lettuce, Blue Lettuce, Prickly Lettuce, Wild Locoweed Lupine Mallow Mallow, Indiana Mallow, Venice Marigold, Roundleafed Marigold, Wild Marshelder Matchweed Medic, Black Mexicanweed Morningglory Morningglory, Woolly Mugwort Mullein, Common Mustard Mustard, Black Mustard, Garlic Mustard, Hedge Mustard, Tansy Mustard, Tumble Mustard, White Mustard, Wild Nettle Nettle, Bull Nettle, Burning Nettle, Dead Nettle, Stinging Nettle, Tall Nutgrass Oak, Poison |
|---|---|---|

Onion, Wild
 Oxalis
 Pansy, Field
 Pantain, Blackseed
 Parsley-pierr
 Parsnip
 Parsnip, Wild
 Pearlwort
 Pennycress
 Pennywort
 Peppergrass
 Pepperweed
 Pepperweed, Virginia
 Pigweed
 Pigweed, Prostrate
 Pigweed, Redroot
 Pigweed, Russian
 Pigweed, Smooth
 Pigweed, Tumble
 Scarlet Pimpernel
 Plantain
 Plantain, Blackseed
 Plantain, Bracted
 Plantain, Broadleaf
 Plantain, Buckhorn
 Plantain, Hoary
 Plantain, Narrowleaf
 Plantain, Slender
 Plantain, Woolly
 Pokeweed
 Poorjoe
 Povertyweed
 Prairie
 Prickly Sida
 Primrose
 Puncture vine

Purslane
 Radish, Wild
 Ragweed
 Rape, Wild
 Rose, Southern Wild
 Rush
 Salsify, Western
 Shepherdspurse
 Smartweed
 Sneezeweed
 Sorrel, Red
 Sorrel, Sheep
 Sowthistle
 Sowthistle, Common
 Spanishneedle
 Spatterdock
 Speedwell
 Spiny
 Spiny Amaranth
 Spurge
 Spurge, Prostrate
 Spurge, Spotted
 Spurweed
 St. Johnswort
 Stinkweed
 Stitchwort
 Strawberry, Wild
 Sumac
 Sunflower
 Sunflower, False
 Sweet Potato, Wild
 Tansy Ragwort
 Tanweed
 Tarweed
 Thistle
 Thistle, Blessed

Thistle, Bull
 Thistle, Canada
 Thistle, Musk
 Thistle, Russian
 Thistle, Scotch
 Toadflax
 Trailing Crownvetch
 Trefoil, Tick
 Tumbleweed
 Velvet leaf
 Veronica
 Vervain
 Vervain, Blue
 Vervain, Hoary
 Vervain, Prostrate
 Vervain, Tall
 Vetch
 Vetch, Bird
 Vetch, Milk
 Vetch, Narrowleaf
 Vetch, Wild
 Virginia Buttonweed
 Virginia Creeper
 Willow
 Witchweed
 Woodsorrel
 Wormseed
 Yarrow
 Yellow rocket
 Yellowflower
 Pepperweed
 And other broadleaf
 weeds