



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 08, 2026

SENT BY EMAIL

Ann Kumari
annu.kumari@nufarm.com
NUFARM AMERICAS, INC.

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Add an optional marketing statement to the label
Product Name: SWEET SIXTEEN WEED AND FEED WITH DISSOLVE
Admin Number: 228-280
EPA Receipt Date: 12/08/2025
Action Case Number: 00678151

Dear Ann Kumari:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Elizabeth Andrews via email at andrews.elizabeth@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Elizabeth Andrews 'for'".

Mindy Ondish, PM 23
HB, RD
Office of Pesticide Programs

Residential

SWEET SIXTEEN WEED AND FEED WITH DISSOLVE®

NOTIFICATION

228-280

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

01/08/2026

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or Covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

| | |
|----------------------------------------------------------------------------|-----------------|
| Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* | 0.700% |
| Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid**‡ | 0.182% |
| Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***‡ | 0.179% |
| OTHER INGREDIENTS: | 98.939% |
| TOTAL: | 100.000% |

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

| | |
|--------------------------------------------------------------------|---------------------|
| *2,4-Dichlorophenoxyacetic Acid..... | 0.58%, 0.0058 ae/lb |
| **(+)-R-2-(2-Methyl-4-Chlorophenoxy) Propionic Acid | 0.15%, 0.0015 ae/lb |
| ***(+)-R-2-(2,4-Dichlorophenoxy) Propionic Acid | 0.15%, 0.0015 ae/lb |
| †CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p. | |

GUARANTEED ANALYSIS:

| | |
|-----------------------------------------------------------|----|
| Total Nitrogen (N) | X% |
| X% Ammoniacal Nitrogen | |
| X% Urea Nitrogen | |
| Available Phosphate (P ₂ O ₅)..... | X% |
| Soluble Potash (K ₂ O)..... | X% |

[Primary Plant Foods] derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum..... X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

Alternative Text: [SEE] [REFER TO] [INSIDE] [LABEL BOOKLET] [BACK PANEL] [FOR] [FIRST AID] [AND] [ADDITIONAL] [PRECAUTIONARY STATEMENTS] [AND] [DIRECTIONS FOR USE]

For Medical Emergencies , Call (877) 325-1840

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

[Note to reviewer: Any text found in brackets "[]" is optional on container label.]

EPA REG. NO. 228-280
EPA EST. NO. 228-IL-1

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



NET WEIGHT: LBS. (Kg)
000228-00280. 20250307

Residential

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

| FIRST AID | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15 to 20 minutes.Call a poison control center or doctor for treatment advice. |
| IF INHALED | <ul style="list-style-type: none">Move person to fresh air.If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth- to-mouth, if possible.Call a poison control center or doctor for further treatment advice. |
| IF IN EYES | <ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information. | |

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

NON-TARGET ORGANISM DVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators. In areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

GROUNDWATER ADVISORY

2,4-DP-p is known to leach through soil into groundwater under certain conditions as a result of label use.

This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow

Residential

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

PRODUCT USE INFORMATION

A minimum of 30 days between applications is required.

This product is a stable mixture of three herbicides. It is for use on all residential turf.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until dusts have settled.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

Residential

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

| | | | |
|-----------------------------|------------------------|-----------------------|----------------------|
| Alder | Coreopsis, Plains | Knapweed, Spotted | Pigweed, Redroot |
| Artichoke | Creeping Jenny | Knawel | Pigweed, Russian |
| Aster | Croton | Knotweed | Pigweed, Smooth |
| Aster, Wild | Croton, Wooly | Knotweed, Prostrate | Pigweed, Tumble |
| Bedstraw | Cudweed | Kochia | Pimpernel, Scarlet |
| Bedstraw, Pineywoods | Curly Indigo | Lambsquarter | Plantain |
| Beggartick | Daisy, English | Lespedeza | Plantain, Blackseed |
| Betony, Florida | Daisy, Oxeye | Lettuce, Blue | Plantain, Bracted |
| Biden | Dandelion | Lettuce, Prickly | Plantain, Broadleaf |
| Bindweed | Dandelion, Fall | Lettuce, Wild | Plantain, Buckhorn |
| Bindweed, Field | Dandelion, False | Locoweed | Plantain, Hoary |
| Bindweed, Hedge | Dock | Lupine | Plantain, Narrowleaf |
| Bitter Wintercress | Dock, Broadleaf | Mallow | Plantain, Slender |
| Bittercress, Hairy | Dock, Curly | Mallow, Indiana | Plantain, Wooly |
| Bitterweed | Dock, Smooth | Mallow, Venice | Pokeweed |
| Black-Eyed Susan | Dogbane | Marigold, Roundleafed | Poorjoe |
| Bloodflower, Milkweed | Dogfennel | Marigold, Wild | Povertyweed |
| Box Elder | Elderberry | Marshelder | Prickly Sida |
| Brassbuttons | Fiddleneck | Matchweed | Primrose |
| Bristly Oxtongue | Fieldcress, Austrian | Medic, Black | Puncturevine |
| Broomweed | Filaree, Redstem | Mexicanweed | Purslane |
| Buckhorn | Flax, False | Morningglory | Radish, Wild |
| Buckwheat, Wild | Flea Bane (Daisy) | Moringlory, Wooly | Ragweed |
| Bullthistle, Wavyleaf | Fleabane, Hairy | Mugwort | Rape, Wild |
| Bur Ragweed | Fleabane, Rough | Mullein, Common | Rose, Southern Wild |
| Burdock | Flixweed | Mustard | Rush |
| Burweed | Florida Pusley | Mustard, Black | Salsify, Western |
| Buttercup | Four-O'Clock, Wild | Mustard, Garlic | Shepherdspurse |
| Buttercup, Bulbous | Frenchweed | Mustard, Hedge | Smartweed |
| Carpetweed | Galinsoga | Mustard, Tansy | Sneezeweed |
| Carrot, Wild | Galinsoga, Smallflower | Mustard, Tumble | Sorrel, Red |
| Catchweed Bedstraw | Garlic, Wild | Mustard, White | Sorrel, Sheep |
| Catnip | Geranium, Carolina | Mustard, Wild | Sowthistle, |
| Catsear | Geranium, Wild | Nettle | Sowthistle, Common |
| Catsear, Spotted | Goathead | Nettle, Bull | Spanishneedle |
| Chamomile, Corn | Goatsbeard | Nettle, Burning | Spatterdock |
| Chickweed | Goldenrod | Nettle, Dead | Speedwell |
| Chickweed, Common | Ground Ivy | Nettle, Stinging | Spiny Amaranth |
| Chickweed, Mouseear | Groundsel | Nettle, Tall | Spurge |
| Chicory | Gumweed | Nutgrass | Spurge, Prostrate |
| Cinquefoil | Hawkweed | Orange hawkweed | Spurge, Spotted |
| Cinquefoil, Rough | Hawkweed, Orange | Oak, Poison | Spurweed |
| Clematis, Western | Healall | Onion, Wild | St. Johnswort |
| Clover | Heartleaf Drymary | Oxalis | Stinkweed |
| Clover, Annual Yellow Sweet | Heathaster | Pansy, Field | Stitchwort |
| Clover, Crimson | Hemlock, Poison | Parsley-Piert | Strawberry, Wild |
| Clover, Hop | Hemp | Parsnip | Sumac |
| Clover, Red | Henbit | Parsnip, Wild | Sunflower |
| Clover, Strawberry | Hoary Cress | Pearlwort | Sunflower, False |
| Clover, Sweet | Honeysuckle | Pennycress | Sweet Potato, Wild |
| Clover, White | Horsenettle | Pennywort | Tansy Ragwort |
| Cockle | Horsetail | Peppergrass | Tanweed |
| Cocklebur | Ironweed | Pepperweed | Tarweed |
| Cocklebur, Spiny | Ivy, Poison | Pepperweed, Virginia | Thistle |
| Coffeebean | Jewelweed | Pigweed | |
| Coffeeweed | Jimsonweed | Pigweed, Prostrate | |

(continued)

Residential

| | | | |
|---------------------|--------------------|----------------------|-------------------------|
| Thistle, Blessed | Trefoil, Tick | Vervain, Tall | Willow |
| Thistle, Bull | Tumbleweed | Vetch | Witchweed |
| Thistle, Canada | Velvet Leaf | Vetch, Bird | Wood sorrel |
| Thistle, Musk | Veronica | Vetch, Milk | Wormseed |
| Thistle, Russian | Vervain | Vetch, Narrowleaf | Yarrow |
| Thistle, Scotch | Vervain, Blue | Vetch, Wild Virginia | Yellow Rocket |
| Toadflax | Vervain, Hoary | Buttonweed Virginia | Yellowflower Pepperweed |
| Trailing Crownvetch | Vervain, Prostrate | Creeper | |

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 pounds for 16 pound bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.7 pound (0.8 pound for 16 pound bag) per 250 square feet.

| SPREADER SETTINGS ONE PASS APPLICATION: | |
|--------------------------------------------|---------|
| MAKE | SETTING |
| Central Calibrated | 5 |
| Cyclone Broadcast | 3-3/4 |
| Jackson | 5 |
| Sears Broadcast | 3 |
| Sears | 5 |
| Scotts Calibrated | 4 |

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Use Precautions for Turf, ornamental (golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas)

Postemergence:

Limited to 2 applications per year.

Maximum of 250 pounds of product /acre per application.

The maximum seasonal rate is 500 pounds of product/acre, excluding spot treatments.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert Fertilizer claimed.

Residential

| | |
|------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Environmental Benefits (Grasscycling Logo) | <p>Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:</p> <p><i>Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.</i></p> <ul style="list-style-type: none">✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day✓ Reduces Glare & Reduces Noise✓ Traps Dust, Smoke Particles & Other Pollutants✓ Purifies Water & Modifies Temperature✓ Improves Soil Condition & Controls Erosion |
|------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV.....)

Dissolve and The Optical Advantage are registered trademarks of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.

Residential

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

Apply with a broadcast or drop spreader.

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Editor's notes: 1. Subregistrant Warranty Statements previously accepted by U.S. EPA for use on their own registered products are not included on this label.

2. Subregistrant's trademark name acknowledgements are not included on this label.

Additional Fertilizer and Marketing Claims

FEEDS UP TO 2 MONTHS [2 calendar logo illustration]

STOP! READ THE ENTIRE LABEL FIRST, OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

For Thick, Green, Weed-Free Lawns

Extended Greening From Controlled Release Nitrogen

For best year round results, this product should be used in conjunction with (product name) Lawn Food.

Refer to the table and picture below to determine the size of your lawn. Refer to "Path A" when using a broadcast spreader and "Path B" when using a drop spreader.

Kills weeds* & promotes hardy, lush lawns

*when used as directed

| | | |
|---|--------------------------------------|--------------------------------|
| - | Length (Feet) | |
| X | Width (Feet) | |
| = | Total Square Feet | Illustration of Property |
| - | Non-Lawn Areas | |
| = | Total Square Feet Coverage Needed | |
| | | Illustration of Path A |
| | | Illustration of Path B |

Contains Dissolve® A Premium Broadleaf Herbicide

Residential

Additional Optional Marketing Statements

- Feeds desirable grasses
- Controls Dandelions and more than 250 other Broadleaf Weeds
- Controlled release Nitrogen feeds grass longer
- Kills dandelions and over 140 other weeds to give you a weed-free lawn in the Spring!
- Kills dandelions and other broadleaf weeds* and promotes a hardy, lush lawn
 - *when used as directed
- [6500] sq. ft coverage
 - { Note to reviewer: sq.ft coverage is based upon size of the package. Bracketed numbers can change as package size changes.}
- [XX[lbs.] [pounds] [treats] [covers] up to [XXXX[[square feet] [sq. ft.] [ft²]
- [For Florida, this [xx] pound bag should be used on [XXXX] square feet.]
- Kills Dandelions and Other Broadleaf Weeds and Promotes a Hardy Lush Lawn
- Kills/Controls most common broadleaf weeds
- 2 in 1 formula [graphic] kills weeds and greens/feeds your lawn
- Promotes lush green growth in lawns & control broadleaf weeds
- [Insert Company Name / Brand Name] Weed & Feed; This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns.
- Controls [Kills] clover, dandelion, plantain, chickweed and [other broadleaf weeds] [others as listed]
- Kills tough lawn weeds
- Kills [Over] [More Than] [XXX] [Listed] Weeds [as Listed]
- Kills dandelion and many other broadleaf weeds.
- Kills Weeds While Feeding Your Lawn
- Wipes out tough weeds such as [chickweed, clover, henbit, and/or dandelion]
- Wipes out Weeds While Feeding Your Lawn
- Eliminates even the toughest weeds!
- Eliminates broadleaf weeds in lawns
- Provides excellent turf response plus outstanding broadleaf weed control.

Best results are obtained [Apply] when weeds are growing actively in spring and early fall.

Formatted: Font: 9 pt

Residential

PLEDGE OF SATISFACTION

We have made every effort in the formulation of this product to insure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn and turf specialists will make every effort to assure your success.

For best results when used on our lawn, this product should be used in conjunction with (product name) Lawn Food.

If, for any reason, you are not completely satisfied with this product, your money will be refunded. Send proof of purchase to the address listed with explanation of your problem.

Additional Marketing and Fertilizer Statements

CONTROLS OVER 60 TYPES OF UGLY WEEDS COMPLETELY - ROOTS AND ALL!

- Feeds your lawn with Superturf to produce a deep green, vigorous, competitive turf.
- Apply anytime during growing season spring - summer - fall.
- Another Quality Product from (Company Name)!
- Ideal For Fall Weeding

| 4 - APPLICATION ANNUAL LAWN CARE PROGRAM | | | |
|-----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------|
| (ILLUSTRATIONS) | | | |
| Apply 1st in my Annual Lawn Care Program in the spring. Controls Crabgrass while feeding your lawn. | Apply 2nd in my Annual Lawn Care Program spring through fall. Controls Broadleaf Weeds when they appear while feeding your lawn. | Apply 3rd in my Annual Lawn Care Program in summer. Controls insects while feeding your lawn. | Apply 4th in my Annual Lawn Care Program in fall. Provides nutrients for a greener, thicker lawn. |

ADDITIONAL FERTILIZER AND MARKETING

SUGGESTED SPREADING APPLICATIONS

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. This bag covers approximately 5,000 square feet (100 feet x 50 feet). Close spreader hopper and adjust the spreader setting using the Spreader Settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Close the spreader hopper whenever spreader stops or turns. For one pass application rate.

Measure the size of the area to be treated and do not over or under apply. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. DO NOT HAND SPREAD.

| | | |
|--------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|
| ILLUSTRATION | Broadcast Spreader Settings For best results, apply in a circular path starting on the outside and working toward the center (see Broadcast Spreader Path diagram). A little overlap of product will insure complete coverage. | ILLUSTRATION |
| ILLUSTRATION | Drop Spreader Settings First apply 2 border strips along the edge of each end of the lawn (see Drop Spreader Path diagram). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips. | ILLUSTRATION |


Residential

Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

Controls over 250 broadleaf weeds, including: Dandelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.

Contains controlled-release nitrogen for extended feeding.

| Nitrogen | Phosphorus | Potassium |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Essential for plant growth and a thick green lawn. (OR) Nitrogen is essential for giving your lawn lush green color and dense foliage. (OR) For a Dense Green Lawn | To promote root growth and grass multiplication. (OR) Phosphorus helps your grass develop and maintain vigorous root growth. (OR) Improves Root Growth | To help grass resist the elements and stay strong. (OR) (Also referred to as Potash). This nutrient is vital for vigorous growth while helping your lawn cope with drought and other extreme weather conditions. (OR) Vital for growth and drought resistance |



FOR THE BEST LAWN IN TOWN

(DISTRIBUTORS) ANNUAL LAWN PROGRAM

For best results this product should be used in conjunction with (Seller's) Crabgrass Preventer Plus Lawn Food, (Seller's) Lawn Food or (Seller's) Fall Lawn Food.

HOW IT WORKS:

Start anytime! No guesswork, just follow these steps for a thick, green, weed-free lawn. It's easy...simply apply the (DISTRIBUTOR) product that corresponds to the season.

| | | | | Residential |
|-----|---------------------------------------------------------|--------------------------------------------------|------------------------|--------------------------------------------------------------|
| | • PREVENTS Crabgrass & other unsightly weeds | apply early spring | apply mid-late spring | • Controls Dandelions & more than 140 other weeds |
| | • ENCOURAGES quick spring greenup | SELLERS Crabgrass Preventer Plus Lawn Food | SELLERS WEED & FEED | • THICKENS & greens your lawn with extended feeding Nitrogen |
| (A) | • GREENS your lawn with extended feeding Nitrogen | apply fall (A) | apply summer | • GREENS with Iron & Sulfur |
| | • THICKENS fall lawns & encourages quick spring greenup | SELLERS FALL LAWN FOOD | SELLERS LAWN FOOD | • THICKENS & promotes a lush green lawn |
| | • NOURISHES your lawn with extended feeding Nitrogen | | | • NOURISHES your lawn with extended feeding Nitrogen |
| | • GREENS with Iron & Sulfur | | | • GREENS with Iron & Sulfur |

(A) BAG ILLUSTRATIONS

Alternate copy

| (A) | (A) | (A) | (A) | (B) |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Early Spring Crabgrass Preventer Plus Lawn Food • Prevents crabgrass & other unsightly weeds. • Encourages quick greenup. • Greens your lawn with extended feeding nitrogen. | Late Spring Weed & Feed • Controls dandelions & more than 140 other weeds. • Thickens & greens your lawn with extended feeding Nitrogen, Iron & Sulfur. | Summer Lawn Food • Thickens & promotes a lush green lawn. • Nourishes your lawn with extended feeding Nitrogen, Iron & Sulfur. | Fall Fall Lawn Food • Thickens lawns & encourages quick spring greenup. OR Fall Weed & Feed • Kills dandelions & more than 140 other weeds while feeding your lawn. | |

(A) Bag Illustrations

(B) Guarantee Illustration - We guarantee that you will be 100% satisfied with the results of this product. If for any reason you are not completely satisfied, bring back the unused portion of the product and your register receipt & we will cheerfully refund your money.

DO NOT HAND APPLY.

SPECIAL CARE NOTICE: This product contains the micronutrient Iron to help prevent Iron deficiency. Iron may stain concrete surfaces such as sidewalks, patios and swimming pools.

Product should not be applied on water dampened concrete and should be removed promptly after application by sweeping or vacuuming.

CONTROLS ALL COMMON WEED TYPES

Where beautiful gardens begin™

- Controls Dandelions & more than 250 other weeds
- Thickens & greens your lawn
- Greens with Iron & Sulfur

Residential

Contains Poly-S® controlled release nitrogen for extended greening (For best results, use with other lawn care products in the [Seller's Name] Fertilizer Program.)

Contains the 3-way power of Dissolve® for Broadleaf, Creeping and Woody weeds.

Controls over 140 common lawn weeds while feeding your lawn with a premium blend of fertilizer nutrients that provide quick and extended greening.

BROADCAST SPREADER SETTINGS

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

DROP SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Do it Best® Lawn Care Program

The following represents the suggested application timing of the products listed in the Do it Best® Program. Abnormal climatic conditions can affect appropriate timing.

U.S. Climate Zones:



■ Bluegrass, Ryegrass, Fine & Tall Fescues

■ Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagrass, Zoysiagrass

■ Bermudagrass, Zoysiagrass, Centipedegrass, Bahiagrass, St. Augustinegrass

Do it Best® 28-3-4 Crabgrass Preventer + Lawn Fertilizer

- Promotes a lush, green lawn while controlling crabgrass and most other annual weed grasses.

- Contains the pre-emergent herbicide, Team®

■ Apply mid March - April

■ Apply early March - mid April

■ Apply mid February - late March

Do it Best® 28-3-3 Weed & Feed

- Controls over 200 common lawn weeds while feeding your lawn with a premium blend of nutrients.

- Contains the premium broadleaf herbicide Dissolve®

■ Apply mid April - late June

■ Apply late March - May

■ Apply mid March - mid May

Do it Best® 18-3-5 Fertilizer + Insect Control

- Controls ants, fleas, ticks, chiggers, crickets, lawn moths and many more.

- Contains slow release nitrogen to extend feeding of the lawn.

■ Apply April - August Apply

■ mid May - early September

■ Apply mid April - September

Do it Best® 22-4-14 Winterizer

- Formulated for fall application to prepare your turf for winter

- Contains slow release nitrogen to extend feeding of the lawn.

■ Apply September - mid October

■ Apply September - late October

■ Apply September - mid November

Other Do it Best® Fertilizers

- 29-3-4 Premium Lawn Fertilizer

- 22-4-11 Winterizer + Weed Control

- 20-27-5 New Lawn Starter

Team is a registered trademark of Dow AgroSciences; Dissolve® is a registered trademark of Nufarm Americas, Inc.

GreenThumb Lawn Care Program

The following represents the suggested application timing of the products listed in the GreenThumb Program. Abnormal climatic conditions can affect appropriate timing.

U.S. Climate Zones:



■ Bluegrass, Ryegrass, Fine & Tall Fescues

■ Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagrass, Zoysiagrass

■ Bermudagrass, Zoysiagrass, Centipedegrass, Bahiagrass, St. Augustinegrass

GreenThumb Premium Crabgrass Preventer + Lawn Fertilizer

■ Apply mid March - April Apply

■ early March - mid April Apply

■ mid February - late March

- Promotes a lush, green lawn while controlling crabgrass and most other annual weed grasses.

- Contains the pre-emergent herbicide, Team®

GreenThumb Premium Weed & Feed Lawn Fertilizer

■ Apply mid April - late June

■ Apply late March - May

■ Apply mid March - mid May

- Controls over 200 broadleaf weeds, including Dandelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.

- Contains the premium broadleaf herbicide Dissolve®

GreenThumb Premium Lawn Fertilizer

■ Apply June - August Apply

■ mid May - early September Apply

■ mid April - September

- Specially formulated to promote even, lush growth of your grass.

- Contains slow release nitrogen to extend feeding of the lawn.

GreenThumb Premium New Lawn Starter

■ Apply anytime before seeding and laying sod

■ Apply anytime before seeding and laying sod

■ Apply anytime before seeding and laying sod

- Formulated with high phosphate to encourage new grass to develop a strong root system

- Contains controlled-release nitrogen for extended feeding

GreenThumb Premium Winterizer Lawn Fertilizer

■ Apply September - mid October

■ Apply September - late October

■ Apply September - mid November

- Formulated for fall application to prepare your turf for winter.

- Contains slow release nitrogen to extend feeding of the lawn.

*Team is a registered trademark of Dow AgroSciences LLC
Dissolve® is a registered trademark of Nufarm Americas Inc.,
Patent No. 5,266,553

Late Spring Application

Kills weeds* & promotes hardy, lush lawns.

*when used as directed

Apply with a broadcast spreader (diagram of spreader) or drop spreader (diagram of spreader).

Residential

Optional Language:

- 1) NOTICE: This product contains the micronutrient iron for extra greening and to help prevent iron deficiency.
Iron may stain concrete and other surfaces. Misapplied product to these surfaces should be removed promptly by sweeping or blowing. Do not wash off with water.
- 2) Formulated for Midwestern Lawns
- 3) Available from:
Hy-Vee Inc.
5820 Westown Parkway West
Des Moines, IA 50266
Ideal for lakeside communities and those areas with phosphorus fertilizer restrictions.
Phosphorus Free Application.
- 4) Note to reviewer: a) Non-misleading graphics depicting sub-registrant company logos may be represented in Final Printed Labeling. b) Distributor labels may have the following: [Exclusively] distributed by: [name of distributor] [address of distributor] [company logo of distributor]
- 5) [Note to reviewer: DISTRIBUTOR SPREADER SETTINGS - Broadcast and Drop Spreader setting/rate table may be adjusted to agree with bag size and distributor partnerships]

- 6) [OPTIONAL SPREADER SETTINGS]
[These settings are furnished as a guide ONLY. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.] [We recommend the cyclone (roto, broadcast) spreaders.] [Calibrate your spreader according to manufacturer's directions]
[They cover [XXXX] sq ft (XX' x XX') in about [XX] minutes.]
[Overlap X" to X" and you will get professional results without streaking.] [On standard models-Set all makes about [X] of total opening.]

[OPTIONAL FERTILIZER STATEMENTS]:

- [For Fertilizer Mixtures Containing Nitrogen]: [It is a violation of law in [Maryland][New Jersey][New Hampshire][list state(s)] to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen, or more than 0.9 lb per 1,000 sq.ft. of total nitrogen, with a least 20% of which shall consist of slow release nitrogen, per application. Additional restrictions apply to total nitrogen applied to turf per season.]
- In Maryland, do not apply this product at more than 0.7 lbs nitrogen per 1,000 sq ft.
- Provides controlled release nutrient feeding from [XXX] Technology.
- Controlled / slow release nitrogen for extended feeding
- Contains (X%) iron for deep greening
- Extended Feeding with [XXX] technology
- Non-staining [formula]
- Slow-release nitrogen feeds your lawn [turfgrass] over [X] months/weeks
- Feeds [X] months for a lush and thick lawn

Residential

- Promotes thick and lush lawns
- Helps develop a deep, dense root system that encourages early spring green-up
- Encourages beneficial soil microbes
- Promotes healthier soil
- Contains, non-burning, slow release nitrogen and iron for deep greening
- [Virginia applications: [Apply no more than 0.7 lb per 1,000 sq. ft. of readily available nitrogen during any given 30-day period. Apply no more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq. ft. of total nitrogen, with at least 20% of which shall consist of slow-release nitrogen, during any given 30-day period.]
- [For Fertilizer Mixtures Containing Phosphorous]: [For starter lawns in MD]: [In Maryland, a soil test is not required for a new lawn. Apply either 2 lbs. of phosphorus per 1,000 sq. ft. initially or apply 1 lb. of phosphorus per 1,000 sq. ft. twice during the starter year. On established turf, this product must be used in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application.]
- [For levels of phosphorus 5% or below]: [This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when (1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; (2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or (3) reestablishing or repairing a turf area. Not for use on established lawns or grass in Maryland.]
- [For all %'s of phosphorus-containing fertilizers in VA]: [Virginia Applications]: [This lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.]
- [Do not apply to any turfgrass in North Florida in winter]
- [State Specific Use Restriction if distributed in Florida]: Do not apply to any [use site] in [season] in [state].

Do It Best Fertilizer Program

The following represents the suggested application timing of the products listed in the Do It Best Program. Abnormal climatic conditions can affect appropriate timing.

U.S. Climate Zones:

| Climate Zone | Grasses |
|-------------------|----------------------------------------------------------------------------|
| NORTHERN ZONE | Bluegrass, Ryegrass, Fine & Tall Fescues |
| TRANSITIONAL ZONE | Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagrass, Zoysia grass |
| SOUTHERN ZONE | Bermudagrass, Zoysia grass, Centipedegrass, Bahiagrass, St. Augustinegrass |

Do It Best® Crabgrass Preventer Plus Lawn Fertilizer

- Promotes a lush, green lawn while controlling crabgrass and most other annual weed grasses.
- Contains the preemergent herbicide, Team®

| Timing |
|---------------------------------|
| Apply mid March - April |
| Apply early March - mid April |
| Apply mid February - late March |

Do It Best® Weed & Feed

- Controls over 140 broadleaf weeds, including Dandelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.
- Contains the premium broadleaf herbicide Dissolve®

| Timing |
|-----------------------------|
| Apply mid April - late June |
| Apply late March - May |
| Apply mid March - mid May |

Do It Best® Lawn Fertilizer

- Specially formulated to promote even, lush growth of your grass.
- Contains slow release nitrogen to extend feeding of the lawn.

| Timing |
|---------------------------------|
| Apply June - August |
| Apply mid May - early September |
| Apply mid April - September |

Do It Best® Summerizer Lawn Fertilizer

- Special non-burning formula, when used as directed, for light summer application.
- Contains slow release nitrogen to extend feeding of the lawn.

| Timing |
|---------------------------------|
| Apply June - August |
| Apply mid May - early September |
| Apply mid April - September |

Do It Best® Winterizer

- Formulated for fall application to prepare your turf for winter.
- Contains slow release nitrogen to extend feeding of the lawn.

| Timing |
|--------------------------------|
| Apply September - mid October |
| Apply September - late October |
| Apply September - mid November |

Team is a reg. TM of Dow AgroSciences LLC; Dissolve - Riverdale Chemical Co., Patent No. 6,295,563

| Commercial | | | |
|------------|-------|---|-----------|
| 2,4-D | GROUP | 4 | HERBICIDE |
| MCPP- P | GROUP | 4 | HERBICIDE |
| 2,4-DP-P | GROUP | 4 | HERBICIDE |

SWEET SIXTEEN WEED AND FEED WITH DISSOLVE®

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or Covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

| | |
|----------------------------------------------------------------------------|-----------------|
| Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* | 0.700% |
| Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid**‡ | 0.182% |
| Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***‡ | 0.179% |
| OTHER INGREDIENTS: | 98.939% |
| TOTAL: | 100.000% |

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

| | |
|--------------------------------------------------------------------|---------------------|
| *2,4-Dichlorophenoxyacetic Acid..... | 0.58%, 0.0058 ae/lb |
| **(+)-R-2-(2-Methyl-4-Chlorophenoxy) Propionic Acid | 0.15%, 0.0015 ae/lb |
| ***(+)-R-2-(2,4-Dichlorophenoxy) Propionic Acid | 0.15%, 0.0015 ae/lb |
| †CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p. | |

GUARANTEED ANALYSIS:*

| | |
|-------------------------------------------------------------------------------------------------------------------------------|----|
| Total Nitrogen (N) | X% |
| X% Ammoniacal Nitrogen | |
| X% Urea Nitrogen | |
| Available Phosphate (P ₂ O ₅)..... | X% |
| Soluble Potash (K ₂ O). | X% |
| Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash. | |
| Chlorine (Cl) Maximum..... | X% |
| The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen. | |
| Fertilizer guaranteed by (Name of fertilizer registrant) | |

Commercial

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS
Alternative Text: [SEE] [REFER TO] [INSIDE] [LABEL BOOKLET] [BACK PANEL] [FOR] [FIRST AID] [AND]
[ADDITIONAL] [PRECAUTIONARY STATEMENTS] [AND] [DIRECTIONS FOR USE]

For Medical Emergencies, Call (877) 325-1840

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

[Note to reviewer: Any text found in brackets “[]” is optional on container label.]

EPA REG. NO. 228- 280
EPA EST. NO. 228-IL-1

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



NET WEIGHT: LBS. (Kg)

000228-00280. 20250307

Commercial

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

| | |
|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice. |
| IF INHALED | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |
| IF IN EYES | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators. In areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

GROUNDWATER ADVISORY

2,4-DP-p is known to leach through soil into groundwater under certain conditions as a result of label use.

This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Commercial

PRODUCT RESTRICTIONS

Do not enter or allow people (or pets) to enter the treated area until dusts have settled.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Only protected handlers may be in the area during application.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

A minimum of 30 days between applications is required.

WEED RESISTANCE MANAGEMENT

For resistance management, this product contains a Group 4 herbicide 2,4-D. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. Appropriate resistance management strategies should be followed.

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the desirable plants and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest controls advisors for additional pesticide resistance-management and/or integrated weed management recommendations for specific types of plants and weed biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. Do not assume that each listed weed is being controlled by this mechanism of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled.

PRODUCT USE INFORMATION

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf such as Lawns (Residential, Industrial, and Institutional), Parks, Cemeteries, Athletic Fields, and Golf Courses (Fairways, Aprons, Tees, and Roughs), and similar turf areas.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggarstick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Black-seed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canadian thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horse nettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Knotweed, Knavel, Kochia, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morning glory, Mouseear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Nutgrass, Orange hawkweed, Oxalis, Oxyeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy,

Commercial

Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanisneedle, Spatterdock, Speedwell, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, and Yellowflower pepperweed.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers, especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 pounds for 16 pound bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.7 pound (0.8 pound for 16 pound bag) per 250 square feet.

| SPREADER SETTINGS ONE PASS APPLICATION [*] : | |
|----------------------------------------------------------|---------|
| MAKE | SETTING |
| Central Calibrated | 5 |
| Cyclone Broadcast | 3-3/4 |
| Jackson | 5 |
| Sears Broadcast | 3 |
| Sears | 5 |
| Scotts Calibrated | 4 |

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Use Precautions for Turf, ornamental (golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas)

Postemergence:

Limited to 2 applications per year.

Maximum of 250 pounds of product /acre per application.

The maximum seasonal rate is 500 pounds of product/acre, excluding spot treatments.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

Commercial

*Based on Inert Fertilizer claimed.

| | |
|--------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| [Environmental Benefits] (Grasscycling Logo) | <p>Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:</p> <p><i>Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.</i></p> <ul style="list-style-type: none">✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day✓ Reduces Glare & Reduces Noise✓ Traps Dust, Smoke Particles & Other Pollutants✓ Purifies Water & Modifies Temperature✓ Improves Soil Condition & Controls Erosion |
|--------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

For nonrefillable bags of granular or dry formulation

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

or

Solid dilutable formulations in nonrefillable plastic containers

CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

CONTAINER DISPOSAL: Nonrefillable containers larger than 5 Gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Commercial

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV)

Dissolve and The Optical Advantage are registered trademarks of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.

Commercial

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

Apply with a broadcast or drop spreader.

OPTIONAL SPREADER INSTRUCTIONS:

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Additional Fertilizer and Marketing Claims

For Thick, Green, Weed-Free Lawns

Extended Greening From Controlled Release Nitrogen

For best year round results, this product should be used in conjunction with (product name) Lawn Food.

Refer to the table and picture below to determine the size of your lawn. Refer to "Path A" when using a broadcast spreader and "Path B" when using a drop spreader.

Kills weeds* & promotes hardy, lush lawns

*when used as directed

| | | |
|---|---------------|--------------------------------------|
| - | Length (Feet) | Illustration of Property |
| X | - | |
| = | - | |
| - | - | |
| = | - | Total Square Feet Coverage Needed |
| | | Illustration of Path A |
| | | Illustration of Path B |

Contains Dissolve® A Premium Broadleaf Herbicide

Additional Marketing Statements

- Feeds desirable grasses
- Controls Dandelions and more than 250 other Broadleaf Weeds
- Controlled release Nitrogen feeds grass longer

Optional language:

Ideal for lakeside communities and those areas with phosphorous fertilizer restrictions.
Phosphorous Free Application.

Commercial

ADDITIONAL FERTILIZER AND MARKETING

SUGGESTED SPREADING APPLICATIONS

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. This bag covers approximately 5,000 square feet (100 feet x 50 feet). Close spreader hopper and adjust the spreader setting using the Spreader Settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Close the spreader hopper whenever spreader stops or turns. For one pass application rate.

Measure the size of the area to be treated and do not over or under apply. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. DO NOT HAND SPREAD.

| | | |
|--------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|
| ILLUSTRATION | Broadcast Spreader Settings For best results, apply in a circular path starting on the outside and working toward the center (see Broadcast Spreader Path diagram). A little overlap of product will insure complete coverage. | ILLUSTRATION |
| ILLUSTRATION | Drop Spreader Settings First apply 2 border strips along the edge of each end of the lawn (see Drop Spreader Path diagram). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips. | ILLUSTRATION |


Commercial

Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

Controls over 250 broadleaf weeds, including: Dandelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.

Contains controlled-release nitrogen for extended feeding.

| Nitrogen | Phosphorus | Potassium |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| Essential for plant growth and a thick green lawn. (OR) NITROGEN For a Dense Green Lawn | To promote root growth and grass multiplication. (OR) PHOSPHATE Improves Root Growth | To help grass resist the elements and stay strong. (OR) POTASH Vital for growth and drought resistance |



- Length (Foot)
- x Width (Foot)
- Total Square Foot
- Non-Lawn Areas
- Total Square Foot coverage you need

FOR THE BEST LAWN IN TOWN