NOTIFICATION

## RIVERDALE®

DISSOLVE® WEED AND FEED

AUG 0 5 2004

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

## GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:  Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*
ISOMER SPECIFIC METHOD, EQUIVALENT TO:  *2,4-Dichlorophenoxyacetic Acid
GUARANTEED ANALYSIS:  Total Nitrogen (N)
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.
Fertilizer guaranteed by (Name of fertilizer registrant)
Riverdale, Dissolve and The Optical Advantage are Registered Trademarks of Nufarm Americas Inc.  Patent No. 5,266,553

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-281

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

## MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0863

Revised 2/14/03	Changed First Aid Statement per PR Notice 2001-1 and Disposal Statement
	per PR Notice 2001-6 via Amendment.
6/13/03	Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.
11/10/03	Added marketing hype via Notification.

3/15/04 Added marketing hype to Residential label only, via Notification.

6/28/04 Corrected per EPA's letter dated 5/17/04.

3/24

(Back Panel)

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

### FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN

Take off contaminated clothing. Rinse skin immediately with

plenty of

OR CLOTHING:

water for 15 to 20 minutes. Call a poison control center or

doctor for treatment advice.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DISSOLVE® wEEL AND YEER is a stable mixture of three herbicides. It is for use on Residential Turf.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine. and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyed Susan, Black medic, Black mustard, Blackseed pantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed,

Page 3

Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 (18 or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 4 pounds (3.6 pounds for 18# bag, 3.2 pounds for 16# bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound (.9 pound for 18# bag, .8 pound for 16# bag) per 250 square feet.

(Residential)

Page 4

SPREADER SETTINGS ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Besides increasing property values, did you know a well-maintained lawn provides the following

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P,O, and K,O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. \*Based in Inert Fertilizer claimed.

## **Environmental Benefits** (Grasscycling

Grass clippings and yard waste are approximately

20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

benefits to your home and environment:

- ✓ A 50' x 50' lawn produces enough
- oxygen for a family of 4 every day Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & other Pollutants
  Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

## STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed, or other pesticides.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 061303C)

(Residential)

Page 5

#### OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and

shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the

outside and working toward the center, overlapping very slightly.

#### OPTIONAL MARKETING STATEMENTS:

The following note will not be part of the actual label copy.

°NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various

commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Editor's notes: 1. Subregistrant Warranty Statements previously accepted by U.S. EPA for use on their own registered products are not included on this label.

2. Subregistrant's trademark name acknowledgments are not included on

this label.

### Alternate (Optional) Marketing and Fertilizer Statements

Contains Dissolve® A Premium Broadleaf Herbicide

Provides Fast Green-Up & Slow Release for Long Feeding.

Easy to Use

Satisfaction Guaranteed

If you are not satisfied with the product results, simply send proof of purchase to (Name of Distributor) for replacement product voucher.

If you apply (product name) as directed, and are not completely satisfied, send proof of purchase from this bag and the cash register receipt to the address provided. We will promptly send your money back.

CONTAINS ELEVEN ESSENTIAL ELEMENTS, INCLUDING EXTENDED TIME-RELEASE NITROGEN, TO PRODUCE EXCELLENT RESULTS REGARDLESS OF GRASS TYPE OR SOIL QUALITY.

DUAL ACTION FEEDS YOUR LAWN WITH TIMED-RELEASE NITROGEN FOR EXTENDED GREENING

SPECIAL FALL GRASS TIME-RELEASE FERTILIZER

GREENS IN AUTUMN HASTENS SPRING RECOVERY ENCOURAGES EARLY SPRING GREEN-UP

KILLS WEEDS AND FORTIFIES YOUR LAWN.

FOR A LONG-LASTING QUICK GREEN LAWN

WITH SLOW-RELEASE NITROGEN FOR EXTENDED FEEDING

HELPS BUILD A STRONGER, DEEPER ROOT SYSTEM

NEVER SPREAD BY HAND (OR) NEVER APPLY BY HAND

DO NOT APPLY WHEN TEMPERATURE EXCEEDS 90°F

SATISFACTION GUARANTEED (OR) MONEY BACK GUARANTEE

(Residential)

KILLS OVER 250 TYPES OF WEEDS (OR) Kills over 250 varieties of weeds

SOLD EXCLUSIVELY THROUGH (NAME OF STORES)

Contains Polyon® Timed-Release Fertilizer

USED BY PROFESSIONAL TURF MANAGERS

It's the GREEN inside

IMPORTANT - It is important to follow the directions as indicated on this quality lawn care product.

LOOK for other quality (brand name) lawn and garden products. For more information about POLYON® and (brand name) products visit us on the INTERNET at www.XXXXXX.com or call us at 1-800-XXX-XXXX.

The POLYON® Advantage (ILLUSTRATION)

(Brand name) products with POLYON® timed-release fertilizers work differently from other slow-release products. POLYON timed-release fertilizers release in response to soil temperature rather than moisture level. Other slow-release products release according to the amount of moisture present in the soil...the more moisture, the quicker the nutrient release...putting your grass at a risk from burning. Because (Brand name) products contains POLYON timed-release fertilizers, your grass is fed for up to 3 months!

If It's Not GREEN...It's Not POLYON®!

The (Brand name) PROMISE Your satisfaction guaranteed or your money back. If you apply (Product name) with POLYON® timed-release nitrogen as directed, and are not completely

satisfied, send proof of purchase from this package and a cash register receipt to the address below. We will promptly send your money back.

## Alternate (Optional) Marketing and Fertilizer Statements

EFFECTIVELY KILLS(OR CONTROLS) DANDELIONS AND BROADLEAF WEEDS

NITROGEN: Slow release for growing a thick green lawn

PHOSPHATE: Improves root growth and plant food storage capabilities.

POTASH: Vital for growth. Improves plant hardiness and drought resistance.

- \*Provides phosphate and potash plus time-released Nitrogen which encourages grass to fill in bare spots weeds leave behind.
- \*Contains Sulfur which helps maintain dark green color and encourages plants and root growth.

Controls a wide range of broadleaf weeds and feeds your lawn to help it grow thick and green!

 $\mathcal{O}_{f_0}$ 

Page 6

(Residential)

Page 7

Length		
X Width .	TLLISTRATION	To determine how much you need, multiply the length
= Total Sq. Ft.		of your lawn by its width. This equals your total
Non-Lawn Areas		square footage. Deduct any non-lawn areas (house,
= Total Sq. Ft. Coverage Needed		garden, etc). One bag covers 5,000 square feet.

Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader at the beginning of the application; adjust spreader if necessary. Consult owner's manual. DO NOT HAND SPREAD. Close spreader openings when loading, turning or when stopped.

	Broadcast Spreader Settings One Pass Application	
ILLUSTRATION	Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.	ILLUSTRATION

	Drop Spread Settings One Pass Application	
ILLUSTRATION	First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.	ILLUSTRATION

Rids your lawn of undesirable weeds and feeds it to help it grow thick and green! For best results, water lawn first. Apply (product name) while lawn is wet. Wait 48 hours, then water lawn as needed. DO NOT APPLY DURING TIME PERIODS WHEN THE MAXIMUM DAILY TEMPERATURE IS IN EXCESS OF 90°F.

	Broadcast Spreader Settings One Pass Application	
ILLUSTRATION	For best results, apply in a circular path starting on the outside and working toward the center, A little overlap of product will insure complete coverage.	ILLUSTRATION

Page 8

Riverdale Dissolve Weed And Feed

(Residential)

	Drop Spread Settings One Pass Application	
ILLUSTRATION	First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.	ILLUSTRATION

Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

(Additional Optional Fertilizer and Advertising Information)

For use on Fescue, Rye, Blue and other listed grasses

Controls over 250 common lawn weeds while feeding your lawn with a premium blend of fertilizer nutrients that provide quick and extended greening.

Contains the 3 way power of Dissolve for broadleaf, creeping and woody weeds.

For thick, green, weed-free turf.

Contains Poly-So controlled released nitrogen for extended greening.

(For best results, use with other lawn care products in the [Seller-name] Fertilizer Program)

(Attach Sam Walton info)

# Sam Walton believed in the ability of American workers to produce the finest products in the world.

That's why we're searching the USA to create something special. Innovative products, made to our own higher standards, that we're proud to call Sam's American Choice<sup>™</sup>. We believe these products, available at Wal-Mart and Wal-Mart Supercenters, offer better value than the leading national brands at the always low Wal-Mart prices. We think you'll agree, or we'll refund your money, no questions asked!

"Better products at the always low Wal-Mart prices..."

Always

Sam m. Walton

We guarantee it.

Sam Walton

## COMPETITIVE EDGE SCHOLARSHIP FUND

In the next 10 years, the United States will have the jobs Americans need, but Americans may not have the skills these jobs demand. To have a competitive edge in the future, American must have a well-

(Residential)

Page 9

educated, trained work force.
To help achieve this goal, Wal-Mart and our vendor partner will each donate 1% of the cost of this product to our Competitive Edge Scholarship Program.

#### OUR ENVIRONMENT:

### (ILLUSTRATION)

- \* Exclusive DeepGreen® Feeding System Helps Improve A Lawn's Winter Hardiness While Eliminating Weeds.
- \* Exclusive DeepGreen® Feeding System Keeps Lawns Green Whatever the Growing Conditions Hot or Cold, Wet or Dry, While Controlling Weeds.
- \* Builds Strong Root Systems And Promotes Spring Green-Up.
- \* Greens up to Three Times Longer than Ordinary Soluble Lawn Fertilizer.

(Logo) DeepGreen - Whatever the Weather - Hot, Cold, Wet, Dry

## EXPERIENCE THE (SELLER'S NAME) ADVANTAGETM

(Seller's Name) has the extraordinary DeepGreen® nutrient delivery system containing an exclusive (or both IBDU® and) polymer-coated time-release nitrogen source(s), assuring you the predictable, continuous feeding ordinary soluble lawn fertilizers can't deliver. So you'll enjoy the advantage of a consistently beautiful lawn week after week.

(Illustration)

Greening Effect

(Seller's) DeepGreen® System Feeds Your Lawn

Continuously When Ordinary Soluble Fertilizers Can't.

Effectively Controls/kills Dandelions and Other Broadleaf Weeds.

The above suggested spreader settings are approximate. In general, spreaders should be calibrated to apply 1 lb. of this fertilizer product per 275 square feet of lawn.

Be sure to close spreader openings when loading, turning or when stopped.

## (SELLER'S NAME) 100% GUARANTEE OF SATISFACTION

If not totally satisfied with the results of this product, simply send proof of purchase and a brief description of your problem to (Seller name & address) for a prompt refund.

#### (SELLER'S NAME) HELP LINE

Questions concerning lawn care or fertilizer application? Just call toll free 1-888-XXXXXX-1

Made in the U.S. (with flag illustration)

(Residential)

Page 10

For Professional Use Only

Weeds controlled by this product are gradually eliminated

Beautiful lawns made easy

Premium formulated herbicide and fertilizer

Controls over 200 broadleaf weeds

Apply to control dandelions, clover and other broadleaf weeds

The best defense against weeds is vigorous turf. Many times a grass plant requires more of a nutrient that the soil can supply. If deprived of these nutrients, the turf becomes thin, weak and off color. Turf left in this state is vulnerable to infestations of weeds. A properly fertilized lawn is less susceptible to weed infestation and will better tolerate cold, heat, drought and traffic. There is NO substitute for proper fertilization.

Rotary Spreader Settings (various spreaders and settings listed)
Drop Spreader Settings (various spreaders and settings listed)

#### OPTIONAL LIABILITY STATEMENTS:

Do not apply this product to "Floratam" St. Augustine grass in Florida.

Do not allow animals to graze on areas treated with this product.

## GreenThumb Fertilizer Program

The following represents the suggested application liming of the products listed in the GreenThumb Program. Abnormal climatic conditions can affect appropriate timing.

## U.S. Climate Zones:



- Bluegrass, Ryegrass,
- Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagnass, Zoysiagrass
- Bermudagrass, Zoyslagrass, Centipedegrass, Babiagrass, S1. Augustinegrass

## Green Tromb Premium Crabgrass Preventer + Lawo Femilizer

- Promotes a lush, green lawn white controlling crabquass and most other annual weed grasses.
- · Contains the pre-emergent berbicide, Team\*
- Apply mid Match Apill
- Apply Barly Merch Sold Appli
- Apply into Fromary late

### Green Thomb Promisor Weed & Feed Lawy Ferlifizer

- Controls over 140 broadlest weeds, including Candelion, Plantain, Chickweed, Knotweed, Henbit and Spurge.
- Contains the premium broadleaf herbicide Dissolve-
- Apply mid April Jala
- Apply late March May
- Applymed March mid Mar

## GreenThumb Premium Lawn Festilizer

- Specially formulated to promote even, lush growth of your grass.
- Contains slow release mitrogen to extend feeding of the lawn.
- Apply June August
- Apply and May early September
- Apply mit April -Seplander

## Green Tromb Premium Summerizer Lawn Fertilizer

- Special non-burning formula, when used as directed, for light swarmer application.
- Contains slow release mitrogen to extend feeding of the lawn.
- Apply Sine August
- Applyment May early September
- Apply said April -Seciender

## Green Thumb Premium Winterizer Lawn Fertilizer

- Formulated for fall application to prepare your turt for winter.
- Contains slow release mitrogen to extend feeding of the lawn.

Apply September -

Apply September - Jale October

Apply:September - min

Tessu la areg. Yill of Bandites III; Dissains - Historiale Cherakai Fa., Patest No. 5,265,563

Additional and/or Alternate map text Northern Zone: April-June & mid August-mid October Transitional Zone: April-May & September-late October Southern Zone: March-April & mid September-mid November

(Residential)

Page 12

Special nutrient formulation greens and thickens fall lawns and encourages root development for quicker greenup in the spring. Controls over 250 weeds.

Contains the 3-Way Power of Dissolve® for fall control of Broadleaf, Creeping and Woody Weeds.

For thick, green lawns this fall and next spring.

Controls dandelions and over 250 other weeds.

X% Nitrogen: Gives lawns their green color. Fertilizers with controlled release nitrogen formulas will feed and green your lawn over a longer period of time.

X% Phosphate: This nutrient is important for normal root development. Good roots are vital for grass to take up water and nutrients.

X% Potassium: Also called potash, helps lawns better tolerate extreme weather conditions. Contributes to the overall vigor or grass.

## Additional Step 4 to Sam's Choice Fertilizer Program:

Step 4. Sam's Choice™ WINTERIZER LAWN FOOD OR WEED & FEED

- \* Winterizer Weed & Feed fertilizes while controlling existing weeds; if weeds are not evident, use Winterizer Lawn Food to nourish your lawn and prepare it for winter.
- \* Both contain a premium blend of fertilizer nutrients that help your lawn through the winter and speed greening in the spring.

Apply mid August-mid October Apply September - late October Apply mid September - mid November

(Diagram of bags)

To determine square footage of application area, follow calculation to the right.

#### THE COUNTRY COTTAGE STORY

Country Cottage products are designed with the home gardener in mind. They are produced with only high quality ingredients and are designed to help you have a better garden and be a better gardener.

When you shop at an authorized Country Cottage Dealer you are getting more than just the best quality products. What you are also getting is years of experience and strong horticultural know-how to answer all your gardening questions and solve all of your gardening problems.

When you think about projects in your lawn and garden, think about your Country Cottage Dealer, the hometown neighbor, a premier garden center. The products you want and the advice you need, all at one convenient location.

- Special Analysis formulated for Southern grasses
   Special for Southern Turf
- PCT Slow Release Nitrogen for extended feeding Poly-Coated technology slow-release nitrogen

Information regarding the contents and levels of metals in the product is available on the Internet at http://www.wa.gov/agr/

THE CHECK MEANS IT'S PROVEN®

- ✓ Four-step program may be started at any stage
- ✓ Contains slow-release nitrogen for extended greening
- ✓ Contains the 3-way power of Dissolve® for broadleaf, creeping and woody weeds

Riverdale Dissolve Weed And Feed (Residential)

Page 13

NITROGEN

PHOSPHORUS

POTASSIUM

For a lush, Improves Vital for growth & green lawn root growth drought resistance

### ILLUSTRATIONS OF FOUR BAGS

CRABGRASS PREVENTER	WEED & FEED	LAWN FERTILIZER	FALL FERTILIZER/NEW LAWN STARTER
turf to control crabgrass, foxtail, goosegrass, barnyardgrass and annual bluegrass.	Promotes lush, green lawn growth and controls dandelions and over 140 broadleaf weeds.	Apply throughout the growing season for a show-place lawn.	Promotes vigorous growth of new grass in spring and encourages root development, winter hardiness and disease resistance in fall.

Controls Dandelions and over 100 other broadleaf weeds

Nitrogen - For vigorous blade, leaf and stem growth and dark green color (or) Contains no Nitrogen

Phosphate - For root development and growth (or) Contains no Phosphorus

Potash - Promotes strong roots and improves turf hardiness (or) Contains no Potassium

## ALTERNATE ARRANGEMENT OF LISTED WEEDS

Alder Artichoke Aster Aster, Wild Bedstraw Bedstraw, Pineywoods Beggartick Betony, Florida Biden Bindweed Bindweed, Field Bindweed, Hedge Bitter wintercress Bittercress, Hairy Bitterweed Black-eyed Susan Bloodflower,	Milkweed Boxelder Brassbuttons Bristly Oxtongue Broomweed Buckhorn Buckwheat, Wild Bullthistle, Wavyleaf Bur ragweed Burdock Burweed Buttercup Buttercup Buttercup, Bulbous Carpetweed Carrot, Wild Catchweed Bedstraw	Catnip Catsear Catsear, Spotted Chamomile, Corn Chickweed Chickweed, Common Chickweed, Mousear Chicory Cinquefoil Cinquefoil, Rough Clematis, Western Clover Clover, Annual Yellow Sweet Clover, Crimson Clover, Hop
Clover, Red Clover, Strawberry Clover, Sweet Coreopsis, Plains Creeping jenny Croton Croton, Wooly	Clover, White Cockle Cocklebur  Cudweed Curly indigo Daisy, English Daisy, Oxeye	Cocklebur, Spiny Coffeebean Coffeeweed  Dandelion Dandelion, Fall Dandelion, False Dock
Dock, Broadleaf Dock, Curly	Dock, Smooth Dogbane	Dogfennel Elderberry

r
_

Mullein, Common Mustard Mustard, Black Mustard, Garlic Mustard, Hedge Mustard, Tansy	
Mustard, Tumble Mustard, White Mustard, Wild Nettle Nettle, Bull	
Nettle, Burning Nettle, Dead Nettle, Stinging Nettle, Tall Oak, Poison	
Onion, Wild Oxalis Pansy, Field Pantain, Blackseed Parsley-piert	
Parsnip Parsnip, Wild Pearlwort Pennycress Pennywort	
Peppergrass Pepperweed Pepperweed, Virginia Pigweed Pigweed, Prostrate Pigweed, Redroot Pigweed, Russian	
Pigweed, Smooth Pigweed, Tumble Scarlet Pimpernel	
Plantain Plantain, Blackseed Plantain, Bracted Plantain, Broadleaf Plantain, Buckhorn Plantain	
Plantain, Hoary Plantain, Narrowleaf Plantain, Slender Plantain, Wooly Pokeweed	
Poorjoe Povertyweed Prairie Prickly Sida Primrose Puncture vine	
Puncture Vine Purslane Radish, Wild Ragweed Rape, Wild Rose, Southern Wild	
Rush Salsify, Western Shepherdspurse Smartweed	

Sorrel, Red Sorrel, Sheep Sowthistle Sowthistle, Common Spanishneedle Spatterdock Speedwell Spiny Spiny Amaranth Spurge Spurge, Prostrate Spurge, Spotted Spurweed St. Johnswort Stinkweed Stitchwort Strawberry, Wild Sumac Sunflower Sunflower, False Sweet Potato, Wild Tansy Ragwort Tanweed Tarweed Thistle Thistle, Blessed Thistle, Bull Thistle, Canada Thistle, Musk Thistle, Russian Thistle, Scotch Toadflax Trailing Crownvetch Trefoil, Tick Tumbleweed Velvet leaf Veronica Vervain Vervain, Blue Vervain, Hoary Vervain, Prostrate Vervain, Tall Vetch Vetch, Bird Vetch, Milk Vetch, Narrowleaf Vetch, Wild Virginia Buttonweed Virginia Creeper Willow Witchweed Woodsorrel Wormseed Yarrow Yellow rocket Yellowflower Pepperweed And other broadleaf weeds

Sneezeweed

## RIVERDALE®

## DISSOLVE® WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Sq. Ft. (50' x 100')

## GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*
ISOMER SPECIFIC METHOD, EQUIVALENT TO:  *2,4-Dichlorophenoxyacetic Acid
GUARANTEED ANALYSIS:  Total Nitrogen (N)
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.
Fertilizer guaranteed by (Name of fertilizer registrant).  Riverdale, Dissolve and The Optical Advantage are Registered Trademarks of Nuferr Americas Inc.  Patent No. 5,266,553

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-281

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

## MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 2/14/03 Changed First Aid Statement per PR Notice 2001-1 via Amendment. 6/13/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

11/10/03 Added marketing hype to Residential label only via Notification. 3/15/04 Added marketing hype to Residential label only, via Notification.

6/28/04 Corrected per EPA's letter dated 5/17/04.

16/21

(Back Panel)

#### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks, shoes and gloves when loading and applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

### FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

## ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP, and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pasticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DISSOLVE® WEED AND FEED is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or

Riverdale Dissolve Weed and Feed (Commercial)

Page 4

browning of grass.

HOW TO APPLY: This 20 (or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds (3.6 pounds for 18# bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound (.9 lb. for 18# bag) per 250 square feet.

SPREADER SETTINGS
ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing  $\frac{*(No. \text{ of lbs.})}{(No. \text{ of lbs.})}$  pounds of nitrogen per 1,000 square feet (optional  $P_2O_5$  and  $K_2O$  levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. \*Based on Inert Fertilizer claimed.

# Environmental Benefits

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Senetits (Grasscycling

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✔ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✔ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion



### STORAGE AND DISPOSAL

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feed and foodstuffs. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 061303C)

## OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and

shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the

outside and working toward the center, overlapping very slightly.

#### OPTIONAL MARKETING STATEMENTS:

Contains Dissolve® A Premium Broadleaf Herbicide

The following note will not be part of the actual label copy.

NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

## ALTERNATE ARRANGEMENT OF LISTED WEEDS

Alder
Artichoke
Aster
Aster, Wild
Bedstraw
Bedstraw, Pineywoods
Beggartick
Betony, Florida
Biden
Bindweed
Bindweed, Field
Bindweed, Hedge
Bitter wintercress
Bittercress, Hairy

Bitterweed

Black-eyed Susan
Bloodflower,
Milkweed
Boxelder
Brassbuttons
Bristly Oxtongue
Broomweed
Buckhorn
Buckwheat, Wild
Bullthistle,
Wavyleaf
Bur ragweed
Burdock
Burweed
Buttercup

Buttercup, Bulbous Carpetweed Carrot, Wild Catchweed Bedstraw Catnip Catsear Catsear, Spotted Chamomile, Corn Chickweed Chickweed, Common Chickweed, Mousear Chicory Cinquefoil Cinquefoil, Rough

Clematis, Western

Plantain Clover Ironweed Plantain, Blackseed Clover, Annual Yellow Ivy, Poison Plantain, Bracted Jewelweed Sweet Clover, Crimson Plantain, Broadleaf Jimsonweed Knapweed, Spotted Plantain, Buckhorn Clover, Hop Plantain, Hoary Clover, Red Knawel Clover, Strawberry Knotweed Plantain, Narrowleaf Clover, Sweet Knotweed, Prostrate Plantain, Slender Kochia Clover, White Plantain, Wooly Cockle Lambsquarter Pokeweed Cocklebur Lespedeza Poorjoe Cocklebur, Spiny Lettuce, Blue Povertyweed Coffeebean Lettuce, Prickly Prairie Coffeeweed Lettuce, Wild Prickly Sida Coreopsis, Plains Primrose Locoweed Puncture vine Creeping jenny Lupine Croton Mallow Purslane Mallow, Indiana Mallow, Venice Croton, Wooly Radish, Wild Cudweed Ragweed Curly indigo Marigold, Rape, Wild Daisy, English Roundleafed Rose, Southern Wild Daisy, Oxeye Marigold, Wild Rush Dandelion Marshelder Salsify, Western Dandelion, Fall Matchweed Shepherdspurse Dandelion, False Medic, Black Smartweed Dock Mexicanweed Sneezeweed Dock, Broadleaf Sorrel, Red Morningglory Sorrel, Sheep Dock, Curly Morningglory, Wooly Dock, Smooth Mugwort Sowthistle Dogbane Mullein, Common Sowthistle, Common Dogfennel Mustard Spanishneedle Elderberry Mustard, Black Spatterdock Fiddleneck Mustard, Garlic Speedwell Mustard, Hedge Mustard, Tansy Fieldcress, Austrian Spiny Filaree, Redstem Spiny Amaranth Flax, False Mustard, Tumble Spurge Spurge, Prostrate Flea bane (daisy) Mustard, White Fleabane, Hairy Mustard, Wild Spurge, Spotted Fleabane, Rough Nettle Spurweed Flixweed Nettle, Bull St. Johnswort Florida pusley Nettle, Burning Stinkweed Four-O'-Clock, Wild Nettle, Dead Stitchwort Frenchweed Nettle, Stinging Strawberry, Wild Galinsoga Nettle, Tall Sumac Galinsoga, Smallflower Oak, Poison Sunflower Sunflower, False Garlic, Wild Onion, Wild Geranium, Carolina Oxalis Sweet Potato, Wild Geranium, Wild Pansy, Field Tansy Ragwort Goathead Tanweed Plantain, Blackseed Goatsbeard Parsley-piert Tarweed Goldenrod Thistle Parsnip Thistle, Blessed Ground-Ivy Parsnip, Wild Groundsel Pearlwort Thistle, Bull Gumweed Pennycress Thistle, Canada Thistle, Musk Hawkweed Pennywort Hawkweed, Orange Thistle, Russian Peppergrass Thistle, Scotch Healall Pepperweed Heartleaf drymary Pepperweed, Toadflax Trailing Crownvetch Heathaster Virginia Trefoil, Tick Hemlock, Poison Pigweed Pigweed, Prostrate Hemp Tumbleweed Pigweed, Redroot Henbit Velvet leaf Hoary cress Pigweed, Russian Veronica Pigweed, Smooth Honeysuckle Vervain Pigweed, Tumble Vervain, Blue Horsenettle Horsetail Scarlet Pimpernel Vervain, Hoary

## Riverdale Dissolve® Weed and Feed (Commercial)

Page 7

Vervain, Prostrate Vervain, Tall

Vetch

Vetch, Bird Vetch, Milk Vetch, Narrowleaf Vetch, Wild

Virginia Buttonweed Virginia Creeper

Willow Witchweed Woodsorrel Wormseed

Yarrow Yellow rocket Yellowflower Pepperweed And other broadleaf

weeds



Nufarm Americas Inc.

1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527-0866 Phone: 708.754.3330 Fax: 708.754.0314 www.us.nufarm.com

UPS

June 28, 2004

Ms. Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Crystal Mall Building #2
1801 Bell Street
Arlington, VA 22202

Dear Ms. Miller:

Subject: Notifications

Riverdale Dissolve Weed and Feed EPA Registration No. 228-281

Revised Label Submitted March 15, 2004

Your letter dated May 17, 2004

Please find enclosed revised labels per your above subject letter. For the record, the deleted phrase had been submitted before the Agency accepted and then later changed its mind. This was not any attempt to bypass our statement that we would no longer change the Disposal Statement, which would include the deleted phrase.

If there is anything else we should do at this time please feel free to contact us immediately.

Thank you for your assistance.

Sincerely yours,

NUFARM AMERICAS INC.

By Russell F. Sawyer Regulatory Affairs Manager

RFS:jb

Enclosures





## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

MAY 1 7 2004

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Russell F. Sawyer Nufarm Americas Inc. 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527-0866

Dear Mr. Sawyer:

Subject:

**Notifications** 

Riverdale Triamine Lawn Weed Killer EPA Registration Number 228-181

Riverdale Triamine Liquid Weed and Feed

**EPA Registration Number 228-189** 

Riverdale Premium 3333 Liquid Weed and Feed

**EPA Registration Number 228-237** 

Riverdale Sweet Sixteen Weed and Feed with Dissolve

EPA Registration Number 228-280 Riverdale Dissolve Weed and Feed EPA Registration Number 228-281

Your Submissions Dated November 10 and December 22, 2003 and

March 15, 2004

The submissions referred to above, submitted as notifications are not acceptable. Delete the phrase, "or offered to others for their use," from the **If partly filled** disposal statement. This phrase is not related to disposal and is misleading to the user; therefore, the product could be deemed to be misbranded with such a phrase.

Marketing claims and adding pests may be submitted as notifications as long as you have data on file in your company to support the efficacy claims and the claims are not implied safety claims. See PR Notice 98-10. Copies of PR Notices are available at the following web site:

http://www.epa.gov/PR Notices/

If you have any questions concerning this letter, please contact me at 703-305-6224.

Sincerely yours,

Joanne I. Miller

Product Manager (23)

Herbicide Branch

Registration Division (7505C)

anne D. Miller