

228-283

06/10/2003

1/9

(MASTER LABEL)

RIVERDALE®

TRIPLET® WATER SOLUBLE

Selective Broadleaf Weed Control For Turfgrasses
To Control Dandelion, Clover, Henbit, Plantains, Wild Onion, and
Many Other Broadleaf Weeds

Contains 2,4-D, Mecoprop and Dicamba

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	60.20%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	24.20%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	6.02%
INERT INGREDIENTS:	9.58%
TOTAL	100.00%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	50%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	20%
***3,6-Dichloro-o-anisic Acid	5%

Riverdale and Triplet are Registered Trademarks of Riverdale - A Nufarm Company

Patent No. 5,266,553

FOR SALE TO AND USE BY PROFESSIONAL PERSONNEL ONLY
NOT FOR SALE TO HOMEOWNER

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

SEE SIDE PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: BULK - FIBER DRUMS
(or) 25 lbs. (10 X 2-1/2 lb. packets)
(or) 5 lbs. (8 X 10 oz. packets)
(or) 1 lb. 11-1/2 oz. (40 X 3-1/4 oz. packets)

EPA REG. NO. 228-283

NOTE: Spanish language is optional.

ACCEPTED

JUN 10 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 228-283

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BURR RIDGE, ILLINOIS 60527-0866

2/9

(Side Panels)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER - PELIGRO**

Corrosive, causes irreversible eye damage. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear face shield, goggles or safety glasses and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow and temple protection.

In addition to the clothing and eye protection listed above, commercial mixers/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN
OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
- NOTE TO
PHYSICIAN:** Probable mucosal damage may contraindicate gastric lavage.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigator waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Triplet® Water Soluble is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs), and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

(For 2-1/2 lb. - Water Soluble Packets)

Riverdale Triplet Water Soluble is a stable mixture of three herbicides. Each 2-1/2 lb. water soluble packet will treat from 40,750 to 54,375 square feet (9/10 to 1-1/4 acres) of cool season grasses or 65,000 to 83,000 square feet (1-1/2 to 1-9/10 acres) of Bahia, Bent, common Bermuda, Centipede, St. Augustine, and Zoysia grasses. Maximum kill of common weeds will be obtained by applying in Spring and early Fall when weeds are actively growing.

(For 10 oz. - Water Soluble Packets)

Riverdale Triplet Water Soluble is a stable mixture of three herbicides. Four 10 oz. water soluble packets will treat from 40,750 to 54,375 square feet (9/10 to 1-1/4 acres) of cool season grasses or 65,000 to 83,000 square feet (1-1/2 to 1-9/10 acres) of Bahia, Bent, common Bermuda, Centipede, St. Augustine, and Zoysia grasses. Maximum kill of common weeds will be obtained by applying in Spring and early Fall when weeds are actively growing.

Just add the recommended water to your spray tank, then remove the water soluble packet from plastic outer bag, add the appropriate Triplet into the spray tank, agitate for three to five minutes. Triplet will make a stable water solution. It is non-flammable.

The one acre coverage is equivalent to 1.88 lbs. AI/Acre; the 1-1/3 acre coverage is equivalent to 1.41 lbs. AI/Acre. The 2 acre coverage is equivalent to .94 lb. AI/Acre.

NOTES: (1) Exposure to moisture prior to mixing may cause water soluble bag to rupture. (2) Individual water soluble bags are not to be offered for sale. (3) Four 10 oz. packets are equal to one 2-1/2 lb. packet.

(For 2-3/4 oz. - Water Soluble Packets)

Triplet® W. S. Backpack is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs), and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

Riverdale Triplet W. S. Backpack is a stable, non-flammable mixture of three herbicides. Each 2-3/4 oz. water soluble packet will treat from 3,000 to 4,000 square feet of Bluegrass, Fescue, Rye, Bahia, Bent, common Bermuda, Centipede, St. Augustine, and Zoysia grasses. Maximum kill of common weeds will be obtained by applying in Spring and early Fall when weeds are actively growing.

Just add half the recommended water to your two, three, or four gallon spray tank or handcan, then remove the water soluble packet from plastic outer bag, add one 2-3/4 oz. Triplet water soluble packet into the spray tank, allow two to three minutes to totally solubilize. Shake well. Add remaining water and shake again. Triplet will make a stable water solution.

NOTES: (1) Exposure to moisture prior to mixing may cause water soluble bag to rupture. (2) Individual water soluble bags are not to be offered for sale.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present.

Except as noted, use only lawn-type sprayers. Avoid fine mists. Coarse sprays are less

likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

(All Packages Except 2-3/4 oz.)

Triplet Water Soluble can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

(For 2-3/4 oz. - Water Soluble Packets)

When spot treating in warm season grasses, applications should be limited directly to the weed surface to reduce potential injury to desirable turf species.

WEEDS CONTROLLED BY TRIPLET® WATER SOLUBLE

Bedstraw	Henbit	Purslane
Black medic	Knotweed	Ragweed
Buckhorn	Lambsquarter	Sheep sorrel
Burdock	Lespedeza	Shepherdspurse
Chicory	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Clover	Peppergrass	Wild carrot
Dandelion	Pigweed	Wild garlic
Dock	Plantain	Wild lettuce
Ground ivy	Poison ivy	Wild onion
Healall	Poison oak	Yarrow

and other broadleaf weeds

ORNAMENTAL LAWNS & TURF

APPLICATIONS

(For Bulk-Fiber Drums)

Cool Season Grasses Such as Bluegrass, Fescue and Rye: Use this product at a rate of 2 to 2-2/3 lbs. per acre (3/4 oz. to 1 oz. per 1000 square feet).

For Normal Application - Use 40 to 120 gallons of water per acre (1 to 3 gallons water per 1,000 square feet).

For Professional Lawn Maintenance - Use 1 to 5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses: Apply Triplet Water Soluble at a maximum rate of 1-1/3 to 1-2/3 lbs. in 40 to 200 gallons of water per acre.

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, up to 2-2/3 lbs. per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass: On closely mowed bentgrass (putting and bowling greens), apply this product at a maximum rate of 3/4 ounce in 5 gallons of water per 1,500 square feet - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

(For 2-1/2 lb. - Water Soluble Packets)

Cool Season Grasses Such as Bluegrass, Fescue and Rye: Each 2-1/2 lb water soluble packet will treat 9/10 to 1-1/4 acres (40,750 to 54,375 square feet).

For Normal Application - Use 40 to 120 gallons of water per acre (1 to 3 gallons of water per 1,000 square feet).

For Professional Lawn Maintenance - Use 1 to 5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses: Each 2-1/2 lb. water soluble packet will treat 1-1/2 to 1-9/10 acre (65,000 to 83,000 square feet) in 40 to 200 gallons of water per acre.

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, one 2-1/2 lb. packet treating up to 9/10 acre (40,750 square feet) may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass: On closely mowed bentgrass (putting and bowling greens), apply this product at a maximum rate of one 2-1/2 lb. packet in up to 267 gallons water per 80,000 square feet, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week. This rate is equivalent to 3/4 oz. in 5 gallons of water per 1500 square feet.

(For 10 oz. - Water Soluble Packets)

Cool Season Grasses) Such as Bluegrass, Fescue and Rye: Four 10 oz. water soluble packets will treat 9/10 to 1-1/4 acres (40,750 to 54,375 square feet).

For Normal Application - Use 40 to 120 gallons of water per acre (1 to 3 gallons of water per 1,000 square feet).

For Professional Lawn Maintenance - Use 1 to 5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts of add.

Warm Season Grasses St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses: Four 10 oz. water soluble packets will treat 1-1/2 to 1-9/10 acre (65,000 to 83,000 square feet) in 40 to 200 gallons of water per acre.

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, four 10 oz. packets treating up to 9/10 acre (40,750 square feet) may be used. However, some

hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass: On closely mowed bentgrass (Putting and bowling greens), apply this product at a maximum rate of four 10 oz. packets in up to 267 gallons of water per 80,000 square feet, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week. This rate is equivalent to 3/4 oz. in 5 gallons of water per 1500 square feet.

NOTE

(For Bulk Fiber Drum)

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. 3/4 ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom, as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area, will result in heavy local over-application and subsequent turf discoloration or injury.

Herbi™ Controlled Droplet Applicator: For cool season grasses - Add 1-1/3 to 2 lbs. of Triplet Water Soluble to the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 square feet (approx. 3/4 acre). Do not overlap spray patterns.

For listed warm season grasses - Add 1 to 1-1/3 lbs. of Triplet Water Soluble to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Other Controlled Droplet Applicators (CDA), Atomizers, Spinning Disk Applicators: For cool season grasses - use Triplet Water Soluble at the rate 2 to 2-2/3 lbs. per acre of (3/4 to 1 oz. per 1000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use 1-1/3 to 1-2/3 lbs. of Triplet Water Soluble per acre (1/2 to 2/3 oz. per 1000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume Equipment. Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

(For 2-1/2 lb. - Water Soluble Packets)

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one 2-1/2 lb. packet in 267 gallons of water per 30,000 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom, as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area, will result in heavy local over-application and subsequent turf discoloration or injury.

Other Controlled Droplet Applicators For cool season grasses - Use Triplet Water Soluble at the rate 2-1/2 lbs. per 9/10 to 1-1/4 acre (1 oz. per 1020 to 1360 square

(CDA), Atomizers,
Spinning Disk
Applicators:

feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use one 2-1/2 lb. water soluble packet per 1-1/2 to 1-9/10 acre (1 oz. per 1625 to 2090 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type equipment); and follow CDA spray instructions or cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume
Equipment:

When using a Low Volume Applicator (such as Lawn Doctor's "Turf Tamer"), empty contents of 2-1/2 lb. water soluble packet into 1/2 gallon of water to treat 9/10 to 1-1/4 acre of cool season grasses.

(For 10 oz. - Water Soluble Packets)

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. four 10 oz. packets in 267 gallons of water per 80,000 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom, as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area, will result in heavy local over-application and subsequent turf discoloration or injury.

Herbi™ Controlled
Droplet Applicator:

For cool season grasses - Empty two to three 10 oz. water soluble packets into the Herbi 5 pint container, then fill with 5 pints of water (or 4-1/2 pints of water and 1/2 pint of a surfactant). Keeping container agitated, spray entire contents over 33,000 square feet (approx. 3/4 acre). Do not overlap spray patterns.

For listed warm season grasses - empty one to two 10 oz. water soluble packets of the Herbi 5 pint container and fill with 5 pints of water (or 4-1/2 pints of water and 1/2 pint of a surfactant). Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat drought, etc.

Other Controlled
Droplet Applicators
(CDA), Atomizers,
Spinning Disk
Applicators:

For cool season grasses - Use Triplet Water Soluble at the rate of four 10 oz. packets per 9/10 to 1-1/4 acre (1 oz. per 1020 to 1360 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use four 10 oz. water soluble packets per 1-1/2 to 1-9/10 acre (1 oz. per 1625 to 2090 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume
Equipment:

When using a Low Volume Applicator (such as Lawn Doctor's "Turf Tamer"), empty contents of four 10 oz. water soluble packets into 1/2 gallon of water to treat 9/10 to 1-1/4 acre of cool season grasses.

8/9

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in original container and place in a secured warehouse or storage building. This product should be stored in a cool, dry location. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spillage. Clean up all spilled material with broom. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Fiber Drum - Completely empty plastic drum liner into application equipment and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Fiber drum may be offered for recycling or dispose in same manner as plastic liner. Water Soluble Packaging - Outer plastic bags, cartons, and shipping cases may be offered for recycling or dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

This product is sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

WARRANTY

Nufarm Americas, Inc. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(RV 050703)

NOTE: For all water soluble packets the following statement will be used on the front panel - **"Individual Soluble Bags Are Not To Be Offered For Sale"**.

For 2-3/4 oz. Water Soluble packets, the following statements will also be used at the top of the front panel - **"TO REMOVE PACKETS, CUT ALONG DOTTED LINE AND OPEN RESEALABLE CLOSURE."**

-----"
and **"PACKAGED SPECIFICALLY FOR SPOT TREATMENT USING BACKPACK SPRAYERS AND
HAZCONS"**.

WATER SOLUBLE PACKET LABEL

(Front Panel of overwrap bags
except 2-3/4 oz. size)

RIVERDALE
TRIPLET® WATER SOLUBLE

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE PRODUCT LABEL ON CARTON FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-283

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BURR RIDGE, ILLINOIS 60527-0866

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	60.20%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	24.20%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	6.02%
INERT INGREDIENTS:	<u>9.58%</u>
TOTAL	100.00%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	50%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	20%
***3,6-Dichloro-o-anisic Acid	5%

Patent No. 5,266,553

Exposure to moisture prior to mixing may cause water soluble bag to rupture.

Individual soluble bags are not to be offered for sale.