

228-288

9-29-2000

1/6

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98



United States  
Environmental Protection Agency  
Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

257281

Application for Pesticide - Section I

1. Company/Product Number 228-288	2. EPA Product Manager J. I. Miller (23)	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Riverdale Triplet® MC SAF	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Riverdale Chemical Company 425 West 194th Street Glenwood, IL 60425-1584 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: <b>NOTIFICATION</b> EPA Reg. No. _____ SEP 29 2000 Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
Changing official product name to Riverdale Triplet® Sensitive

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes * <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted	If "Yes" Unit Packaging wgt. _____ No. per container _____	If "Yes" Package wgt. _____ No. per container _____		<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product			
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Russell F. Sawyer	Title Regulatory Affairs Manager	Telephone No. (Include Area Code) 708/754-3330
2. Signature 		3. Title Regulatory Affairs Manager
4. Typed Name Russell F. Sawyer		5. Date July 10, 2000
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)

2/16

RIVERDALE

TRIPLET® SENSITIVE

SELECTIVE BROADLEAF WEED CONTROL FOR TURFGRASSES  
TO CONTROL DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND  
MANY OTHER BROADLEAF WEEDS

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

GET THE OPTICAL ADVANTAGE

NOTIFICATION  
SEP 29 2000

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	10.86%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	18.91%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	4.62%
INERT INGREDIENTS:	65.61%
TOTAL	100.00%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	9.02%	0.82 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	15.63%	1.43 lbs./gal.
***3,6-Dichloro-o-anisic Acid	3.84%	0.35 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

Triplet® Is A Registered Trademark of Riverdale Chemical Company  
THE OPTICAL ADVANTAGE™ is a Trademark of Riverdale Chemical Company

For Sale To And Use By Professional Personnel Only. Not For Sale To Homeowners.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que  
se la explique a usted en detalle.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-288


NET CONTENTS GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY  
RIVERDALE CHEMICAL COMPANY  
GLENWOOD, ILLINOIS 60425-1584

Note: Spanish Language optional.

Revised 10/23/98 Converted MCPP to Optical Isomer. Corrected 1/14/99  
7/10/00 Changed official name (via notification)



## (Side Panels)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive, causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed, absorbed through skin or inhaled. Wear face shield or goggles and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention at once.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water. If irritation develops, get medical attention.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Triplet<sup>®</sup> Sensitive is for use on Lawns, Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**USE PRECAUTIONS**

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Triplet Sensitive Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

**WEEDS CONTROLLED**

- |             |               |                |
|-------------|---------------|----------------|
| Bedstraw    | Henbit        | Purslane       |
| Black medic | Knotweed      | Ragweed        |
| Buckhorn    | Lambsquarters | Sheep sorrel   |
| Burdock     | Lespedeza     | Shepherdspurse |
| Chicory     | Mallow        | Speedwell      |
| Chickweed   | Morningglory  | Spurge         |
| Clover      | Peppergrass   | Wild carrot    |
| Dandelion   | Pigweed       | Wild garlic    |
| Dock        | Plantain      | Wild lettuce   |
| Ground ivy  | Poison ivy    | Wild onion     |
| Healall     | Poison oak    | Yarrow         |

and other broadleaf weeds.

**ORNAMENTAL LAWNS AND TURF**

**APPLICATIONS:**

Cool Season Grasses -

**RATES:**

Use this product at the rate of 3 to 4 pints per acre (1.5 to 2.0 fluid ounces per 1,000 square feet).

For Normal Application - Use 40 to 80 gallons of water per acre (1 to 2 gallons of water per 1,000 square feet).

For Professional Lawn Maintenance - Use 1 to 5 gallons of water

per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses - St. Augustine and Centipede - Apply Triplet Sensitive at a rate of 1 to 1-1/2 pints in 40 to 200 gallons of water per acre. Repeat after 3 weeks if necessary. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on these grasses. St. Augustine and Centipede Turfgrasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.

For Bahia, Common Bermuda, and Zoysia Grasses - Apply Triplet Sensitive at a maximum rate of 2 to 2-1/2 pints in 40 to 200 gallons of water per acre. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

#### NOTE

Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

Herbi™ Controlled Droplet Applicator: For cool season grasses - Add 2 to 3 pints of Triplet Sensitive to the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 square feet (approx. 3/4 acre). Do not overlap spray patterns.

For listed warm season grasses - Add 1.0 to 2 pints of Triplet Sensitive to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Other Controlled Droplet Applicators: (CDA), Atomizers, and Spinning disk Applicators: For cool season grasses - Use Triplet Sensitive at the rate of 3 to 4 pints per acre (1.1 to 1.5 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use 1 to 2-1/2 pints of Triplet Sensitive per acre (.4 to .9 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume Equipment Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface. Take special care to prevent overdosing.

Herbi™ is a trademark of North American Micron.

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY**

Riverdale Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(ERP 102393) (PR938&11 042194/RP 102396A)

Additional Approved (Optional) Uses:

**Bentgrass Turf (Putting and Bowling Greens):** On closely mowed bentgrass, apply this product at a maximum rate of 1 fluid ounce in 5 gallons of water per 1,500 square feet - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.