
ବର୍ଷ-କଥ

d instructions on reverse before completing form. Copy to Maurian Shedting golded. OMB No. 2070-0060

≎EPA	Un Environmental	nited States	n Agency		Registra Amenda Other	ation	OPP Identifier Number
	F	Application	ı for Pesticio	de - Sect	ion I		
1. Company/Product Numb 228-288	ier			Product Mana e Miller	ager	3. Pr	roposed Classification
4. Company/Product (Name Triplet Sensitive Selecti			PM# 23				J
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 150 Harvester Drive Suite 200 Burr Ridge, IL 60527		oj.	6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.				
X Check if th	his is a new address		 	ct Name _			
	/		Section - I	<u> </u>		 	
Amendment - Explain Resubmission in res	sponse to Agency letter d	fated		Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.			
PR Notice 95-2 and 98-10 statement of formula of the	ent with PRN 95-2 and 98 0 and EPA regulations at nis product. I understand diffication is not consistent	8-10, see cover t 40 CFR 152.40 I that it is a viola t withe terms of	r letter for detailed 6, and no other d ation of 18 U.S.C f PR Notice 95-2	changes have c. Sec. 1001 t and 98-10 an	e been made to th to willfully make fa nd 40 CFR 152.46	e labeling alse stater	
			Section - II	Ī			
1. Meterial This Product W	/ill Be Packaged In:						
Child-Resistant Peckaging Yes No * Certification must	Unit Packaging Yes No If "Yes" Unit Packaging wgt.		Yes No If "Yes" Package wgt	No. per	2. Type of	Metal Plastic Glass Paper	Specify)
be submitted				1		<u></u>	
3. Location of Net Contents Label	s Information 4 Container	4. Size(s) Retail	Container	Container 5. Location of Label Directions			ond .
6. Menner in Which Label is	Affixed to Product	Lithograp Paper plu Stenciled	ph ued d	Other			
			Section - IV	<i></i>			
1. Contact Point (Complete	e items directly below for	r identification c	of individual to be	contected, i	if necessary, to pr	ocess this	application.)
Name George Meindl george.meindl@us.nufarm.com			I				e No. (Include Area Code) 55.2017
i certify that the state i acknowledge that a both under applicable	ements I have made on th iny knowlinglly false or mi s law.	Certificatio his form and all hisleading staten	attachments the	reto are true, ishable by fin	, accurate and cor ne or imprisonmen	mpiete. t c <i>i</i>	6. Date Application Recolved {Stamped}
2. Signature			з. ти• Registration Manager			·	1
4. Typed Nam		5, (Date				
George Meindl		\ \footnote{2}	3/24/2006			Į	



NOTIFICATION
APR 2 8 2006

TRIPLET® SENSITIVE

Selective Broadleaf Weed Control For Turfgrasses
To Control Dandelion, Clover, Henbit, Plantains, Wild Onion,
and Many Other Broadleaf Weeds
Contains 2,4-D, Mecoprop-p and Dicamba

GET THE OPTICAL ADVANTAGE®

ACTI'	VE IN	GRED	ENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡		
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-anisic Acid)***		
OTHER INGREDIENTS:	6	5.61%
TOTAL:	100	0.00%
Isomer Specific AOAC Method, Equivalent to:		
*2,4-Dichlorophenoxyacetic Acid	9.02%, 0.82 lb	s./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	15.63%, 1.43 lb	s./gal.
***3,6-Dichloro-o-anisic Acid	3.84%, 0.35 lb	s./gal.
+CONTAINS THE SINGLE ISOMER FORM OF MECOPROP.0		_

FOR SALE TO AND USE BY PROFESSIONAL PERSONNEL ONLY.
NOT FOR SALE TO HOMEOWNERS.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

SEE SIDE PANELS FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

Note: Spanish language optional.

EPA REG. NO. 228-268 EPA EST. NO. 228-IL-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



NET CONTENTS

GALS.

000228-00288.20060323.clean

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive, causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed, absorbed through skin or inhaled. Wear face shield or goggles and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

	FIRST AID
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
	HOT LINE NUMBER ntainer or label with you when calling a poison control center or doctor, or going for treatment. t 1-877-325-1840 for emergency medical treatment information.
Probable mucosal da	NOTE TO PHYSICIAN mage may contraindicate use of gastric lavage.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemereries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray culutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except



as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only.

Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Bedstraw	Ground ivy	Pigweed	Spurge
Black medic	Healall	Plantain	Wild carrot
Buckhorn	Henbit	Poison ivy	Wild garlic
Burdock	Knotweed	Poison oak	Wild lettuce
Chicory	Lambsquarters	Purslane	Wild onion
Chickweed	Lespedeza	Ragweed	Yarrow
Clover	Mallow	Sheep sorrrel	and other broadleaf weeds.
Dandelion	Morningglory	Shepherdspurse	
Dock	Peppergrass	Speedwell	

ORNAMENTAL LAWNS AND TURF

APPLICATIONS:	RATES:
Cool Season Grasses	Use this product at the rate of 3 to 4 pints per acre (1.1 to 1.5 fluid ounces per 1,000 square feet).
	For Normal Application - Use 40 to 80 gallons of water per acre (1 to 2 gallons of water per 1,000 square feet.)
	For Professional Lawn Maintenance - Use 1 to 5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.
Warm Season Grasses	St. Augustine and Centipede - Apply this product at a rate of 1 to 1-1/2 pints in 40 to 200 gallons of water per acre. Repeat after 3 weeks if necessary. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on these grasses. St. Augustine and Centipede Turfgrasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.
	For Bahia, Common Bermuda, and Zoysia Grasses - Apply this product at a maximum rate of 2 to 2-1/2 pints in 40 to 200 gallons of water per acre. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.
	NOTE: If Bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.
Bentgrass Turf (Putting and Bowling Greens)	On closely mowed bentgrass, apply this product at a maximum rate of 1 fluid ounce in 5 gallons of water per 1,500 square feet - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE

Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and

subsequent turf discoloration or injury.

Herbi™ Controlled Droplet Applicator	For cool season grasses - Add 2 to 3 pints of this product to Droplet Applicator: the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 square feet (approx. 3/4 acre). Do not overlap spray patterns. For listed warm season grasses - Add 1.0 to 2 pints of this product to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.
Other Controlled Droplet Applicators: (CDA), Atomizers, and Spinning disk Applicators	For cool season grasses - Use this product at the rate of 3 to 4 pints per acre (1.1 to 1.5 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns. For listed warm season grasses - Use 1 to 2-1/2 pints of this product per acre (.4 to .9 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.
Lower Volume Equipment	Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface. Take special care to prevent overdosing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably to the user assumes all risks of any such use.

(RV032306N)

TRIPLET and THE OPTICAL ADVANTAGE are Registered Trademarks of Nufarm Americas Inc. Herbi™ is a trademark of North American Micron.





Nufarm Americas, Inc. George Meindl Registration Manager 150 Harvester Drive, Suite 200 Burr Ridge, IL 60527 Phone: 630.455-2017 Fax: 630.455.2030 george.meindl@us.nufarm.com

Via Overnight Courier

March 24, 2006

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 Bell Street
Arlington, VA 22202-4501

Subject:

Triplet® Sensitive Selective Herbicide

EPA Reg. No. 228-288

Label notification per PR98-10 and PR95-2

Dear Ms. Hobgood:

Nufarm Americas, Inc would like to make label changes to the subject registration. We feel that these label changes are permitted and consistent with PRN95-2 section II (Notifications) part A (Labeling) subsection 4 (Revising Advisory Statements) and PRN98-10 section II (Labeling Notifications) part D (Adding Advisory Statements).

Specifically we have replaced the following sentence "Do not spray when air temperature exceeds 90°F" with the following 2 sentences "Avoid broadcast applications when air temperature exceeds 90 degrees. When using small spot treatment applications in temperatures over 90 degrees, turf injury may occur." We have also added in a separate paragraph with the following two advisory statements: "When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only."

The applicators of this product have told us the current wording was incomplete and confusing. We feel the suggested changes gives the applicator a clearer understanding on how to use this product in the most efficacious manner possible. These changes also support Nufarm's product stewardship initiatives and overall responsible care efforts.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling clean (1 copy)

If you should have any questions regarding this matter, please feel free to contact me at 630.455.2017 or email at qeorge.meindl@us.nufarm.com.

Sincerely, A

George Meindl (\)
Registration Manager
Nufarm Americas, Inc.

H:\228-288\label notification per PRN95-2 and 98-10 submitted 3.24.2006.doc