


228-296

8/10/2009

Jacket
1/10



U.S. ENVIRONMENTAL PROTECTION
AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: 228-296
Date of Issuance: AUG 10 2009

NOTICE OF PESTICIDE:
 Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:
Name of Pesticide Product:
Veteran 2010 Herbicide

Name and Address of Registrant (include ZIP Code):
Nufarm Americas Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Change the Signal word to "WARNING" and "Peligro" to "AVISO".

Signature of Approving Official:

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

AUG 10 2009

3) To "SPRAY DRIFT MANAGEMENT" add

"Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions."

"Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles."

"Apply only when the wind speed is 2-10 mph at the application site."

Delete conflicting text

4) The text "Do not apply more than 6.4 pints of product (.99 lbs ae Dicamba, 2 lbs ae MCPA) per application ..." appearing on Page 3 of the label allows rates in excess of the maximum allowable rate of 1.5 lbs ae per acre per application for MCPA. The label must be revised.

5) To page 4, NonCropland Use add "Do not apply more than 1.5 lb ae MCPA / acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

6) To page 1, specify "similar noncropland areas" same on page 4. Reduce the rates on page 5 to "Do not apply more than 1.5 lb ae MCPA / acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days." For all uses.

7) Revise the Tank Mix Rates on page 5, simazine to 4 lbs and tebuthiuron to 6 lbs. On page 6, change "Do not exceed 16 lbs." to "Do not exceed 3 lbs."

You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supercedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

VETERAN® 2010 HERBICIDE

ACCEPTED
with COMMENTS
In EPA Letter Dated:

AUG 10 2009
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-296

WATER SOLUBLE INDUSTRIAL HERBICIDE FOR CONTROL OF BRUSH AND
BROADLEAF WEEDS ON RIGHTS-OF-WAY, RAILROADS, PIPELINES, FOREST
BRUSH, FENCE ROWS, NON-IRRIGATION DITCHBANKS, WASTELAND,
GENERAL FARMSTEAD, AND SIMILAR NONCROPLAND AREAS. ALSO FOR
PASTURES AND RANGELANDS.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	32.6%
Dimethylamine salt of Dicamba (3,6-Dichloro-o-Anisic Acid)**	16.0%
OTHER INGREDIENTS:	<u>51.4%</u>
TOTAL:	100.0%

BY ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	26.62%, 2.50 lbs./gal.
**Dicamba (3,6-Dichloro-o-Anisic Acid)	13.23%, 1.25 lbs./gal.

FOR COMMERCIAL USE ONLY

**KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

NOTE: Spanish language is optional


EPA REG. NO. 228-296
EPA EST. NO.

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



Net Contents Gal. (L)

000228-00296.20090511.EPA.Dicamba RED.Pending



4/10

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER / PELIGRO**

Causes substantial but temporary eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants,
- protective eyewear (goggles, face shield or safety glasses),
- chemical-resistant gloves (made of any water-proof material), and
- shoes plus socks.

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland. These mixers/loaders must also wear: a chemical-resistant apron, and a NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-84A. It is recommended that you require that respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See engineering controls for additional requirements.

Engineering Control Statements:

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)].

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN
Probable mucosal damage may contraindicate the use of gastric lavage. Chlorophenoxy compounds are irritating to eyes, skin and mucous membranes.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

5/10

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Not for use on (1) turf being grown for sale or other commercial use as sod; (2) trees being grown for sale or other commercial use, or for the production of timber or wood products; (3) turf, trees, crops, timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply more than 6.4 pints of product per application and no more than 2 applications per year.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or green houses.

Do not allow people (other than applicator) or pets on treatment area during application.

Do not enter treatment areas until sprays have dried.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

SENSITIVE CROP PRECAUTIONS

IMPORTANT: Do not apply this product directly to or allow spray drift to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, sugar beets, beans, or other desirable plants. Coarse sprays are less likely to drift. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not apply, except as recommended, or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

SHAKE WELL BEFORE USING.

NONCROPLAND USE: (Roadsides, fence rows, pastures, rangelands, railroads, brush control for forest site preparation and similar areas.) For top growth control of field bindweed, Canada thistle, perennial sowthistle, Russian knapweed and leafy spurge, apply 3.2 to 4.75 pints of this product. For spot treatment, apply 1 teaspoonful of this product per 275 to 550 square feet. Apply as a foliar spray to actively growing plants.

RESTRICTIONS AND LIMITATIONS FOR USE ON NON-CROPLAND RIGHTS-OF-WAY:

Do not apply more than 4.75 pints of product per acre per year.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

BRUSH AND BROADLEAF WEED CONTROL

SPECIES CONTROLLED

When used as directed, this product will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as suppress many unwanted woody plant and vine species. Species controlled include:

ANNUALS

Buckwheat	Knawl	Purslane
Carpetweed	Kochia	Ragweed
Chickweed	Lambsquarter	Russian Thistle
Clover	Morningglory	Smartweed
Cocklebur	Mustard	Velvetleaf
Daisy, English	Pigweed	
Henbit	Puncturevine	

BIENNIALS

Ragwort, Tansy	Thistle, Musk
----------------	---------------

PERENNIALS

Bindweed, Field	Dogfennel	Sorrel, Sheep
Carrot, Wild (Queen Anne's Lace)	Knapweed, Russian	Spurge, Leafy
Dock, Curly	Milkweed	Thistle, Canada
	Ragweed, Perennial	Toadflax, Dalmatian

WOODY BRUSH AND VINES*

Alder	Honeysuckle	Sassafras
Ash	Ivy, Poison	Schinus
Aspen	Kudzu	(Florida Holly,
Basswood	Locust	Brazil Peppertree,
Beech	Maple	Christmas-berry)
Birch	Oak	Serviceberry
Blackberry	Olive, Russian	Snowberry
Cherry	Persimmon	Spruce
Creeper, Virginia	Pine	Sumac
Creosotebush	Plum, Wild	Sycamore
Cucumber tree	Poplar	Trumpet creeper
Dogwood	Puncturevine	Waxmyrtle
Elderberry	Raspberry	Willow
Elm	Redcedar, Eastern	Witchhazel
Gum	Redvine	Yaupon
Hawthorn	Rose, Multiflora	
Hemlock	Sagebrush	

*Suppression

APPLICATION DIRECTIONS

Regardless of the species to be controlled, spray volumes should be high enough to allow for good spray coverage. Make applications when weeds and brush are actively growing. The addition of surfactants¹ can increase control. A drift control agent² may also be used. Biennials are best controlled when treated in the rosette stage. Regrowth may occur on resistant species. Retreatments may be made at least 30 days after the previous application; however, do not exceed a total of 0.8 gallon (1 lb. a.i. dicamba + 2 lbs. a.i. MCPA) of this product per treated acre during a growing season. For control of woody species, a tank mix may be necessary. Tank mix recommendations are for use only in states where the tank mix product and application site are registered. Read and follow all label directions.

HERBACEOUS BROADLEAF WEED CONTROL

Apply 3/4 to 3-1/4 quarts of this product in 5 to 100 gallons of water per treated acre. 3/4 to 2 quarts of this product is recommended for annuals, 2 to 3-1/4 quarts for biennials and easy-to-kill (or control) perennials, and 3-1/4 quarts for established perennials. Do not apply more than 1.6 gallons of product per treated acre per year.

WOODY BRUSH AND VINE CONTROL

High Volume Foliar Spot Applications: Mix 0.8 gallon of this product in enough water to make 100 gallons of spray mix. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 1.6 gallons of product per treated acre per year. Direct the spray to treat all foliage, stems, and root collars to wet.

WOODY BRUSH AND VINE CONTROL/SUPPRESSION

Broadcast Applications with Ground Equipment: Apply 0.8 gallon of this product in 20 to 100 gallons of water per treated acre. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 1.6 gallons of product per treated acre per year. Spray all foliage, stems, and root collars to wet. For optimum control of woody species, a tank mix may be necessary.

AERIAL APPLICATIONS

Aerial applications may be made to control either herbaceous or woody plants. Apply 1 to 3-1/4 quarts of this product (for herbaceous weeds) or 0.8 gallon of this product (for woody brush and vines) in 5 to 40 gallons of water per treated acre. Coverage is important, so increase spray volume when treating dense stands of brush or weeds. Do not apply more than 1.6 gallons of product per treated acre per year. For optimum control of woody species, a tank mix may be necessary.

TANK MIX TREATMENTS

TANK MIX RECOMMENDATIONS ARE FOR USE ONLY IN STATES WHERE THE TANK MIX PRODUCT AND APPLICATION SITE ARE REGISTERED. READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, this product may be tank mixed with other herbicides registered for noncropland uses (e.g. railroad, highway, pipeline, pastures, rangeland, forest brush control, etc.), including but not limited to the following herbicides. Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

<u>HERBICIDE</u>	<u>RATE</u>	
	<u>(LBS. A.I./TREATED ACRE)</u>	
Asulam (Asulox®)	2.92	to 6.68
Bromacil (Hyvar®)	1.60	to 12.00
Chlorsulfuron (Telar®)	0.1875	to 2.25 oz.**
Dicamba (Diablo™)	0.25	to 1.00***
Diuron (Karmex®)	4.00	to 12.00
Fosamine ammonium (Krenite®)	6.00	to 12.00
Glyphosate (Razor®)	0.75	to 3.75
Hexazinone (Velpar®)	0.675	to 10.80
flmazapyr (Arsenal®)	0.50	to 1.50
Metsulfuron Methyl (Patriot®)*	0.30	to 0.90**
MSMA	1.00	to 2.475
Picloram* (Tordon®)	0.50	to 1.00
Simazine* (Princep®)	4.80	to 40.00
Sulfometuron methyl (Spyder™)	0.75	to 9.00 oz.**
Tebuthiuron (Spike®)	1.00	to 16.00
Triclopyr (Tahoe™)	0.75	to 9.00
2,4-D	0.475	to 4.00****
2,4-DP-p	0.50	to 0.75

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST as described on this label is recommended prior to actual tank mixing.

* Restricted use pesticides limited to certified applicators

- ** Telar = 0.1875 to 2.25 oz. a.i./A (0.25 to 3.0 oz. product/A)
Patriot = 0.3 to 0.9 oz. a.i.(0.5 to 1.5 oz product/A)
Spyder = 0.75 to 9.0 oz. a.i./A (1.0 to 12.0 oz. product/A)
- *** Do not exceed a total of 1 lb. a.i. dicamba per treated acre per growing season. (This product contains 1.25 lbs. a.i. dicamba per gallon).
- **** Use of extremely hard water (500 ppm) may cause this product to form a precipitate when used in a tank mix with Veteran 2010 Herbicide. Do not exceed 16 lbs. total MCPA acid equivalent per treated acre per growing season. (Veteran 2010 Herbicide contains the equivalent of 2.5 lbs./gal. MCPA acid).
- † Not registered for use in California
- a Using this product and Patriot to reduce the development and spread of resistant biotypes and problem weeds such as Kochia and Russian thistle: some commonly resistant weeds typically require multiple spray applications to obtain adequate control. To reduce the number of applications required, applicators can utilize a mixture of Patriot and this product. Apply as a tank mix for post-emergence as follows: Add 5 ounces of Patriot to 1 quart of this product per acre. A non-ionic or silicone surfactant may be used for wetting, penetration or drift control.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**AMOUNT OF HERBICIDE TO ADD TO
ONE PINT OF SPRAY CARRIER
(ASSUMING VOLUME IS 25 GALLONS PER ACRE)**

<u>HERBICIDE FORMULATIONS</u>	<u>RATE PER ACRE</u>	<u>LEVEL TEASPOONS</u>
Dry	1 pound	1-1/2
Liquid	1 pint	2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested components are compatible. Usually incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of spray solution).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of this product or tank mixes of this product plus MCPA amine.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferable overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply this product as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. This product tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- 5) Complete step 1.
- 6) Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 7) Flush the detergent solution out of the spray tank through the boom.
- 8) Repeat step 1, and follow with steps 2, 3 and 4.

9/2

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperature above 32°F. If allowed to freeze, warm to at least 40°F and remix before using. Freezing does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: [Nonrefillable Containers 5 Gallons or Less:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Nonrefillable Containers Larger than 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable Container Larger than 5 Gallons:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV051109R)

Veteran, Diablo, Patriot, Razor, Spyder, and Tahoe are registered trademarks of Nufarm Americas Inc. All other trademarks are the property of their respective owners.

10
/
10

Optional Brand Names and Cleaning Instructions

¹Such as R-11®, Syl-tac™, Hasten®, or Mor-Act®

²Such as Bivert®, or Redi-Vert

³The addition of a tank cleaner, such as Neutral-Clean spray tank cleaner and neutralizer. In a 100 to 300 gallon spray tank, add approximately 50 gallons of water and 1/2 pound of Neutral-Clean. Agitate and circulate for approximately 15 minutes. Read and follow complete Neutral-Clean label instructions.