

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

	United States	<input type="checkbox"/> Registration	OPP Identifier Number <b>228041</b>
	<b>Environmental Protection Agency</b>	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

**Application for Pesticide - Section I**

1. Company/Product Number 228-299	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Riverdale Tri-Power (R) 4000 Weed and Feed	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Riverdale Chemical Company 425 West 194th Street Glenwood, Illinois 60425-1584 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Adding Fertilizer & Marketing hype.  
Also corrected Commercial Label which was missing groundwater statement and incorrect Storage and Disposal Statement.

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Russell F. Sawyer	Title Regulatory Affairs Manager	Telephone No. (Include Area Code) 708/754-3330
2. Signature 		6. Date Application Received (Stamped)
3. Title Regulatory Affairs Manager		
4. Typed Name Russell F. Sawyer		
5. Date July 13, 1995		

(Commercial Label)

*Extra cost  
with Highland  
Chicago 2 9 9*

RIVERDALE

TRI-POWER® 4000  
WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD  
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF  
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 4,000 Sq. Ft. (40' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	. . . . .	.646%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	. . . . .	.255%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	. . . . .	.064%
INERT INGREDIENTS:	. . . . .	<u>94.035%</u>
	TOTAL . . . . .	100.000%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid . . . . .	0.528%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid . . . . .	0.211%
***3,6-Dichloro-o-Anisic Acid . . . . .	0.053%

GUARANTEED ANALYSIS:°

Total Nitrogen (N) . . . . .	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	X%
Soluble Potash (K <sub>2</sub> O) . . . . .	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum . . . . . X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

†TRI-POWER® - is a Registered Trademark of Riverdale Chemical Company

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.

EPA REG. NO. 228-299

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Causes moderate eye irritation, harmful if absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Wear long pants, long sleeved shirt and shoes when applying this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not re-enter or permit workers to re-enter areas until dust has settled.

## STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.  
 IF IN EYES: Flush eyes with plenty of water. Call physician if irritation persists.  
 IF ON SKIN: Wash promptly with plenty of soap and water and rinse thoroughly. If irritation develops, get medical attention.

## ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate, nor pour the washwaters on the ground; spray or drain over a large area away from well, other water sources and desirable plants.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRI-POWER® WEED AND FEED is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, tree or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggartick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dogfennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbert-leaved saltbush, Hawkweed, Healall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter, Mallow Marshelder, Mexican poppy, Morningglory, Mustard, Nettle, Oxalis, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanish needle, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow charlock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or

early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application, (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 4000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 5 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader of 200 sq. ft. and adjust to apply at the rate of 1 lb per 200 sq. ft.

#### SPREADER SETTINGS

ONE PASS APPLICATION:°

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1000 sq. ft.; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

#### STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feeds, fertilizers, insecticides or fungicides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local

5 7.9

authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(PR938 042194/RV 071395)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL ADVERTISING STATEMENT

CONTAINS Tri-Power® A Premium Broadleaf Herbicide

Environmental Benefits



Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

\*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

(Non-Commercial Label)

6 7 9  
EX. 7203  
with analytical  
changes

RIVERDALE

TRI-POWER® 4000  
WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD  
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF  
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEED ACTION.

Weed contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 4,000 Sq. Ft. (40' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	. . . . .	.646%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	. . . . .	.255%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	. . . . .	.064%
INERT INGREDIENTS:	. . . . .	99.035%
	TOTAL . . . . .	100.000%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid . . . . .	.528%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid . . . . .	.211%
***3,6-Dichloro-o-Anisic Acid . . . . .	.053%

GUARANTEED ANALYSIS:°

Total Nitrogen (N) . . . . .	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	X%
Soluble Potash (K <sub>2</sub> O) . . . . .	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum . . . . . X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

†TRI-POWER® - is a Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.

EPA REG. NO. 228-299

EPA EST. NO. 228-IL-1.

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Wear long pants, long sleeved shirt and shoes when applying this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow persons and pets on treated areas until dusts have settled.

## STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.  
 IF IN EYES: Flush eyes with plenty of water. Call physician if irritation persists.  
 IF ON SKIN: Wash promptly with plenty of soap and water and rinse thoroughly. If irritation develops, get medical attention.

## ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRI-POWER® WEED AND FEED is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggartick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dogfennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbert-leaved saltbush, Hawkweed, Healall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter, Mallow, Marshelder, Mexican poppy, Morningglory, Mustard, Nettle, Oxalis, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanishneedle, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow charlock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers, especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture

conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 4000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 5 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce settings for slower speed; raise settings for higher speed. Check your spreader on 200 sq. ft. and adjust to apply at the rate of 1 lb. per 200 sq. ft.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE	SETTINGS
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1000 sq. ft.; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert Fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(PR938 042194/RV 071395)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.



(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL ADVERTISING STATEMENT

CONTAINS TRI-POWER® A Premium Broadleaf Herbicide

# Environmental Benefits



Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

\*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.