

1023

228 301

1051

(Commercial Label)

RIVERDALE

TRI-POWER® DRY WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENTS:*†

2-Methyl-4-Chlorophenoxyacetic Acid	0.63%
2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.25%
Dicamba (3,6-Dichloro-o-Anisic Acid)	0.06%
INERT INGREDIENTS:	<u>99.06%</u>
TOTAL	100.00%

*ISOMER SPECIFIC AOAC METHOD

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer registrant)

†TRI-POWER® - A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT LBS.


EPA REG. NO. 228-301

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.
- IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.
- IF ON SKIN:** Wash promptly with plenty of soap and water and rinse thoroughly. Get medical attention.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters or rinsate.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRI-POWER® DRY WEED AND FEED is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and to control such broad-leaf weeds as: Beggartick, Bindweed, Black medic, Buckhorn, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, English daisy, Fat hen, Fennel, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbertleaved saltbush, Hawkweed, Healall, Heartleaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter, Mallow, Marshelder, Mexican poppy, Morningglory, Mustard, Nettle, Oxalis, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Redstem, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sow thistle, Spanish-needle, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Toadflax, Veronica, Vetch, Whitebrush, White top, Wild aster, Wild carrot, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild onion, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Wood sorrel, Yarrow, Yellow charlock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 (or 18) pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds (3.6 pounds for 18# bag) per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. (.9 lb. for 18# bag) per 250 sq. ft.

SPREADER SETTINGS

ONE PASS APPLICATION:°

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1000 sq. ft.; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Iherf fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feed or foodstuffs. Do not stack more than two pallets high. To prevent cross-contamination, do not store near open containers of fertilizers, seeds or other pesticides. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(PR938 042194/RV 091995)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS:

Contains Tri-Power® Dry a Premium Broadleaf Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

