



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
228-302

Date of Issuance:
SEP 29 1995

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Riverdale Triplet
L.A. Selective
Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Company
425 West 194th Street
Glenwood, IL 60425-1584

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. "228-302".
 - b. Revise the statement under Hazards to Humans and Domestic Animals which begins "Harmful if swallowed ..." to read as follows: "Harmful if swallowed, inhaled or absorbed through skin."
 - c. Delete the phrase "if irritation develops" from the dermal statement of practical treatment.

Signature of Approving Official:

Date:

SEP 29 1995

287

page 2
EPA Reg. No. 228-302

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

RD:STANTON:PM Team 23:Rm. 235:CM-2:305-5218:Disk #1:228_302.REG

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	S. Stanton							
DATE ▶	Sep 26, 1995							

387

RIVERDALE

TRIPLET® L.A.
SELECTIVE HERBICIDE

Selective Broadleaf Weed Control For Turfgrasses
To Control Dandelion, Clover, Henbit, Plantains, Wild Onion, and
Many Other Broadleaf Weeds.
Also for Highways, Rights-of-Way and Other Similar Non-Crop Areas

Contains 2,4-D, Mecoprop and Dicamba

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	27.6%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	13.9%
Dimethylamine Salt of Dicamba(3,6-Dichloro-o-anisic Acid)***	2.7%
INERT INGREDIENTS:	55.8%
TOTAL	100.0%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	22.9%	2.20 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	11.4%	1.10 lbs./gal.
***3,6-Dichloro-o-anisic Acid	2.2%	0.22 lbs./gal.

Triplet® Is A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

**SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT**

EPA REG. NO. 228-GNE

NET CONTENTS GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

NOTE: Spanish language is optional

ACCEPTED
with COMMENTS
In EPA Letter Dated
SEP 29 1995
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-302

(Inside Booklet)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER - PELIGRO

Corrosive, causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed or inhaled. Wear face shield, goggles or safety glasses and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements and Directions For Use instructions.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention at once.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water and rinse thoroughly. If irritation develops, get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LA-

BEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Triplet L. A. Selective Herbicide is for use on Lawns, Turf, Golf Courses, (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

.....

ADDITIONAL PRECAUTIONS FOR INDUSTRIAL APPLICATIONS

When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands, face and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Triplet L. A. Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Bedstraw	Dock	Morningglory
Black medic	Ground ivy	Peppergrass
Buckhorn	Healall	Pigweed
Burdock	Henbit	Plantain
Chicory	Knotweed	Poison ivy
Chickweed	Lambsquarter	Poison oak
Clover	Lespedeza	Purslane
Dandelion	Mallow	Ragweed
Sheep sorrel	Spurge	Wild lettuce
Shepherdspurse	Wild carrot	Wild onion
Speedwell	Wild garlic	Yarrow

and other broadleaf weeds.

ORNAMENTAL LAWNS & TURF

APPLICATIONS:

RATES:

Cool Season Grasses - Use this product at the rate of 3-1/3 to 4-1/2 pints per acre (1.2 to 1.6 fl. oz. per 1000 sq. ft.).

For Normal Application - Use 40 to 80 gallons of water per acre. (1 to 2 gallons water per 1,000 square feet)

For Professional Lawn Maintenance - Use 1 to 2 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses - Such as St. Augustine, Bahia, Common Bermuda, Centipede, and Zoysia grasses - Apply Triplet L. A. Selective Herbicide at a maximum rate of 2-1/4 to 2-4/5 pints in 40 to 200 gallons of water per acre. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: If Bermudagrass is dormant, up to 4-1/2 pints per acre may be used. However, some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Bentgrass: On closely mowed bentgrass (putting and bowling greens), apply this product at a maximum rate of 1-1/8 fluid ounce in 5 gal. water per 1,500 square ft. - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE

Care should be taken to avoid overdosing bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. 1-1/8 fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

Herbi™ Controlled Droplet Applicator: For cool season grasses - add 2-1/4 to 3-1/3 pints of Triplet L. A. Selective Herbicide to the Herbi 5 pint container. Fill balance of container with water. Keeping container agitated, spray entire contents over 33,000 sq. ft. (approx. 3/4 acre). Do not overlap spray patterns.

For listed warm season grasses - add 1-2/3 to 2-1/4 pints of Triplet L. A. Selective Herbicide to the Herbi 5 pint container and fill balance of container with water. Follow Herbi spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Other Controlled Droplet Applicators: (CDA), Atomizers, and Spinning Disk Applicators: For cool season grasses - Use Triplet L. A. Selective Herbicide at the rate of 3-1/3 to 4-1/2 pints per acre (1.2 to 1.6 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1 to 4 gal. water per acre is normal for this type of equipment). Do not overlap spray patterns.

For listed warm season grasses - Use 2-1/4 to 2-4/5 pints of Triplet L. A. Selective Herbicide per acre (.84 to 1.0 fl. oz. per 1000 sq. ft.) in sufficient water to assure coverage (1 to 4 gal. of water per acre is normal for this type equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.

Lower Volume Equipment Use as little as 5 gallons water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting

each weed surface.

Herbi™ is a trademark of North American Micron

NON-TURF AREAS

Roadsides (including aprons and guardrails) rights-of-way, and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 3 pints to 1-1/8 gallon of Triplet L. A. Selective Herbicide per 50 to 300 gallons of water. This mixture will cover 43,500 sq. ft. Thoroughly saturate all weeds with spray mixture. Apply anytime between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4-1/2 ounces Triplet L. A. Selective Herbicide per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage anytime from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 1-1/8 to 1-2/3 gallons of Triplet L. A. Selective Herbicide to 100 gallons of water applying 200 to 600 gallons of spray mixture per 43,500 sq. ft. depending upon the height and thickness of the brush. Mix thoroughly before spraying.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(ERP 102393) (PR938&11 042194)