

228-309

07/23/2003

1/4

RIVERDALE®

VETERAN® 10G

GRANULAR HERBICIDE

FOR NONCROPLAND AREAS
SUCH AS FENCE ROWS, ROADWAYS, WASTELAND, GENERAL FARMSTEAD,
AND SIMILAR NONCROPLAND. ALSO FOR PASTURES AND RANGELAND

ACTIVE INGREDIENTS:

Dicamba (3,6-dichloro-o-anisic acid)	10.0%
INERT INGREDIENTS	90.0%
TOTAL	100.0%

Riverdale and Veteran are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

WARNING



**SEE SIDE PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA REG. NO. 228-309

NET CONTENTS LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****WARNING**

Harmful if swallowed. Avoid breathing vapors. Dust may cause eye irritation. Avoid contact with skin, eyes and clothing.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
- IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate irrigation ditches or water used for domestic purposes. Do not contaminate water by the cleaning of equipment or disposal wastes. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Veteran® 10G Herbicide may cause injury to desirable trees or plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems, or foliage.

Do not treat areas where the possible downward movement into the soil or surface washing may cause contact of herbicide with the roots of desirable plants such as trees and shrubs.

To avoid injury to desirable plants, equipment used for VETERAN 10G Herbicide should be thoroughly cleaned before reuse to handle or apply any other chemicals.

VETERAN 10G Herbicide may temporarily stunt or eliminate the growth of grass in the treated areas.

Pasture and rangeland grasses may be grazed or harvested for feed (green or dry) for

lactating dairy animals 60 days following treatment. There is no waiting period between treatment and grazing for animals other than lactating dairy animals. Remove meat animals from treated areas 30 days prior to slaughter.

WEEDS AND BRUSH CONTROLLED

VETERAN 10G Herbicide, when applied in accordance with this label, will give control or growth suppression of many ANNUAL, BIENNIAL and PERENNIAL broadleaf weeds as well as many WOODY brush and vine species including:

ANNUAL BROADLEAF WEEDS

Black Nightshade	Pigweeds
Cocklebur	Prickly lettuce
Jimsonweed	Puncturevine
Kochia	Ragweed
Lambsquarter	Russian thistle
Morningglory	Sunflower
	Tansy mustard

BIENNIAL and PERENNIAL BROADLEAF WEEDS

Bindweed, Field	Milkweed
Blueweed, Texas	Musk thistle
Broom snakeweed	Nightshade, Silverleaf
Burdock	Plumeless thistle
Bursage, Skeletonleaf	Ragwort, Tansy
(Bur Ragweed, Povertyweed)	Sowthistle, Perennial
Bursage, Woolyleaf	*Spurge, Leafy
(Lakeweed)	Starthistle, Yellow
Chicory	Stinging nettle
Curly dock	Sweetclover
Dalmatian toadflax	Teasel
Diffuse knapweed	Thistle, Artichoke
Dogbane	Thistle, Bull
Dogfennel	Thistle, Canada
*Fern, Bracken	Tropical Soda Apple
Goldenrod	Trumpet creeper
Knapweed, Russian	Yarrow
Milk thistle	

*Noted weeds require higher rates of VETERAN 10G than other listed weeds. See APPLICATION RATES AND TIMING section of this label.

WOODY BRUSH and VINES

Alder	Hickory	Poison ivy
Ash	Honeylocust	Poison oak
Aspen	Honeysuckle	Rabbitbrush
Basswood	Hornbeam	Sagebrush
Beech	Huckleberry	Saltcedar
Black locust	Huisache	Sassafras
Birch	Kudzu	Serviceberry
Cherry	Maple	Spicebrush
Chinquapin	Mesquite	Spruce
Cottonwood	Oak	Sumac
Creosotebush	Olive, Autumn	Sycamore
Elm	Olive, Russian	Tarbrush
Grape	Persimmon, Eastern	Willow
Hemlock	Plum, Sand	Witchhazel

APPLICATION RATES AND TIMING: VETERAN 10G Herbicide may be applied using standard ground or air granular application equipment for broadcast or spot treatments. VETERAN 10G Herbicide is activated when there is sufficient moisture to move the herbicide into the root zone of plants to be controlled. Application may be made any time when soil is not frozen. However, best performance usually occurs when treatments are made in the Spring just before or during active growth of the weeds and brush. CONSULT YOUR LOCAL OR STATE AUTHORITIES FOR THE BEST TIMING OF APPLICATION.

Application rates are given below. Use the higher level of listed rate range when treating dense vegetative growth.

- For spot treatments, use 1/2 to 1 pound of product per 1,000 square feet for control of listed ANNUAL, BIENNIAL and PERENNIAL weeds.
- For spot treatments, use 1 to 2 pounds of product per 1,000 square feet for noted (*) ANNUAL, BIENNIAL and PERENNIAL weeds.
- For spot treatments, use 2 pounds of product per 1,000 square feet for control of listed WOODY BRUSH and VINES. Uniformly spread granules over the area where the plant roots are growing.

In most cases, these treatments will not control weed seedlings which germinate one to two years after treatment. Once the effect of the herbicide has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

For broadcast application, do not use more than 20 pounds of VETERAN 10G per treated acre per year.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a well-ventilated area away from fertilizer, feed or foodstuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) plastic pails. For paper or plastic bags, or for fiber drums with liners, completely empty into application equipment by shaking and tapping container. Offer pails or drums for recycling or reconditioning or puncture before disposal. Dispose of bags, liners, or punctured containers in sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer and the limit of liability of any such use. The exclusive remedy of user or buyer and the limit of liability of Seller is the purchase price paid for the quantity of product involved. (RV 051503)