11/19/2004

31 -1 2 - /1/8

Form Approved. OMB No. 2070-0060

≎EPA ■	460			Registration X Amendment Other		OPP Identifier Number		
1. Company/Product Number	Application	00 101 t	Pesticide - Sect	·		3 0	d Classi	Paskan
228-312	•		2. EPA Product Man Joanne Miller	eger		3, Proposed Classification X None Restricted		
4. Company/Product (Name) RiverdaleTriplet SF		PM# 23			X	None	Nestricted	
5. Name and Address of Application Nufarm Americas Inc. 1333 Burr Ridge Parkway, Burr Ridge, IL 60527 Check if this is						nd labeling		
		Sec	tion - II					
Resubmission in respons X Notification - Explain bel	Amendment - Explain below. Final printed labels in repsonse to Agency letter dated "Me Too" Application.							
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Addition of an alternate brand name - Primera Triplet SF. -								
		Sect	ion - III					
1. Material This Product Will Be	Packaged In:							
Yes No	Init Peckaging Yes No No. per Init Packaging wgt. container	If "Yes"	Weter Soluble Packaging 2. Type o Yes No If "Yes" No. per			Container Metal Plastic Glass Paper Other (Specify)		
be submitted	ubmitted Container Package wgt Container				ation of Lebel			
	Label Container							
Section - IV								
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)								
Name Jeannie Smith	Title VP Registrations & Regulatory Affairs Telephone No. (Include Area Code) 708 755-7332							
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete, I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 6. Date Ap, lication Received (Stamped)								
2. Signatuse	3. Title VP Registrations & Regulatory Affairs							
4. Typed Name	5. Date							
Jeannie Smith	November 15, 2004 .							

Primera Triplet® SF

NOTIFICATION

NOV 1 9 2004

SELECTIVE BROADLEAF WEED CONTROL FOR TURFGRASSES INCLUDING USE ON SOD FARMS. TO CONTROL CLOVER, DANDELION, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER BROADLEAF WEEDS. ALSO FOR HIGHWAYS, RIGHTS-OF-WAY AND OTHER SIMILAR NON-CROP AREAS.

CONTAINS 2,4-D MECOPROP-p AND DICAMBA GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	30.56%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid **‡	8.17%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-anisic Acid)***	2.77%
OTHER INGREDIENTS:	58.50%
TOTAL:	100.00%
Isomer Specific Method, Equivalent to:	
*2,4-Dichlorophenoxyacetic Acid	88 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	
***3,6-Dichloro-o-anisic Acid	22 lbs./gal.
±CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-D	•

FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY. NOT FOR SALE TO OR USE BY HOMEOWNERS.

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS


For Chemical Spill, Leak, Fire, or Exposure, Cail CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-312 EPA EST. NO. 228-IL-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



NET WEIGHT GALS.

000228-00312.20041111.name



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive, causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Avoid inhalation of spray mists. Harmful if swallowed or inhaled. Remove saturated clothing as soon as possible and shower. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

NON-WPS TURF USES: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) - in general only agricultural plant uses are covered by the WPS - must wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

NON-WPS INDUSTRIAL USES: When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry.

WPS USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) - in general, agricultural plant uses are covered - must wear: coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure and chemical-resistant apron when cleaning equipment, mixing or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing of PPE must not be reused until it has been cleaned. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Engineering Controls Statements: If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe or pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should

- · Wash hands, face, and arms with soap and water before eating, smoking, drinking or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID					
 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Çall a poison control center or doctor for treatment advice. 					
 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 					
Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.					
 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 					

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use the closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For turf use, the maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas. Also for use on Sod Farms.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpétgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Bedstraw	Dandelion	Lespedeza	Poison oak	Wild carrot
Black medic	Dock	Mallow	Purslane	Wild garlic
Buckhorn	Ground ivy	Morningglory	Ragweed	Wild lettuce
Burdock	Healall	Peppergrass	Sheep sorrel	Wild onion
Chicory	Henbit	Pigweed	Shepherdspurse	Yarrow
Chickweed	Knotweed	Plantain	Speedwell	and other
Clover	Lambsquarter	Poison ivy	Spurge	broadleaf weeds

ORNAMENTAL LAWNS & TURF

APPLICATION RATES

SITE	USE RATE (Fluid Ounces)	SPRAY VOLUME (Gallons)	USE RATE (Pints)	SPRAY VOLUME (Galions)	USE DIRECTIONS
	PER 1,00	0 SQ. FT.	PER	ACRE	
Bahiagrass,					Normal Applications
Bluegrass, Common Bermudagrass, Fescue, Ryegrass, Zoysiagrass	1.1 to 1.5	0.5 to 5	3 to 4	5 to 220	For Professional Lawn Maintenance - Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.
St. Augustinegrass and Centipedegrass	0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200	Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses. If Bermudagrass is dormant, up to 4 pints per
			ļ.,		acre may be used. However, some hybrid Bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.
Bentgrass (Putting and Bowling Greens)	0.67 Maximum	3.3	1.8 Maximum	145	Apply on closely mowed Bentgrass, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

LOW VOLUME SPRAY APPLICATION EQUIPMENT

EQUIPMENT	APPLICATION	USE RATE			
		Fluid Ounces per 1,0000 Square Feet	Pints per Acre	USE DIRECTIONS	
Controlled Droplet Applicators (CDA), Atomizers and Spinning Disk Applicators	Cool season grasses	1.1 to 1.5	3 to 4	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.	
	Listed warm season grasses	0.75 to 0.9	2 to 2.5	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.	
Lower Volume Equipment	Cool Season and Listed Warm Season Grasses	1.1 to 1.5	3 to 4	Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.	

NON-TURF AREAS

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1/3 to 1 gallon of this product per 50 to 300 gallons of water. This mixture will cover 43,500 square feet. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with the highwater volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 1 to 1-1/2 gallons of this product to 100 gallons of water applying 200 to 600 gallons of spray mixture per 43,500 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(RV 062003)

Triplet and The Optical Advantage are registered Trademarks of Nufarm Americas, Inc.





Nufarm Americas, Inc.
Jeannie K. Smith
Vice President
Registrations & Regulatory Affairs
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527
Phone: 708.755-7332 Fax: 708.754.0314
jeannie.smith@us.nufarm.com

November 15, 2004

Via Overnight Courier

Document Processing Desk (NOTIF)
Office of Pesticide Programs 7504C
US Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 South Bell Street
Arlington, VA 22202-4501

Subject:

Riverdale Triplet SF

EPA Reg. No. 228-312

Enclosed are the following items to support our request for an alternate brand name:

1. EPA Application Form 8570-1

2. Two (2) Copies of Amended Labeling with the new name, Primera Triplet SF.

Let me know if you have any questions or concerns. Thank you.

Regards,

Jannie K. Smith
Vice President

Designations (Designations

Registrations & Regulatory Affairs



Nufarm Americas, Inc.
Jeannie K. Smith
Vice President
Remistrations & Regulatory Affairs
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527
Phone: 708.755-7332 Hax: 708.754.0314
jeannie.smith@us.nufarm.com

Fax: 703 305-6929

NOTIFICATION

NOV 1 9 2004

November 23, 2004

Owen Beeder
Office of Pesticide Programs
US Environmental Protection Agency

Subject:

Riverdale intelessa

EPARCO, NO. 228-812

Notification Amendment Dated November 15, 2004

Addition of an allemate brand name (Primera hiple) SE

This politication is consistent with the provisions of PR Notice 9840 and EPA regulations at 40 GFR 152.46, and no other changes have been made to like labeling of the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. See, 1001 to will ally make any take statement of EPA. I taking understand that it this notification is not consistent with the terms of ER. Notice 98-10 and 40 GFR 152.46, this product may be in violation of EIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of EIFRA.

Letina know liyon hava any quesilons of concerns. Thank you

Regards

alim**ê** Xiênael

Vice President

Registrations & Regulatory Atlans