


228-312

10/1/2009

3 or 4 oct 10/9



**U S ENVIRONMENTAL PROTECTION AGENCY**  
**Office of Pesticide Programs**  
**Registration Division (7505P)**  
**Ariel Rios Building**  
**1200 Pennsylvania Ave , NW**  
**Washington, D C 20460**

EPA Reg Number  
228-312

Date of Issuance  
**OCT - 1 2009**

NOTICE OF PESTICIDE  
 \_ Registration  
X Reregistration  
 (under FIFRA as amended)

Term of Issuance

Name of Pesticide Product  
Triplet SF

Name and Address of Registrant (include ZIP Code)  
 Nufarm Americas Inc  
 150 Harvester Drive Suite 200  
 Burr Ridge Illinois 60527

**Note** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you

1 Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data

2 The Agency recommends that additional text be added to the Note to Physician that addresses eye irritation concerns

3 The following revisions are needed to the handler PPE section

-For Category A products, the first sentence of the PPE section should be revised to read

“Some materials that are chemical-resistant to this product are made of any waterproof material ”

-Per the 2,4-D, Dicamba and MCPP-p combined label table, the following text must be added to this section of the label

“Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate ”

-The text “Protective eyewear (goggles, face shield or safety glasses) should be revised to read “goggles or face shield ”

Signature of Approving Official

Joanne I Miller  
 Product Manager 23  
 Herbicide Branch  
 Registration Division (7505P)

Date

T -

**Droplet Size**

Use a medium or coarser spray nozzles according to (ASAE standard 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles

**Wind Speed**

Do not apply at wind speeds greater than 10 mph Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind

**Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height Do not make applications into areas of temperature inversions or stable atmospheric conditions

**Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants

**Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides Where states have more stringent regulations, they must be observed

**Equipment**

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates

Do not apply with a nozzle height greater than 4 feet above the crop canopy ”

11 The retreatment interval for all uses must be revised to 30 days per the MCPP-p RED

12 Add the following statements to the labeling

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al v EPA, C01- 0132C, (W D WA) For further information, please refer to [http //www epa gov/espp](http://www.epa.gov/espp)

A stamped copy of your labeling is enclosed for your records You must submit one (1) copy of the final printed label before you release the product for shipment Products shipped after 12 months from the date of this Notice must bear the new revised label If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e) Your release for shipment of the product constitutes acceptance of these conditions

Rec'd 9/11/08  
3089

# TRIPLET® SF

ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
OCT - 1 2009  
Under the Federal Insecticide  
Fungicide and Rodenticide Act  
as amended for the pesticide  
registered under EPA Reg. No

228-312

SELECTIVE BROADLEAF WEED CONTROL FOR TURFGRASSES INCLUDING  
USE ON SOD FARMS TO CONTROL CLOVER, DANDELION HENBIT, PLANTAINS,  
WILD ONION, AND MANY OTHER BROADLEAF WEEDS ALSO FOR HIGHWAYS  
RIGHTS-OF WAY AND OTHER SIMILAR NON CROP AREAS

CONTAINS 2 4 D MECOPROP p AND DICAMBA  
GET THE OPTICAL ADVANTAGE®

**ACTIVE INGREDIENTS**

Dimethylamine Salt of 2 4 Dichlorophenoxyacetic Acid	30.56%
Dimethylamine Salt of (+) R 2 (2 Methyl 4 Chlorophenoxy) propionic Acid ‡	8.17%
Dimethylamine Salt of Dicamba (3 6 Dichloro o anisic Acid)	2.77%

**OTHER INGREDIENTS**

<b>TOTAL</b>	<u>58.50%</u>
	100.00%

Isomer Specific Method Equivalent to

2 4 Dichlorophenoxyacetic Acid	25.38%	2.38 lbs /gal
(+) R 2 (2 Methyl-4 Chlorophenoxy)propionic Acid	6.75%	0.63 lbs /gal
3 6 Dichloro o anisic Acid	2.30%	0.22 lbs /gal

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP p

**FOR USE BY TURF MAINTENANCE PERSONNEL LANDSCAPING OR COMMERCIAL APPLICATORS ONLY**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER - PELIGRO**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle  
(If you do not understand the label find someone to explain it to you in detail )

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill Leak Fire or Exposure Call CHEMTREC (800) 424 9300  
For Medical Emergencies Only Call (877) 325 1840

EPA REG NO 228-312  
EPA EST NO 228 IL 1

MANUFACTURED BY  
NUFARM AMERICAS INC  
150 HARVESTER DRIVE  
BURR RIDGE IL 60527



NET WEIGHT                      GALS

000228 00312 20080401 EPA 24d Pending

4099

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER PELIGRO**

Corrosive causes irreversible eye damage Do not get in eyes or on skin or clothing Harmful if swallowed

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart

**All mixers, loaders applicators and other handlers must wear**

- Long sleeved shirt and long pants
- Shoes plus socks and
- Protective eyewear (goggles, face shield or safety glasses)
- Chemical resistant gloves

See engineering controls for additional requirements

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

**Engineering Controls Statements** If this container is over one gallon and less than five gallons then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron If this container is five gallons or more in capacity do not open pour product from this container A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container If the contents of a non refillable pesticide container are emptied the probe must be rinsed before removal If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240 (d) (4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS

When handlers use closed systems enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240 (d) (4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should</b>	
Wash hands face and arms with soap and water before eating smoking drinking or using the toilet	
Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing	
<ul style="list-style-type: none"> <li>• Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing</li> </ul>	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> </ul> Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person
<b>IF ON SKIN OR CLOTHING</b>	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 877 325 1840 for emergency medical treatment information	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage	

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of the closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas. Also for use on Sod Farms. For turf use, the maximum number of broadcast applications per treatment site is 2 per year. Product should not be used in or near greenhouses.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Aerial application is prohibited.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

**NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For turf use, do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

**SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

**Droplet Size**

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles.

**Wind Speed**

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at downwind edge of treated area.

**Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if 1) conditions of temperature inversion exist, or 2) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food forage or other plantings that might be damaged or crops thereof rendered unfit for sale use or consumption Susceptible crops include but are not limited to cotton okra flowers grapes (in growing stage) fruit trees (foliage) soybeans (vegetative stage) ornamentals sunflowers tomatoes beans and other vegetables or tobacco Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants

**Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2 4 D herbicides Where states have more stringent regulations they must be observed

**Equipment**

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates Do not apply with a nozzle height greater than 4 feet above the crop canopy

**USE PRECAUTIONS**

Avoid mist to vegetables flowers ornamentals shrubs trees and other desirable plants Do not pour spray solutions near these plants Do not spray on carpetgrass dichondra nor on lawns or turf where desirable clovers are present Avoid fine mists Except as noted use only lawn type sprayers Coarse sprays are less likely to 'wind drift' Use coarse spray droplets Do not spray roots of ornamentals and trees Do not exceed specified dosages for any area be particularly careful within the dripline of trees and other species Do not apply to newly seeded grasses until well established Avoid broadcast applications when air temperature exceeds 90 F When using small spot treatment applications in temperature over 90 F turf injury may occur

When treating Carpetgrass avoid broadcast applications when air temperature exceeds 80 F When air temperatures exceed 80 F limit application to spot treatment only

Apply only to dormant St Augustine (Common Raleigh and Seville varieties grown in Texas Louisiana Mississippi only) and dormant Centipede grasses (Texas Louisiana Mississippi only)

Avoid applying during excessively dry or hot periods unless irrigation is used For optimum results (1) turf should not be mowed for 1 to 2 days before and after application (2) do not apply if rain is expected within 4 hours after the application delay irrigation cycle for 24 hours Reseed no sooner than three to four weeks after application of this product Failure to observe all precautions may result in injury to turf and/or susceptible plants

This product can be mixed with some liquid fertilizers or liquid iron materials Because liquid fertilizer and liquid iron differ in pH free ammonia content density salt concentration and percentage of water a compatibility test is recommended prior to mixing in application equipment All regulations either State or Federal relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed

**WEEDS CONTROLLED**

- |             |              |              |                |                 |
|-------------|--------------|--------------|----------------|-----------------|
| Bedstraw    | Dandelion    | Lespedeza    | Poison oak     | Wild carrot     |
| Black medic | Dock         | Mallow       | Purslane       | Wild garlic     |
| Buckhorn    | Ground ivy   | Morningglory | Ragweed        | Wild lettuce    |
| Burdock     | Healall      | Peppergrass  | Sheep sorrel   | Wild onion      |
| Chicory     | Henbit       | Pigweed      | Shepherdspurse | Yarrow          |
| Chickweed   | Knotweed     | Plantain     | Speedwell      | and other       |
| Clover      | Lambsquarter | Poison ivy   | Spurge         | broadleaf weeds |

**TURF, ORNAMENTAL**  
 (golf courses cemeteries parks sports fields, turfgrass lawns and other grass areas)  
**AND**  
**TURF (grown for seed or sod)**

**APPLICATION RATES**

SITE	USE RATE (Fluid Ounces)	SPRAY VOLUME (Gallons)	USE RATE (Pints)	SPRAY VOLUME (Gallons)	USE DIRECTIONS
	PER:1 000 SQ FT		PER ACRE		
Bahiagrass Bluegrass Common Bermudagrass Fescue Ryegrass Zoysiagrass	1.1 to 1.5	0.5 to 5	3 to 4	5 to 220	<p><b>Normal Applications</b></p> <p><b>For Professional Lawn Maintenance</b> Higher water volumes may be used when tank mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.</p> <p>Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green up and from green up to dormancy. Some temporary discoloration may occur on warm season grasses.</p> <p><b>If Bermudagrass is dormant</b> up to 4 pints per acre may be used. However, some hybrid Bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.</p>
Dormant St Augustine (Common Raleigh and Seville varieties grown in Texas Louisiana Mississippi only) Dormant Centipede grasses (Texas Louisiana Mississippi only)	0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200	Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green up and from green up to dormancy. Some temporary discoloration may occur on warm season grasses.
Bentgrass (Putting and Bowling Greens)	0.67 Maximum	3.3	1.8 Maximum	145	<p>Apply on closely mowed Bentgrass preferably in May or mid August through September. Slight turf yellowing will disappear after about one week.</p> <p>On closely mowed golf course fairways and tees or bowling greens, apply when weeds are actively growing and temperatures are low. Exercise care to avoid over application to prevent injury. Temporary discoloration may occur.</p>

**NOTE** Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over application and subsequent turf discoloration or injury.

Ornamental Turf Make no more than 2 applications per year

Turf Grown for Seed/Sod Make no more than 2 applications per year. Minimum interval of 21 days between applications

**LOW VOLUME SPRAY APPLICATION EQUIPMENT**

EQUIPMENT	APPLICATION	USE RATE		USE DIRECTIONS
		Fluid Ounces per 1 000 Square Feet	Pints per Acre	
<b>Controlled Droplet Applicators (CDA) Atomizers and Spinning Disk Applicators</b>	Cool season grasses	1 1 to 1 5	3 to 4	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment) Do not overlap spray patterns
	Listed warm season grasses	0 75 to 0 9	2 to 2 5	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment) and follow CDA spray instructions for cool season grasses Use reduced rates if grass is stressed from heat drought etc
<b>Lower Volume Equipment</b>	Cool Season and Listed Warm Season Grasses	1 1 to 1 5	3 to 4	Use as little as 5 gallons of water per acre Use only application equipment that is capable of spreading a uniform droplet wetting each weed surface

**NON CROPLAND**

(fencerows hedgerows roadsides ditches rights of way utility powerlines railroads airports and industrial sites)

**Roadsides (including aprons and guardrails) rights-of way and other similar non-crop areas** For control of broadleaf weeds mix at a rate of 1 5 to 3 5 quarts of this product per acre in enough water to wet all parts of the foliage This may require 50 to 300 gallons of water per acre This mixture will cover 43 560 square feet Thoroughly saturate all weeds with spray mixture Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant Best results are obtained when weeds are young and actively growing Do not cut weeds until herbicide has translocated throughout the plant causing root death For small broadleaf weeds use the lower rate Heavy dense stands require the higher rate with the high water volume For small (spot) applications with small tank sprayers apply at the rate of 4 ounces of this product per gallon of water and spray to thoroughly wet all foliage

Annual/Perennial Control Make no more than 2 applications per year

For control of woody plants apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant All leaves stems and suckers must be completely wet to the ground line for effective control Regrowth may be anticipated on the more resistant species

**Woody Brush** Add 1 to 1 5 gallons of this product per acre to wet all parts of the brush foliage stem and bark This may require up to 200 to 600 gallons of water per 43 560 square feet depending upon the height and thickness of the brush Mix thoroughly before spraying

Woody Plant Control Make no more than 1 application per year

This product will either kill control or suppress the weeds listed in the label booklet for this product Some of these species may require repeat spot applications even under ideal conditions for application

**Maximum Seasonal Application Rate to non crop sites is 4 pounds 2 4 D Acid equivalent per acre per application site**

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**PESTICIDE STORAGE** Always use original container to store pesticides in a secured warehouse or storage building Store at temperatures above 32 F If allowed to freeze remix before using This does not alter this product Containers should be opened in well ventilated areas Keep container tightly sealed when not in use Do not stack cardboard cases more than two pallets high Do not store near open containers of fertilizer seed or other pesticides

**PESTICIDE DISPOSAL** Pesticide wastes are toxic If container is damaged or if pesticide has leaked contain all spillage Absorb and clean up all spilled material with granules or sand Place in a closed labeled container for proper disposal Improper disposal of excess pesticide spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

**CONTAINER DISPOSAL** Triple rinse (or equivalent) Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities Plastic containers are also disposable by incineration or if allowed by State and local authorities by burning If burned stay out of smoke



**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

**If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations on liability do not use the product and return it unopened to the Seller and the purchase price will be refunded**

(RV040108R)

Triplet and The Optical Advantage are registered Trademarks of Nufarm Americas Inc