228-313 3/15/2006

Page 136

Piease read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

\$epa	United St Environmental Prot Weshington, D	ection Age		X	Registration Amendme Other		OPP Identifier Number
	Appli	cation for I	Pesticide	- Section	I		
1. Company/Product Numbar			2. EPA Product Manager			3. Proposed Classification	
228-313			Joanne Miller			[]	None Restricted
4. Company/Product (Name) Horsepower Selective Herbicide			РМ# 23				
5. Name and Address of App Nufarm Americas, Inc. 1333 Burr Ridge Parkw Burr Ridge, IL 60527		6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No Product Name					
		Sec	tion - II				
Amendment - Explain	below.		F		els in repsonse to	,	
Resubmission in resp	onse to Agency letter dated			gency letter da Me Too" Applic		ľ	NOTIFICATION
X Notification - Explain	below.		<u> </u>	lther - Explain E	oolaw.	I	MAR 1 5 2006
Label notification consistent with PRN 95-2 and 98-10, see cover letter for detailed explanation. This notification is consistent with the provisions of PR Notice 95-2 and 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statements to EPA. I further understand that if this notification is not consistent withe terms of PR Notice 95-2 and 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.							
		Sect	ion - III				
1. Material This Product Will	Be Packaged In:		<u> </u>		_ 		
Child-Resistant Packaging	Unit Packaging	Water	Solubia Paci	laging	2. Type of Co	ntainer	
Yes	Yes		Yes			Aetal Iastic	
No	No		No			lass	
 Certification must be submitted 	If "Yes" No. p. Unit Packaging wgt, conta			No, per container		aper Ither (S	pecify)
3. Location of Net Contents	nformation 4. Size(s) Retail Contair	Hat.	5. L	cation of Label	Directio	ns
Label	ontainer						
6. Manner in Which Label is	Affixed to Product	Lithograph Paper glued Stanciled	[Other			
		Sect	on - IV				
1. Contact Point /Complete	items directly below for identit	fication of indivi	dual to be c	ontacted, if nai	cessary, to proce	ss this	application.)
Name Title Telephone No. (Include Area Code) George Meindl george.meindl@us.nufarm.com Registration Manager 708.755.7277							
Certification I certify that the statemente I have made on this form and all attachments thereto are true, accurate and complete. I ecknowledge that any knowlingity false or misleading statement may be purishable by fine or imprisonment or both under applicable law.							
2. Sightfore	nel	3. Title Regi	stratio	n Mana	iger		
4. Typed Nam George Meindl		5, Date 2/27	^{5, Date} 2/27/2006				

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

White - EPA File Copy (original)

Yesiuw - Applicant Copy



Nufarm Americas, Inc. George Meindl Registration Manager 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527 Phone: 708.755-7277 Fax: 708.754.0314 george.meindl@us.nufarm.com

February 27, 2006

Via Overnight Courier

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) U. S. Environmental Protection Agency Room 266A, Crystal Mall 2 1801 Bell Street Arlington, VA 22202-4501

Subject: Horsepower® Selective Herbicide EPA Reg. No. 228-313 Label notification per PR98-10 and PR95-2

Dear Ms. Hobgood:

Nufarm Americas, Inc would like to make label changes to the subject registration. We feel that these label changes are permitted and consistent with PRN95-2 section II (Notifications) part A (Labeling) subsection 4 (Revising Advisory Statements) and PRN98-10 section II (Labeling Notifications) part D (Adding Advisory Statements).

Specifically we have replaced the following sentence " Do not spray when air temperature exceeds 90°F" with the following 2 sentences "Avoid broadcast applications when air temperature exceeds 90 degrees. When using small spot treatment applications in temperatures over 90 degrees, turf injury may occur." We have also added in a separate paragraph with the following two advisory statements: "When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only."

The applicators of this product have told us the current wording was incomplete and confusing. We feel the suggested changes gives the applicator a clearer understanding on how to use this product in the most efficacious manner possible. These changes also support Nufarm's product stewardship initiatives and overall responsible care efforts.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling clean (1 copy)

If you should have any questions regarding this matter, please feel free to contact me at 708.755.7277 or email at george_meindl@us_nufarm.com.

Since ely,

George Meindl Registration-Manager Nufarm Americas, Inc. H:\228-313Vabel notification per PRN95-2 and 98-10 submitted 2.27.2006.doc



Clean

0

NOTIFICATION MAR 1 5 2006

RIVERDALE® HORSEPOWER® SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES. CONTROLS BLACK MEDIC, CHICKWEED, CLOVER COCKLEBUR, DANDELION, KNOTWEED, OXALIS, PLANTAIN, THISTLE AND MANY OTHER SPECIES OF BROADLEAF WEEDS; SOME OF WHICH ARE LISTED ON THIS LABEL.

CONTAINS MCPA, TRICLOPYR AND DICAMBA

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*		48.99%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**		5.59%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***		4.82%
OTHER INGREDIENTS:		40.60%
	TOTAL:	100.00%

By Isomer Specific AOAC Method, Equivalent to:	
*2-Methyl-4-Chlorophenoxyacetic Acid	40.0%, 3.80 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	4.0% 0.38 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	4.0%, 0.38 lbs./gal.

FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

NOTE: Spanish language is optional

EPA REG. NO. 228-313 EPA EST. NO. 228-IL-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

000228-00313.20060227.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

436

Corrosive. This concentrate causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear face shield or goggles when mixing and loading this product. After product has been diluted in accordance with Directions for Use, eye protection is not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Keep and wash protective clothing and equipment separately from other laundry.

FIRST AID		
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
IF SWALLOWED	 Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vorniting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
IF ON SKIN OR CLOTHING	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 	
	HOT LINE NUMBER Intainer or label with you when calling a poison control center or doctor, or going for treatment. t 1-877-325-1840 for emergency medical treatment information.	
	NOTE TO PHYSICIAN	

Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of imgation system. Do not contaminate water used for imgation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is for use on Ornamental Turl Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turl areas. Not for use on turf being grown for resale or other commercial use as sod, or for commercial seed production or for research purposes. *Excluding Bentgrass Tees.

Entry Restrictions: Do not allow people or pets to enter treated area until sprays have dried.

Application Restrictions: Avoid contact with eyes, skin, or clothing during and after application. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. This product may cause skin sensitization reactions in some people.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present.

Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on nonrecommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used;(2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sconer than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

Amaranth Aster Beadstraw **Bedstraw** Beggartick Beggarweed Bindweed Bitter sneezeweed Black medic Bladder campton Bracken fern Broomweed Buckhorn **Buckwheat** Buffalobur Burclover Burcucumber Burdock Bursage Buttercup Canada thistle Carpetweed Cathip Catsear Chamise Chamomile Chickweed Chicory Cinquefoil Clover Cocklebur Coffeeweed Cornflower Cornspeedwell Croton Daisv Dandelion Dock

Alfalfa

Dog fennel Dragonhead mint Evening primrose Field pennycress Fleabane Florida pusley Frenchweed Galinsoga Geranium Goatsbeard Goldenrod Goosefoot Gromweil Ground ivv Halfshrub sundrop Hawkweed Healall Heartleaf drymary Hemo sesbania Hempnettle Henbit Horsenettle Horseweed Ironweed Jerusalem artichoke Jimsonweed Knapweed Knapwood Knawel Knotweed Kochia Ladysthumb Lambsguarter Lespedeza Little starwort Mallow Marc's Tail Marshelder Matchweed

WEEDS CONTROLLED Mayweed Spurry Spurweed Milkweed Morningglory Starbur Stinging nettle Mustard Nightflowering catchfly Nightshade Oxalis(Stricta and corniculata) Parsley-piert Pennywort Peppergrass Pepperweed Pigweed Plantain Poison hemlock Poison ivv Poison oak Pokeweed Poorjoe Prickly sida Prickly lettuce Puncturevine Purslane Ragweed Red sorrel Red clover Redevine Redstem Rubberweed Sheep sorrel Shepherdspurse Sicklepod Smartweed Snakeweed Sowthistle Speedwell Spiderwort Spikeweed Spiny

Stitchwort Stunkweed Sumpweed Sunflower Tansy ragwort Teasel Teaweed Texas blueweed Thistle Toadflax Trumpetcreeper Velvetleaf Veronica Vetch Waterhemlock Waterhemp Whitebrush Wild violet Wild radish Wild onion Wild aster Wild carrot Wild lettuce Wild garlic Wild geranium Wood sorrel Wormwood Yankeeweed Yarrow Yellow starthistle Yellow rocket and many other broadleaf weeds.

7

ORNAMENTAL LAWNS AND TURF

Spurge

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

Herbi™ Controlled Droplet Applicator: Add 1-1/2 to 2-1/4 pints of this product to the 5 pint Herbi bottle, then fill with water to make

5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution.

While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of this product must be applied when grass is stressed from heat, drought, etc.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

NOTE: For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

- 1. Pour 18 ounces of water into a quart jar.
- 2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
- 3. Add 1 ounce of this product.
- 4. Close jar and shake well.
- 5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
- 6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
- 7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Co-op.'s Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
- 8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in a secured warehouse or storage building. Store at temperatures above 32 F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PRODUCT DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label. (RV022706N)

Riverdale and Horsepower are Registered Trademarks of Nufarm Americas Inc.

Herbi™ is a trademark of Micron Sprayers Limited Corporation, United Kingdom.