

PM 23

228-315

7/16/96

yellow P71/10



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

228-315

JUL 16 1996

Term of Issuance:

Conditional

Name of Pesticide Product:

Riverdale Sweet  
Sixteen Weed and Feed  
with Triplet MC Dri

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Company  
425 West 194th Street  
Glenwood, IL 60425-1584

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes before you release the product for shipment:
  - a. Revise the EPA Registration Number to read, "EPA Reg. No. "228-315".
  - b. Delete the first sentence under Environmental Hazards ("This product is toxic to aquatic invertebrates.") and the text "aquatic invertebrates and" from the second sentence. These statements are not required for acid forms of 2,4-D.
  - c. On the Commercial product label only, add the following statement to the Environmental Hazards section, preferably at the end of the first paragraph:

Signature of Approving Official:

Date:

JUL 16 1996

page 2  
EPA Reg. No. 228-315

"When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources."

- d. Delete the reference to 2,4-DP in the groundwater advisory statement (second paragraph under Environmental Hazards), since there is no 2,4-DP in this product.

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (7505C)

Enclosure

RD:STANTON:PM Team 23:Rm. 235:CM-2:305-5218:Disk #3:228-GRL.REG

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	S. Stanton							
DATE ▶	Jul 11, 1996							

(Non-Commercial Label)

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

3/10

JUL 16 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
228-315

RIVERDALE

SWEET SIXTEEN

WEED AND FEED WITH TRIPLET® MC DRI

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD  
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF  
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS: \*†

2,4-Dichlorophenoxyacetic Acid . . . . .	0.761%
2-(2-Methyl-4-Chlorophenoxy)propionic Acid . . . . .	0.324%
Dicamba (3,6-Dichloro-o-Anisic Acid) . . . . .	0.063%
INERT INGREDIENTS: . . . . .	98.852%
TOTAL . . . . .	100.000%

\*ISOMER SPECIFIC AOAC METHOD

GUARANTEED ANALYSIS: °

Total Nitrogen (N) . . . . .	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) . . . . .	X%
Soluble Potash (K <sub>2</sub> O) . . . . .	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum . . . . . X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer registrant)

†TRIPLET® - A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 16 LBS.


EPA REG. NO. 228-GRL

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

SWEET SIXTEEN WEED AND FEED WITH TRIPLET® MC DRI is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf, Drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or

5/10

early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .8 lb. per 250 square feet.

#### SPREADER SETTINGS

ONE PASS APPLICATION:°	MAKE	SETTING
	Central Calibrated	5
	Cyclone Broadcast	3-3/4
	Jackson	5
	Sears Broadcast	3
	Sears	5
	Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

#### STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizers, seeds or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

6/10

(ERP 061594) (PR938 042194)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Triplet® MC Dri - A Premium Broadleaf Herbicide

Environmental Benefits

(Grasscycling Logo)


Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

\*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration is optional except where mandated by State fertilizer regulation.



(Commercial Label)

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

7/10

JUL 16 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-315

RIVERDALE  
SWEET SIXTEEN  
WEED AND FEED WITH TRIPLET® MC DRI

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD  
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF  
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS: \*†

2,4-Dichlorophenoxyacetic Acid . . . . .	0.761%
2-(2-Methyl-4-Chlorophenoxy)propionic Acid . . . . .	0.324%
Dicamba (3,6-Dichloro-o-Anisic Acid) . . . . .	0.063%
INERT INGREDIENTS: . . . . .	98.852%
TOTAL . . . . .	100.000%

\*ISOMER SPECIFIC AOAC METHOD

GUARANTEED ANALYSIS: °

Total Nitrogen (N) . . . . .	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) . . . . .	X%
Soluble Potash (K <sub>2</sub> O) . . . . .	X%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum . . . . . X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer registrant)

†TRIPLET® - A Registered Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 16 LBS.


EPA REG. NO. 228-GRL

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCP, and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH TRIPLET® MC DRI is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Carpetweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Fenfuchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed,



Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .8 lb. per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 lb. nitrogen/1000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.  
\*Based on Inert fertilizer claimed.

**STORAGE AND DISPOSAL**

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide

To: FO

has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (ERP 061594) (PR938 042194)

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Triplet® MC Dri - A Premium Broadleaf Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy except as noted.

\*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. If Spreader Settings are deleted, the following sentence must be added: "Read your spreader product manual for proper settings and use instructions." Chlorine declaration is optional except where mandated by State fertilizer regulation.

