



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 14, 2016

Ms. Carrie M. Tackema
Nufarm Americas Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

Subject: Label Amendment – To add various weeds controlled and make other minor labeling changes.
Product Name: Turflon II Amine
EPA Registration Number: 228-316
Application Date: 09/09/2015
Decision Number: 509750

Dear Ms. Tackema:


The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Karen Samek by phone at (703) 347-8825, or via email at samek.karen@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Kathryn V. Montague". The signature is written in black ink and is positioned above the typed name.

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

ACCEPTED

06/14/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 228-316

TURFLON® II AMINE

[Alternate Brand Names: Aquasweep]

FOR THE CONTROL OF UNWANTED TREES AND BRUSH, AS WELL AS ANNUAL AND PERENNIAL BROADLEAF WEEDS ON RANGELAND, PASTURES, FENCEROWS, NON-IRRIGATION DITCHBANKS,ROADSIDES, SEASONAL IRRIGATION CANALS AND DITCHES, OTHER NON-CROP AREAS, NATIVE GRASS AREAS, RIGHTS-OF-WAY, ORNAMENTAL TURF, GOLF COURSES AND SOD FARMS.

FOR THE CONTROL OF EMERGENT, FLOATING AND SUBMERGED AQUATIC WEEDS IN THE FOLLOWING AQUATIC SITES: PONDS, LAKES, RESERVOIRS, MARSHES, BAYOUS, NON-IRRIGATION AND SEASONAL IRRIGATION CANALS AND DITCHES, SEASONAL IRRIGATION CANALS AND DITCHES WHICH HAVE LITTLE OR NO CONTINUOUS OUTFLOW AND IMPOUNDED RIVERS AND STREAMS THAT ARE QUIESCENT OR SLOW MOVING.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic Acid, Dimethylamine salt*	34.2%
Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid** as the triethylamine salt	15.2%

OTHER INGREDIENTS:	50.6%
TOTAL:	100.0%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	28.4%, 2.78 lbs./gal.
**Triclopyr acid	10.9%, 1.07 lbs./gal.

Notice: Read the entire label. Use only according to label directions.

Before buying or using this product, read "Warranty" elsewhere on this label.

**KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-316
EPA EST. NO. 228-IL-1

Manufactured For
NUFARM AMERICAS INC.
11901 S. Austin Avenue
Alsip, IL 60803



NET CONTENTS: _____ Gals. (_____ L)

000228-00316.20160607.draft

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER / PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Pilots must wear:

- long-sleeved shirt and long pants, and
- shoes and socks

All mixers, loaders, flaggers, and other applicators and handlers must wear:

- protective eyewear (goggles or face shield or shielded safety glasses),
- long sleeved shirt and long pants,
- shoes plus socks, and
- chemical-resistant gloves made of any waterproof material
- chemical resistant apron when mixing or loading, cleaning up spills, or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement:

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides. [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	
FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate use of gastric lavage.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Water having limited and dense weed infestations may not require partial treatments.

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application

around a cistern or well may result in contamination of drinking water or groundwater

Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable particularly where the water table is shallow, may result in groundwater contamination.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not cut or weld container.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides. (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **Restricted Entry Interval:** Do not allow people (other than applicator) or pets on treatment area during application. Do not enter into treated areas until sprays have dried or dust has settled.

PRODUCT INFORMATION

This product is for use on fence rows, non-irrigation ditchbanks, roadsides, other non-crop areas, native grass areas, industrial sites, rights-of-way, rangeland, seasonal irrigation canals and ditches, pastures, ornamental turf, golf courses, sod farms, and aquatic sites (including ponds, lakes, reservoirs, marshes, bayous, non-irrigation and seasonal irrigation canals and ditches, seasonal irrigation canals and ditches which have little or no continuous outflow and impounded rivers and streams that are quiescent or slow moving).

USE RESTRICTIONS

Do not apply this product directly to, or otherwise permit it to come into direct contact with, cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants and do not permit spray mists containing it to drift onto them. Very small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

Do not use for manufacturing or formulating.

Do not apply to exposed roots of shallow rooted trees or shrubs.

Do not apply on ditches used to transport irrigation water, except where canals and ditches are seasonal. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

USE PRECAUTIONS

Be sure that use of this product conforms to all applicable regulations.

Avoid Injurious Spray Drift. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured.

With ground broadcast equipment, drift can be reduced by keeping the spray boom as low as possible; by applying no less than 20 gallons of spray per acre; by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; and by spraying when wind velocity is low.

In handgun applications drift can be reduced by using low pressure Teejet flat fan, Raindrop or other drift controlling nozzles. Standard flat fan nozzles may also be used at no more than 20 psi. At this pressure the spray pattern may be narrower.

Foliar sprays should be applied during warm weather when brush and weeds are actively growing. Application under drought conditions may provide less than desirable results.

Note: Do not use spray equipment for other applications to land planted, or to be planted to susceptible crops or desirable plants, unless it has been determined that all phytotoxic herbicide residue has been removed by thorough cleaning of the equipment.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial application:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

APPLICATION INSTRUCTIONS – NONCROP

CONSERVATION RESERVE PROGRAMS AND GENERAL FARMSTEAD

This product is recommended for use for Conservation Reserve Programs, general farmstead (non-cropland only), weed and brush control, or use in State Recognized Noxious Weed areas (noncropland areas). Refer to 'Weeds Controlled' section for list of susceptible species. Some weed species require tank mixes for adequate control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For these uses, mix at a rate of 2 to 5 pints of this product per acre. Retreatment may be needed; however, the maximum seasonal application rate is 2 pounds 2,4-D acid equivalent per acre per application. Minimum of 30 days between applications, limited to 2 applications per year. The preharvest interval (PHI) is 14 days (for cut forage for hay). For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Weeds Controlled and Application Rates – Annual and Perennial Weeds

WEEDS CONTROLLED		RATE/ACRE	RATE/1,000 SQUARE FEET
black medic carpet weed catnip chamise chickweeds chicory cinquefoil clovers cocklebur coffee weed cornflower creeping beggarweed dandelion dayflower docks dwarf beggarweed field bindweed goldenrod	henbit hop clover knawel lambquarters lespedeza little starwort mallow matchweed plantain purslane speedwell smartweed sowthistle spiderwort spotted catsear Spurweed vetch yarrow	2-1/2 pints	1 fluid ounce or 2 tablespoons
burdock buttercup mustard	poison ivy poison oak purple loosestrife wild carrot	3 to 3-1/2 pints	1.3 fluid ounces or 2-1/2 tablespoons
cornspeedwell ground ivy oxalis (stricta and corniculata)	parsley-piert prostrate spurge wild violet	3 pints (2 applications, 4 to 6 weeks apart)	1.3 fluid ounces or 2-1/2 tablespoons (2 applications, 4 to 6 weeks apart)

Weeds Controlled – Tree and Brush Species

Alder	<i>Ceanothus</i> spp.	Maples (except bigleaf and vine*)	Sumac
Ash	Cherry (except black)	Multiflora Rose	Sycamore
Beech	Cottonwood	Poison Ivy	Tamarack
Birch	Dogwood	Poison Oak	Wax Myrtle (top growth)
Blackberry	Elderberry	Sassafras (top growth)	White Oak
Black Locust	Hawthorn	Scotch broom	Wild Grape
Boneset	Honeysuckle		Willow
Cascara			

*basal or dormant stem application only

Hard to Control Species

Buckbrush (<i>Symphoricarpos</i> spp.) (suppression)	Hazel	Salt Brush Salt Cedar (Tamarisk)
Brazilian Pepper Tree	Honeylocust (suppression)	Sweetgum
Common Persimmon (suppression)	Pine (suppression)	Trumpet creeper (suppression)
Elm (except winged elm)	Russian Olive	Virginia Creeper (suppression)
	Salmonberry (suppression)	

FOR PASTURE, RANGELAND, and NATIVE GRASS AREAS

Restrictions:

- 14 day pre-harvest interval (PHI) for grass hay
- 3 day pre-slaughter interval
- Do not allow lactating dairy animals to graze on grass forage until the next growing season after application.

- Do not make more than 2 applications per year with a minimum of 30 days between applications.

For susceptible annual, biannual and perennial weeds: Apply 2 ½ to 3 ½ pints per acre (see table above for rate recommendation).

For moderately susceptible biennial and perennial broadleaf weeds, difficult to control weeds and woody plants: Do not apply more than 5-1/2 pints of this product (2.0 lbs of 2,4-D ae) per acre per application.

Harder-To-Control Species: High-volume applications, 1.5% solution..

FOR FENCEROWS AND RIGHTS-OF-WAY

Maximum of 5-1/2 pints of this product (2.0 lbs of 2,4-D ae) per acre per application for treatment of annual and perennial weeds.

Maximum of 1 gallon, 3-1/2 pints of this product (4.0 lbs of 2,4-D ae) per year for treatment of woody plants.

Limited to 1 application per growing season.

14-day pre-harvest interval (PHI) for grass hay.

3-day pre-slaughter interval.

Do not allow lactating dairy animals to graze on grass forage until the next growing season.

FOR ROADSIDES

For annual and perennial plants:

Limit to 2 applications per year.

Maximum of 5-1/2 pints of this product (2.0 lbs of 2,4-D ae) per acre per application.

Minimum of 30 days between applications.

For woody plants:

Limit to 1 application per year.

Maximum of 1 gallon, 3-1/2 pints of this product (4.0 lbs of 2,4-D ae) per acre per year.

PREPARING THE SPRAY

Add about one-half the desired amount of clean water to spray tank. Add this product and complete addition of water with agitation running. Mix thoroughly and continue moderate agitation while spraying.

Size of Sprayer (Gallons)	AMOUNT OF THIS PRODUCT REQUIRED FOR SPRAY MIXTURE		
	1% Solution	1.5% Solution	4% Solution
1	1-1/3 fluid ounces	2 fluid ounces	5-1/3 fluid ounces
3	4 fluid ounces	6 fluid ounces	1 pint
5	6-2/3 fluid ounces	10 fluid ounces	1-2/3 pints
50	2 quarts	3 quarts	2 gallons
100	1 gallon	1.5 gallons	4 gallons

APPLICATION INSTRUCTIONS - TURF

BROADCAST TREATMENT OF ORNAMENTAL TURF, GOLF COURSES AND SOD FARMS

Apply 1 to 2 quarts of this product in enough water to make 20 to 200 gallons total spray per acre to control broadleaf weeds growing in perennial bluegrass, tall fescue, or perennial ryegrass. **Do not** use on other grass species, such as bentgrass or St. Augustinegrass, unless injury can be tolerated. Apply from early spring through fall when weeds are actively growing. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled. Additional applications should be made four weeks apart to minimize grass injury. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not reseed for 3 weeks after application.

SPOT TREATMENT OF ORNAMENTAL TURF, GOLF COURSES AND SOD FARMS USING PORTABLE SPRAYERS

Mix 1 to 2 fluid ounces of this product in enough water to make 3 gallons of spray (1 to 2 quarts per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of undesirable plants to point of runoff.

Ornamental Turf and Sod Farm Restrictions

Limited to 2 applications per year with a 21-day minimum interval.

Do not exceed 8.6 pints of this product (3 lbs of 2,4-D ae) per acre per year from all combined uses.
The maximum application rate for ornamental turf is 4-1/4 pints of this product (1.5 lbs of 2,4-D ae) per acre per application.
Do not exceed 11.5 pints of this product (4 lbs of 2,4-D ae) per acre per year from all combined uses.
The maximum application rate for sod farm turf is 5-1/2 pints of this product (2 lbs of 2,4-D ae) per acre per application.

APPLICATION INSTRUCTIONS - AQUATICS

FOR THE CONTROL OF EMERGENT, FLOATING AND SUBMERGED AQUATIC WEEDS IN THE FOLLOWING AQUATIC SITES: PONDS, LAKES, RESERVOIRS, MARSHES, BAYOUS, NON-IRRIGATION AND SEASONAL IRRIGATION CANALS AND DITCHES, SEASONAL IRRIGATION CANALS AND DITCHES WHICH HAVE LITTLE OR NO CONTINUOUS OUTFLOW AND IMPOUNDED RIVERS AND STREAMS THAT ARE QUIESCENT OR SLOW MOVING.

USE RESTRICTIONS FOR AQUATIC AND WETLAND SITES

Notice to Applicators: Before application, coordination and approval of local and state authorities may be required, either by letter or agreement or issuance of special permits for aquatic applications.

Use Requirements for Aquatic Areas:

When this product is applied to aquatic areas, follow PPE and reentry instructions in the "Non-Agricultural Use Requirements" section of this label.

Do not apply to salt water bays or estuaries.

Do not apply where runoff water may flow onto agricultural land as injury to crops may result.

Do not apply to unimpounded rivers and streams.

Do not apply to ditches or canals currently being used to transport irrigation water or that will be used for irrigation within 120 days following treatment or until triclopyr residue levels are determined to be 1.0 ppb triclopyr or less and 100 ppb 2,4-D or less.

RECREATIONAL USE OF WATER IN TREATMENT AREAS

There are no restrictions on the use of treated water for fishing. Do not swim in water treated with this product for (3) hours after treatment.

LIVESTOCK USE OF WATER FROM TREATMENT AREAS

There are no restrictions on consumption of treated water for potable use by livestock, pets or other animals.

CONTROL OF WEEDS AND BRUSH ON BANKS OF NON-IRRIGATION CANALS AND DITCHES

Application Rate

Apply 44 to 88 ounces of this product per acre to control annual weeds.

Apply 88 ounces of this product per acre for control of biennial and perennial weeds and susceptible woody plants.

Specific Use Directions

Apply using low pressure spray (10 to 40 psi) in a spray volume of 20 to 100 gallons per acre using power operated spray equipment.

Apply when wind speed is low, 5 mph or less.

Apply working upstream to avoid accidental concentration of spray into water. Cross-stream spraying to opposite banks is not permitted and do not allow boom spraying over water surface. When spraying shoreline weeds, allow no more than 2 foot overspray onto water surface with an average of less than 1 foot of overspray to prevent significant water contamination.

Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before stalks appear. For hard to control weeds, a repeat application after 30 days at the same rate may be needed.

For woody species and patches of perennial weeds, mix 1 gallon of this product per 64 to 150 gallons of total spray. Wet foliage by applying about 3 to 4 gallons of spray per 1000 sq ft (10.5 x 10.5 steps).

Restrictions and Limitations

Do not apply more than 2 treatments per season or reapply within 30 days.

Use 2 or more gallons of spray solution per acre.

Do not apply more than 92 oz/acre (2.0 lb of 2,4-D acid equivalent) per application or more than 184 oz/acre (4.0 lb of 2,4-D acid equivalent) per use season.

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance. Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS. Average Width (ft.) x Average Depth (ft.) x Average Velocity (ft. per sec.) = CFS

For ditchbank weeds, do not spray cross-stream to opposite bank. Do not allow boom spray to be directed onto water.

For shoreline weeds, boom spraying onto water surface must be held to a minimum and allow no more than 2 foot overspray onto water with an average of less than 1 foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Aerial Application

Use drift control spray equipment or thickening agent mixed in the spray mixture. Apply 175 oz of this product per acre using standard boom systems using a minimum spray volume of 5 gallons per acre. For Microfoil (r) - drift control spray systems, apply this product in a total spray volume of 12 to 20 gallons per acre. Refer to the "Spray Drift Management" section.

CONTROL OF EMERGENT AND FLOATING AQUATIC WEEDS: Including Water Hyacinth (*Eichhornia crassipes*) and Alligatorweed (*Alternanthera philoxeroides*)

Weeds Controlled

Emerged or Floating Weeds	Scientific Name
Alligatorweed	<i>Alternanthera philoxeroides</i>
Arrowhead	<i>Sagittaria spp.</i>
Bulrush	<i>Schoenoplectus spp.</i>
Duckweed	<i>Lemna spp.</i>
Frogsbit, American	<i>Limnobium spongia</i>
Lotus, American	<i>Nelumbo lutea</i>
Parrots feather	<i>Myriophyllum aquaticum</i>
Spatterdock	<i>Nuphar lutea</i>
Pickerelweed	<i>Pontederia cordata</i>
Pennywort	<i>Hydrocotyle spp.</i>
Purple Loosestrife	<i>Lythrum salicaria</i>
Soda Apple, Aquatic	<i>Solanum tampicense</i>
Water hyacinth	<i>Eichhornia crassipes</i>
Water lily	<i>Nymphaea spp.</i>
Water primrose	<i>Ludwigia spp.</i>
Watershield	<i>Brasenia schreberi</i>
Water chestnut	<i>Trapa natans</i>

SPECIFIC USE DIRECTIONS FOR EMERGENT AND FLOATING AQUATIC WEEDS

Application Rate

Apply 88 to 175 oz product per acre.

Application Timing

Spray weed mass only. Apply when plants are actively growing. A second application may be made 21 days after the initial application to kill regrowth and plants missed in previous operation. Use 175 oz/acre rate when plants are mature or when weed mass is dense.

Surface Application

Use power operated sprayers with boom or spray gun mounted on boat, tractor or truck. Thorough wetting of foliage is essential for maximum control. Use 100 to 400 gallons of spray mixture per acre. Special precautions such as use of low pressure, large nozzles and spray thickening agents should be taken to avoid spray drift to susceptible crops. Follow label directions for use of any drift control agent.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product may be tankmixed with other herbicides such as Clipper®, and Aqua Neat® for use against floating and aquatic weeds.

Restrictions and Limitations for Surface Applications to Emergent Aquatic Weeds

Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Waters having limited and less dense weed infestations may not require partial treatments. Other

local factors such as water exchange and sediment load can also influence the dissolved oxygen level. Coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for aquatic applications.

Do not exceed 184 oz/acre (4.0 lb of acid equivalent) per surface acre per use season. Limited to 2 applications per season.

Do not make a broadcast application within 21 days of previous broadcast application. Spot treatments are permitted.

**CONTROL OF SUBMERGED AQUATIC WEEDS:
Including Watermilfoils**

Weeds Controlled

Submersed Weeds	Scientific Name
Bladderwort	<i>Utricularia spp.</i>
Coontail	<i>Ceratophyllum demersum</i>
Parrots Feather	<i>Myriophyllum aquaticum</i>
Watermilfoil, Eurasian	<i>Myriophyllum spicatum</i>
Watermilfoil, Northern	<i>Myriophyllum sibiricum</i>
Watermilfoil, Hybrid	<i>Myriophyllum spicatum + sibiricum</i>
Watermilfoil, Variable leaf	<i>Myriophyllum heterophyllum</i>
Water Stargrass	<i>Heteranthera dubia</i>

SPECIFIC USE DIRECTIONS FOR EMERGENT AND FLOATING AQUATIC WEEDS

Application Rate

Apply up to 3.88 gallons (10.8 lb of acid equivalent) per acre foot of water.

Application Timing

Apply in spring or summer when aquatic weeds appear. Check for weed growth in areas heavily infested the previous year. A second application may be needed when weeds show signs of recovery.

Subsurface Application

Apply this product undiluted directly to the water through a boat mounted distribution system. Shoreline areas should be treated by subsurface injection application by boat to avoid aerial drift.

Apply to attain a concentration of 2 to 4 ppm 2,4-D and 0.75 to 1.5 ppm triclopyr (see table below).

Amount of 2,4-D and Triclopyr to Apply for a Target Subsurface Concentration					
2,4-D ae ppm	2	2.5	3	3.5*	4*
Triclopyr ae ppm	0.75	1.0	1.2	1.3*	1.5*
Avg Depth (ft)	This product gallons per surface acre at specified depth				
1	1.9	2.4	2.9	3.4	3.9
2	3.9	4.9	5.8	6.8	7.8
3	5.8	7.3	8.7	10.2	11.7
4	7.8	9.7	11.7	13.6	15.5
5	9.7	12.1	14.6	17.0	19.4

* For difficult conditions, for example, spot treatment of pioneer colonies of Eurasian Water Milfoil and certain difficult to control aquatic species.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product may be tankmixed with other herbicides such as Clipper®, for use against submersed weeds.

Restrictions and Limitations for Aquatic Sites with Submersed Weeds

Do not exceed 10.8 lbs. 2,4-D acid equivalent and 6.9 lbs triclopyr acid equivalent per acre foot.

Fish breathe oxygen in the water and a water-oxygen ratio must be maintained. Decaying weeds use up oxygen, but during the period when applications should be made, the weed mass is fairly sparse and the weed decomposition rate is slow enough that the water-oxygen ratio is not disturbed by treating the entire area at one time. If treatments must be applied later in the season when the weed mass is dense and repeat treatments are needed, apply product in lanes, leaving buffer strips which can then be treated when vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment.

Do not apply within 21 days of previous application. Limited to 2 applications per season.

When treating slow moving bodies of water, applications must be made while traveling upstream to prevent concentration of 2,4-D and triclopyr downstream from the application.

**WATER USE
Drinking water (potable water)**

POTABLE WATER INTAKE SETBACKS FOR CONTROL OF SUBMERGED WEEDS – LAKES, RESERVOIRS, OR PONDS

Minimum setback distances from functioning potable water intakes for human consumption for the application of this product must be observed when controlling submerged weeds in lakes, reservoirs or ponds.

Drinking water setback distances do not apply to terrestrial applications made adjacent to water bodies with potable water intakes.

Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits. The potable water use restrictions on this label are to ensure that consumption of water by the public is allowed only when the concentration of triclopyr in water is less than 400 ppb and the concentration of 2,4-D in water is less than the MCL (Maximum Contaminant Level) of 70 ppb. Applicators should consider the unique characteristics of the treated waters to assure that the triclopyr and 2,4-D concentrations in potable water do not exceed 400 ppb and 70 ppb, respectively, at the time of consumption.

For submersed weed applications, the drinking water setback distances from functioning potable water intakes are provided in the following table.

Drinking water setback distance for submersed weed applications				
application concentration and minimum setback distance (ft) from functioning potable water table intake				
2,4-D ae ppm	≤ 0.5	0.6 to 1	1.1 to 2	2.1 to 4
Triclopyr ae ppm	≤ 0.2	0.2 to 0.4	0.5 to 0.7	0.8 to 1.5
Minimum Setback Distance (ft)	600	1200	1800	2400

If no setback distance from the above Drinking Water Setback Distance Table is to be used for the application, applicators or the authorizing organization must provide a drinking water notification and an advisory to shut off all potable water intakes prior to a 2,4-D application. Notification to the party responsible for a public water supply or to individual private water users must be done in a manner to assure that the party is aware of the water use restrictions when this product is applied to potable water.

The following is an example of a notification via posting, but other methods of notification which convey the above restrictions may be used and may be required in some cases under State or local law or as a condition of a permit.

Example: Posting notification should be located every 250 feet including the shoreline of the treated area and up to 250 feet of shoreline past the application site to include immediate public access points. Posting should include the day and time of application.

Posting may be removed if analysis of a water sample collected at the intake shows that the concentration of triclopyr in water is 400 ppb or less and the 2,4-D is 70 ppb or less, or after 21 days following application, whichever occurs first. Use the *Sampling for Drinking Water Analysis After 2,4-D Application for Submersed Weed Applications Table* below to determine the minimum numbers of days to wait between application and water sampling for a given application target concentration.

Text of notification: Wait 21 days before diverting functioning surface water intakes from the treated aquatic site to use as drinking water, unless water at functioning drinking water intakes is tested and is demonstrated by assay to contain triclopyr concentrations of 400 ppb or less and concentrations of 2,4-D of 70 ppb or less. Use the *Sampling for Drinking Water Analysis After 2,4-D Application for Submersed Weed Applications Table* above to determine the minimum numbers of days to wait between application and water sampling for a given application target concentration.

Application Date: _____ Time: _____ .

Following each application of this product, treated water must not be used for drinking water unless one of the following restrictions has been observed:

- i. A setback distance described in the Drinking Water Setback Distance Table was used for the application, or
- ii. A waiting period of 21 days from the time of application has elapsed, or,
- iii. An approved assay indicates that the concentration of triclopyr is 400 ppb or less and the concentration of 2,4-D is 70 ppb or less at the water intake. Sampling for drinking water analysis should occur no sooner than stated in Sampling for Drinking Water Analysis After 2,4-D Application for Submersed Weed Applications Table above. Analysis of samples must be completed by a laboratory that is certified under the Safe Drinking Water Act to perform drinking water analysis using a currently approved version of analytical Method Number 515, 555, other methods for 2,4-D as may be listed in Title 40 CFR, Part 141.24, or Method Number 4015 (immunoassay of 2,4-D) from U.S. EPA Test Methods for Evaluating Solid Waste SW-846.

Except as stated above, there are no restrictions on using water from treated areas for fishing or watering livestock.

Coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for aquatic applications.

Minimum days after application before initial water sampling at the functioning potable water intake			
2,4-D ae ppm	≤0.5	0.6 to 2	2.1 to 4
Triclopyr ae ppm	≤0.2	0.3 to 0.7	0.8 to 1.5
Minimum Days	5	10	14

Note: These are general guidelines; the amount of time required for residues to reach concentrations acceptable for drinking or irrigation will depend on the total acres treated relative to water body size, application rates, water exchange rates, weed density, and various other factors.

WATER USE Irrigation Purposes

Irrigation:

Do not use water treated with the product for irrigating greenhouse or nursery plants unless the triclopyr and 2,4-D residues are confirmed to be less than 1 ppb by laboratory analysis.

Do not use water treated with this product for irrigating hydroponic crops.

Do not apply under circumstances where spray drift may occur to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Do not use treated water for irrigation for 120 days following application or until residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb triclopyr or less and 100 ppb 2,4-D or less.

There is no restriction on use of water from the treatment area to irrigate established grasses. Do not apply this product through any type of irrigation system.

Seasonal Irrigation Waters:

This product may be applied during the off-season to surface waters that are used for irrigation on a seasonal basis, provided that there is a minimum of 120 days between applying this product and the first use of treated water for irrigation purposes, or until residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb triclopyr or less and 100 ppb 2,4-D or less.

Irrigation Canals/Ditches:

Do not apply this product to irrigation canals/ditches unless the 120 day restriction on irrigation water usage can be observed or residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb triclopyr or less and 100 ppb 2,4-D or less.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition et al vs. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web site: <http://www.epa.gov/espp>.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: (This product is available in multiple containers. Follow the container handling instructions below that apply to your container type/size)

[Nonrefillable Containers 5 Gallons or Less:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds.

Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Nonrefillable Containers Larger than 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable Container Larger than 5 Gallons:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Or

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV060716)

Turflon is a trademark of Dow AgroSciences, LLC.

Aquasweep is a trademark of Nufarm Americas