

228-323

12-27-99

DEC 27 1999

Russell F. Sawyer
 Riverdale Chemical Company
 425 West 194th Street
 Glenwood, IL 60425-1584

Dear Mr. Sawyer:

Subject: Use Dilution Amendment - Reduced PPE
 Riverdale TruPower Selective Herbicide
 EPA Registration No. 228-323
 Your Submission Dated August 27, 1998

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the following change:

1. Correct the spelling of the word, "to", in the phrase, "in sufficient water to assure coverage (1 too 4 gallons of water per acre)", under the heading, "Controlled Droplet Applicators" within the "Ornamental Lawns and Turf" section.

A stamped copy of the label is enclosed for your records. Submit one copy of your final printed labeling before you release the product for shipment. We apologize for the delay in responding to this submission.

Sincerely yours,

Joanne I. Miller
 Product Manager 23
 Herbicide Branch
 Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	JIMiller							
DATE ▶	Dec 27, 1999							

RIVERDALE

TRUPOWER™ SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES

COPY

CONTAINS MCPA, CLOPYRALID AND DICAMBA

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis,
Clover, Cocklebur, Thistle and many other species of
broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	48.13%
Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid**	5.18%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.73%
INERT INGREDIENTS:	41.96%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	39.30%, 3.75 lbs./gal.
**3,6-Dichloro-2-Pyridinecarboxylic Acid	3.93%, 0.37 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	3.93%, 0.37 lbs./gal.

TruPower™ - Is A Trademark of Riverdale Chemical Company

For Use By Professional Turf Maintenance Personnel,
Landscaping or Commercial Applicators Only.
Not For Sale to or Use By Homeowners.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-323

NET CONTENTS GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

NOTE: Spanish language is optional

ACCEPTED
with COMMENTS
In EPA Letter Dated

DEC 27 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-323

Revised 8/27/98 Revised PC & SPT for eyes & skin.

3076

(Inside Booklet)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER - PELIGRO**

Corrosive. This concentrate causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses) when mixing and loading this product. After product has been diluted in accordance with directions for use below, eye protection is not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: If contacted with this concentrate, hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention. If contact is made after product has been diluted, flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: If contacted with this concentrate, wash with plenty of soap and water. Get medical attention. If contact is made after product has been diluted, wash with plenty of soap and water. Rinse thoroughly.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, eggwhites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this product will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TruPower™ SELECTIVE HERBICIDE is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees* and Roughs), and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.*Excluding Bentgrass Tees.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are

present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

The suitable use of TruPower on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED BY TRUPOWER™ SELECTIVE HERBICIDE

alfalfa, Amaranth, Aster, Bachelor button, Bedstraw, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campion, Bracken fern, Broomweed, Buckhorn, Buckwheat, Buffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground ivy, Groundsel, Halfshrub sundrop, Hawksbeard, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle, Henbit, Horsenettle, Horseweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Locoweed, Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red sorrel, Red clover, Redvine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse, sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny, Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sumpweed, Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Toadflax, Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild radish, Wild onion, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow earthistle, Yellow rocket, and many other broadleaf weeds.

ORNAMENTAL LAWNS AND TURF

Apply TruPower Selective Herbicide at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

Herbi Controlled Droplet Applicator: Add 1-1/2 to 2-1/4 pints of TruPower Selective Herbicide to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution. While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Reduce application rates to stressed turf unless temporary injury can be tolerated.

Herbi™ is a trademark of North American Micron.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use TruPower Selective Herbicide at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) Avoid double coverage resulting from overlap unless temporary

Riverdale TruPower Selective Herbicide

turf injury can be tolerated. (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

COMPATIBILITY

TruPower Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of TruPower Selective Herbicide.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Marcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding the TruPower Selective Herbicide.)
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds or other pesticides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings and replace all caps. Contact Riverdale Chemical's Customer Service Department at 1-708/754-3330, to arrange for return of the empty refillable container.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label. (PR938&11 042194/RV 082798)

COPY