

228-325

11/29/2006

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

NOV 29 2006

George Meindl  
Nufarm Americas Inc.  
150 Harvester Drive, Suite 200  
Burr Ridge, IL 60527

Dear Mr. Meindl:

Subject: Revised Labeling – CAUTION Signal Word  
Sweet Sixteen Weed and Feed with Horsepower  
EPA Registration Number 228-325  
Riverdale Horsepower Weed and Feed  
EPA Registration Number 228-326  
Your Submissions Dated July 26, 2006

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. In the Hazards to Humans section add:

Causes moderate eye irritation.

b. In the PPE section add:

Safety glasses

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.


A stamped copy of the labeling is enclosed for your records.

2/11

-2-

If you have any questions concerning this letter, please contact Mr. James Stone at 703-305-7391. For your assistance enclosed are copies of the scientific reviews.

Sincerely yours,

A handwritten signature in black ink that reads "Joanne I. Miller". The signature is written in a cursive style with a large, looping initial "J".

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure

3/11

(Residential Label)

# RIVERDALE® SWEET SIXTEEN WEED AND FEED WITH HORSEPOWER®

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS MCPA, TRICLOPYR, AND DICAMBA.  
Weeds contacted by this product gradually die and disappear from the lawn.  
Treats (or covers) 5,000 Square Feet (50' x 100')

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid (MCPA)*	1.10%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid (Triclopyr)**	0.12%
Dimethylamine Salt of 3,6-Dichloro-o-Anisic Acid (Dicamba)***	0.11%

<b>OTHER INGREDIENTS:</b>	<b>98.67%</b>
<b>TOTAL</b>	<b>100.00%</b>

**ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:**

*2-Methyl-4-Chlorophenoxyacetic Acid	0.897%
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.089%
***3,6-Dichloro-o-Anisic Acid	0.089%

**GUARANTEED ANALYSIS:\***

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	X%
Soluble Potash (K <sub>2</sub> O)	X%
Primary Plant Foods derived from (List sources)	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	
Fertilizer guaranteed by (Name of fertilizer registrant)	

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
**SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-325  
EPA EST. NO. 228-IL-1

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

MANUFACTURED BY  
NUFARM AMERICAS INC.  
BURR RIDGE, IL 60527



NET CONTENTS      LBS.

NOV 29 2006  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

000228-00325.20060718.FIN

228-325

4/11

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Applicators and all other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Keep and wash protective clothing and equipment separately from other laundry.</li> </ul>	

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is a stable mixture of three herbicides. It is for use on residential turf.

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area until after application. Avoid contact with eyes, skin, or clothing during and after application.

The maximum application rate is 9 lbs. active ingredient per acre per year.

5/11

This product may be used on Bahia, Bent, Bermuda, Blue Grass, Fescue, Rye, and Zoysia grass. Avoid applications to Centipede and St. Augustine grass unless injury can be tolerated. Do not use on Dichondra, Carpetgrass, or where desirable clovers are present.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Beggarweed	Dog fennel	Mustard	Spurge
Bindweed	English daisy	Oxalis (Stricta and corniculata)	Spurweed
Black medic	Florida pusley	Parsley-piert	Stinging nettle
Buckhorn	Frenchweed	Pennywort	Stitchwort
Burdock	Goldenrod	Pepperweed	Thistle
Buttercup	Ground ivy	Pigweed	Toadflax
Canada thistle	Hawkweed	Plantain	Veronica
Carpetweed	Healall	Poison ivy	Vetch
Catnip	Heartleaf drymary	Poison oak	Wild aster
Chamise	Henbit	Purslane	Wild carrot
Chickweed	Jimsonweed	Ragweed	Wild garlic
Chicory	Knawel	Red clover	Wild geranium
Cinquefoil	Knotweed	Red sorrel	Wild lettuce
Clover	Kochia	Sheep sorrel	Wild onion
Cocklebur	Lambsquarter	Shepherdspurse	Wild radish
Coffeeweed	Lespedeza	Smartweed	Wild violet
Cornflower	Little starwort	Sowthistle	Wood sorrel
Cornspedwell	Mallow	Speedwell	Yarrow
Dandelion	Matchweed	Spiderwort	and many other broadleaf
Dock	Morningglory	Spotted catsear	weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, trees, vegetables, flowers, or desirable plants; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two-pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 lbs. for 16 pound bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .7 pound (.8 lb. for 16 pound bag) per 250 square feet.

**SPREADER SETTINGS**

ONE PASS APPLICATION:°	MAKE	SETTING
	Central Calibrated	5
	Cyclone Broadcast	3-3/4
	Jackson	5
	Sears Broadcast	3
	Sears	5
	Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product

6/11

will supplement your regular fertilizer program by providing \*(No. of lbs.)\* pounds of nitrogen per 1,000 square feet (optional P2O5 and K2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

**Environmental Benefits**

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

**STORAGE AND DISPOSAL**

**STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

**DISPOSAL:** If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV071806)

Riverdale and Horsepower are registered trademarks of Nufarm Americas Inc.

**OPTIONAL SPREADER INSTRUCTIONS:**

*(Illustration)* For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

*(Illustration)* For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

**OPTIONAL MARKETING STATEMENTS:**

Contains Horsepower® - A Premium Broadleaf Herbicide

The following note will not be part of the actual label copy.

\*Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

7/11

(Commercial Label)

# RIVERDALE® SWEET SIXTEEN WEED AND FEED WITH HORSEPOWER®

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS MCPA, TRICLOPYR, AND DICAMBA.

Weeds contacted by this product gradually die and disappear from the lawn.  
Treats (or covers) 5,000 Square Feet (50' x 100')

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid (MCPA)*	1.10%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid (Triclopyr)**	0.12%
Dimethylamine Salt of 3,6-Dichloro-o-Anisic Acid (Dicamba)***	0.11%

<b>OTHER INGREDIENTS:</b>	<u>98.67%</u>
<b>TOTAL</b>	<b>100.00%</b>

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.897%
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.089%
***3,6-Dichloro-o-Anisic Acid	0.089%

**GUARANTEED ANALYSIS:<sup>a</sup>**

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	X%
Soluble Potash (K <sub>2</sub> O)	X%
Primary Plant Foods derived from (List sources)	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	
Fertilizer guaranteed by (Name of fertilizer registrant)	

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-325  
EPA EST. NO. 228-IL-1

NET CONTENTS      LBS.

000228-00325.20060718.FIN

MANUFACTURED BY  
**ACCEPTED** NUFARM AMERICAS INC.  
with COMMENTS BARR RIDGE, IL 60527  
In EPA Letter Dated:



2007  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No

228-325

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Applicators and all other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Keep and wash protective clothing and equipment separately from other laundry.

**FIRST AID**

<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is a stable mixture of three herbicides. It is for use on residential turf.



9/11

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Avoid contact with eyes, skin, or clothing during and after application.

The maximum application rate is 9 lbs. active ingredient per acre per year.

This product may be used on Bahia, Bent, Bermuda, Blue Grass, Fescue, Rye, and Zoysia grass. Avoid applications to Centipede and St. Augustine grass unless injury can be tolerated. Do not use on Dichondra, Carpetgrass, or where desirable clovers are present.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Beggarweed	Dog fennel	Mustard	Spurge
Bindweed	English daisy	Oxalis (Stricta and corniculata)	Spurweed
Black medic	Florida pusley	Parsley-piert	Stinging nettle
Buckhorn	Frenchweed	Pennywort	Stitchwort
Burdock	Goldenrod	Pepperweed	Thistle
Buttercup	Ground ivy	Pigweed	Toadflax
Canada thistle	Hawkweed	Plantain	Veronica
Carpetweed	Healall	Poison ivy	Vetch
Catnip	Heartleaf drymary	Poison oak	Wild aster
Chamise	Henbit	Purslane	Wild carrot
Chickweed	Jimsonweed	Ragweed	Wild garlic
Chicory	Knawel	Red clover	Wild geranium
Cinquefoil	Knotweed	Red sorrel	Wild lettuce
Clover	Kochia	Sheep sorrel	Wild onion
Cocklebur	Lambsquarter	Shepherdspurse	Wild radish
Coffeeweed	Lespedeza	Smartweed	Wild violet
Cornflower	Little starwort	Sowthistle	Wood sorrel
Cornspeedwell	Mallow	Speedwell	Yarrow
Dandelion	Matchweed	Spiderwort	and many other broadleaf
Dock	Morningglory	Spotted catsear	weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, trees, vegetables, flowers, or desirable plants; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 14 (or 16) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two-pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 2.8 pounds (3.2 pounds for 16 lb. bag) per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .7 lbs. (.8 lbs. for 16 lb. bag) per 250 square feet.

**SPREADER SETTINGS**

ONE PASS APPLICATION:°	MAKE	SETTING
	Central Calibrated	5
	Cyclone Broadcast	3-3/4
	Jackson	5
	Sears Broadcast	3
	Sears	5
	Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader area with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *\*(No. of lbs.)\** pounds of nitrogen per 1,000 square feet (optional P2O5 and K2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

## Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV071806)

Riverdale and Horsepower are registered trademarks of Nufarm Americas Inc.

### OPTIONAL SPREADER INSTRUCTIONS:

*(Illustration)* For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

*(Illustration)* For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

### OPTIONAL MARKETING STATEMENTS:

Contains Horsepower® - A Premium Broadleaf Herbicide



The following note will not be part of the actual label copy.

\*Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.