

(Commercial Label)

RIVERDALE

HORSEPOWER™ GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS MCPA, TRICLOPYR, AND DICAMBA.

Weed contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	1.10%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	0.12%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.11%
INERT INGREDIENTS:	98.67%
TOTAL	100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.897%
**3,5-6-Trichloro-2-Pyridinyloxyacetic Acid	0.089%
***3,6-Dichloro-o-Anisic Acid	0.089%

†Horsepower™ Is A Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT LBS.


EPA REG. NO. 228-328

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH HORSEPOWER™ is a stable mixture of three herbicides. It is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, and Golf Courses (Fairways, Aprons, Tees, and Roughs).

This product may be used on Bahia, Bent (excluding golf course greens and tees), Bermuda, Blue Grass, Fescue, Rye, and Zoysia grass. Avoid applications to Centipede and St. Augustine grass unless injury can be tolerated. Do not use on Dichondra, Carpetgrass, or where desirable clovers are present.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggarweed, Bindweed, Black medic, Buckhorn, Burdock, Buttercup, Canada thistle, Carpetweed, Catnip, Chamise, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Dandelion, Dock, Dog fennel, English daisy, Florida pusley, Frenchweed, Goldenrod, Ground ivy, Hawkweed, Healall, Heartleaf drymary, Henbit, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarter, Lespedeza, Little starwort, Mallow, Matchweed, Morningglory, Mustard, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, pepperweed, Pigweed, Plantain, Poison ivy, Poison oak, Purslane, Ragweed, Red clover, Red sorrel, Sheep sorrel, Shepherdspurse, Smartweed, Sowthistle, Speedwell, Spiderwort, Spotted catsear, Spurge, Spurweed, Stinging nettle, Stitchwort, Thistle, Troadflax, Veronica, Vetch, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild violet, Wood sorrel, Yarrow and many other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, trees, vegetables, flowers, or desirable plants; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after

application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 lb. per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

STORAGE AND DISPOSAL


STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(PR938 042194/RV 022798)



4 2.8

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Horsepower™ - A Premium Broadleaf Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

*Note: Spreader Settings will vary among various commercial manufacturers.

598

(Non-Commercial Label)

RIVERDALE

HORSEPOWER™ GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS MCPA, TRICLOPYR, AND DICAMBA.

Weed contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	1.10%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	0.12%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.11%
INERT INGREDIENTS:	98.67%
TOTAL	100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.897%
**3,5-6-Trichloro-2-Pyridinyloxyacetic Acid	0.089%
***3,6-Dichloro-o-Anisic Acid	0.089%

†Horsepower™ Is A Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT LBS.


EPA REG. NO. 228-328

EPA EST. NO. 228-IL-1


MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



Revised 2/27/98 Changed Statement of Practical Treatment to agree with 228-325, -326 & -327.



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH HORSEPOWER™ is a stable mixture of three herbicides. It is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, and Golf Courses (Fairways, Aprons, Tees, and Roughs).

This product may be used on Bahia, Bent (excluding golf course greens and tees), Bermuda, Blue Grass, Fescue, Rye, and Zoysia grass. Avoid applications to Centipede and St. Augustine grass unless injury can be tolerated. Do not use on Dichondra, Carpetgrass, or where desirable clovers are present.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Beggarweed, Bindweed, Black medic, Buckhorn, Burdock, Buttercup, Canada thistle, Carpetweed, Catnip, Chamise, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Dandelion, Dock, Dog fennel, English daisy, Florida pusley, Frenchweed, Goldenrod, Ground ivy, Hawkweed, Healall, Heartleaf drymary, Henbit, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarter, Lespedeza, Little starwort, Mallow, Matchweed, Morningglory, Mustard, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, pepperweed, Pigweed, Plantain, Poison ivy, Poison oak, Purslane, Ragweed, Red clover, Red sorrel, Sheep sorrel, Shepherdspurse, Smartweed, Sowthistle, Speedwell, Spiderwort, Spotted catsear, Spurge, Spurweed, Stinging nettle, Stitchwort, Thistle, Toadflax, Veronica, Vetch, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild violet, Wood sorrel, Yarrow and many other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, trees, vegetables, flowers, or desirable plants; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 or 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to listed Warm Season Grasses - under conditions of inadequate

moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 lb. per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE

SETTING

Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

STORAGE AND DISPOSAL


STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

PR938 042194/RV 022798)



898

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Horsepower™ - A Premium Broadleaf Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 22% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 30' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

*Note: Spreader Settings will vary among various commercial manufacturers.

