riease read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

<b>≎EPA</b>	United States Environmental Protection Agency Washington, DC 20480  Application for Pesticide - Section I				OPP Identifier Number						
		Application	n tot P				<u> </u>	<del></del>	-		
1. Company/Product Number 228-332				2. EPA P	roduct Man Miller	eger		3.	3. Proposed Classification		
4. Company/Product (Name)				PM#				[	None	Restu	ricted
Millennium Ultra 2 Herbicio	de		23								
5. Name and Address of Applicant (Include ZIP Code)  Nufarm Americas, Inc. 1333 Burr Ridge Parkway Suite 125A  Burr Ridge, IL 60527  Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No.  Product Name					ling			
<u> </u>	, , , , , , , , , , , , , , , , , , , ,		Sact	ion - II	t Name _	<u></u>		=			
1		<del></del>	Seci	1011 - 11			<del></del> _				
Amendment - Explain t	)alow,				Final printed Agency lett		ls in repson: ed	e to	NO	TIFICAT	ION
Resubmission in respon	nse to Agency letter	dated	<del></del>	- []	"Me Too" A					R 15 2	
X Notification - Explain be	elow.				Other - Expl	ain be	olow.		61 <b>7</b> p	1 20 2	ovu.
Explanation: Use additional	negate if necessary	- /For section	n I and Sec	tion II I	<del></del>				·		
PR Notice 95-2 and 98-10 an statement of formula of this p understand that if this notifica	Label notification consistent with PRN 95-2 and 98-10, see cover letter for detailed explanation. This notification is consistent with the provisions of PR Notice 95-2 and 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statements to EPA. I further understand that if this notification is not consistent withe terms of PR Notice 95-2 and 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.						er				
			Secti	on - II							
1. Material This Product Will B			T				T				
Child-Resistant Packaging   Ves	Unit Packaging		(	Water Soluble Packaging 2. Type of Contains Yes Metal							
No	No		11	No.			Plasti	astic			
Constitution was a	lf "Yes" Unit Packaging wgt.	No. per container	If "Yes" No. per Paper								
		4. Size(s) Ret	- Contain		L	E 100	cation of Le	hal Direc	-siana		
3. Location of Net Contents Information 4. Size(s) Retail  Label Container			ail Container   5. Location of Label Directions								
8. Manner in Which Label is Affixed to Product   Lithograph   Other   Paper plued   Stencied											
Section - IV											
1. Contact Point (Complete its	1. Contact Point										
Name George Meindl george.meindl@us.nufarm.com								755.727		ode)	
Certification  I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.  I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.  6. Data Application Received  (Stamped)											
2. Signature ( ) 1 ) ( ) 3.			3. Title				7				
Top I had			Registration Manager				_				
4. Typod Narh George Meindl			2/27/2006						į		



Nufarm Americas, Inc. George Meindl Registration Manager 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527

Phone: 708.755-7277 Fax: 708.754.0314 george.meindl@us.nufarm.com

February 27, 2006

Via Overnight Courier

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 Bell Street
Arlington, VA 22202-4501

Subject:

Millennium Ultra™ 2 Herbicide

EPA Reg. No. 228-332

Label notification per PR98-10 and PR95-2

Dear Ms. Hobgood:

Nufarm Americas, Inc would like to make label changes to the subject registration. We feel that these label changes are permitted and consistent with PRN95-2 section II (Notifications) part A (Labeling) subsection 4 (Revising Advisory Statements) and PRN98-10 section II (Labeling Notifications) part D (Adding Advisory Statements).

Specifically we have replaced the following sentence "Do not spray when air temperature exceeds 90°F" with the following 2 sentences "Avoid broadcast applications when air temperature exceeds 90 degrees. When using small spot treatment applications in temperatures over 90 degrees, turf injury may occur." We have also added in a separate paragraph with the following two advisory statements: "When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only."

The applicators of this product have told us the current wording was incomplete and confusing. We feel the suggested changes gives the applicator a clearer understanding on how to use this product in the most efficacious manner possible. These changes also support Nufarm's product stewardship initiatives and overall responsible care efforts.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling clean (1 copy)

If you should have any questions regarding this matter, please feel free to contact me at 708.755.7277 or email at <a href="mailto:george.meindl@us.nufarm.com">george.meindl@us.nufarm.com</a>.

Sincerply,

George Meihdl Registration Manager Nufarm Americas, Inc.

H:\228-332\label notification per PRN95-2 and 98-10 submitted 2.27.2006.doc

Clean

3 7 9

MAR 1 5 2006

# MILLENNIUM ULTRA<sub>M</sub><sup>2</sup>

### FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS, TURF GRASSES, AND ON SOD FARMS

#### **CONTAINS 2,4-D, CLOPYRALID AND DICAMBA**

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*.  Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid** Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***  OTHER INGREDIENTS:  TOTAL	2.54% 4.65% 55.49%
Isomer Specific Method, Equivalent to:  *2,4-Dichlorophenoxyacetic Acid  *3,6-Dichloro-2-Pyridinecarboxylic Acid  *3,6-Dichloro-o-Anisic Acid  3.86%, 0.375	lbs./gal.

FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY. NOT FOR SALE TO OR USE BY HOMEOWNERS.

<sup>1</sup>NOT FOR SALE OR USE IN NASSAU AND SUFFOLK COUNTIES IN N.Y. STATE
IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY.
IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE
AGRICULTURAL, RIGHTS-OF-WAY, GOLF COURSE OR CEMETERY SITES.

## KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-332 EPA EST. NO. 228-1L-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes, and rubber gloves when mixing, loading, or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

NON-WPS TURF USES: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170)--in general only agricultural plant uses are covered by the WPS--must wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes and chemical-resistant gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

NON-WPS INDUSTRIAL USES: When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

WPS USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170)—in general, agricultural plant uses are covered—must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear, when cleaning equipment, mixing or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other taundry. After each day of use, clothing of PPE must not be reused until it has been cleaned. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Engineering Controls Statements: If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **USER SAFETY RECOMMENDATIONS**

#### Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID					
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>				
IF SWALLOWED	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>				
IF ON SKIN OR CLOTHING	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>				
	HOT LINE NUMBER				

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertigal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system, Do not contaminate water used for irrigation or domestic purposes,

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions partaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

#### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

#### **USE PRECAUTIONS**

This product is for use on Non-Residential Ornamental Turf Lawns (Industrial and Institutional) Athletic/Recreational Fields (Parks, Playgrounds and Sports Fields) Cemeteries, Sod Farms, Golf Courses (Fairways, Aprons, Tees\* and Roughs), Industrial Sites, Non-Cropland, Rights-of-Way and Roadsides. \*Excluding Bentgrass Tees.

Do not use on residential turfgrass. Do not collect grass clippings for mulch or compost. Do not send clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for mulching or composting.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipedegrass, St. Augustinegrass, Dichondragrass, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not use grass clippings from turf treated with this product for composting and/or mulch. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

#### COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

- 1. Pour 18 fluid ounces of water into a quart jar.
- 2. Add 1 fluid ounce of either the liquid fertilizer or liquid iron to be used.
- 3. Add 1 fluid ounce of this product.
- 4. Close jar and shake well.
- 5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
- 6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application,
- 7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 fluid ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
- 8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

#### **WEEDS CONTROLLED**

Alder	Bloodflower, milkweed	Catsear	Clover, strawberry
Artichoke	Boxelder	Catsear, spotted	Clover, sweet
Aster	Brassbuttons	Chamomile, corn	Clover, white
Aster, wild	Bristly oxtongue	Chickweed	Cockle
Bedstraw	Broomweed	Chickweed, common	Cocklebur
Bedstraw, pineywoods	Buckhorn	Chickweed, mousear	Cocklebur, spiny
Beggartick	Buckwheat, wild	Chicory	Coffeebean
Betony, Florida	Bur ragweed	Cinquefoil	Coffeeweed
Biden	Burdock	Cinquefoil, rough	Coreopsis, plains
Bindweed	Burweed	Clematis, Western	Creeping Jenny
Bindweed, field	Buttercup	Clover	Croton
Bindweed, hedge	Buttercup, bulbous	Clover, annual yellow	Croton, wooly
Bitter wintercress	Carpetweed	sweet	Cudweed
Bittercress, hairy	Carrot, wild	Clover, crimson	Curly indigo
Bitterweed	Catchweed bedstraw	Clover, hop	Daisy, English
Black-eyed Susan	Catnip	Clover, red	Daisy, oxeye

Dandelion Knotweed Peppergrass St. Johnswort Dandelion, fall Knotweed, prostrate Pepperweed Stinkweed Dandelion, talse Kochia Pepperweed, Virginia Stitchwort Dock Lambsquarter Strawberry, wild Pigweed Dogbane Lespedeza Pigweed, prostrate Sumac Dogfennel Lettuce, blue Pigweed, regroot Sunflower Elderberry Lettuce, prickly Pigweed, Russian Sunflower, false Fiddleneck Lettuce, wild Pigweed, smooth Sweet Potato, wild Fieldcress, Austrian Locoweed Pigweed, tumble Tansy ragwort Filaree, redstem Lupine Scarlet pimpernel Tanweed Mallow Flax, false Plantain Tarweed Fleabane (daisy) Mallow, Indiana Plantain, black-seed Thistle Fleabane, hairy Plantain, bracted Mallow. Venice Thistle, blessed Fleabane, rough Marigold, roundleafed Plantain, broadleaf Thistle, bull Marigold, wild Thistle, Canada Flixweed Plantain, buckhorn Florida pusley Marshelder Plantain, hoary Thistle, musk Thistle, Russian Matchweed Four-o'-clock, wild Plantain, narrowleaf Frenchweed Medic, black Plantain, slender Thistle, Scotch Galinsoga Mexicanweed Plantain, woolly Toadflax Galinsoga, smallflower Morningglory Pokeweed Trailing crownvetch Garlic, wild Morningglory, woolly Trefoil, tick Poorjoe Geranium, Carolina Mugwort Povertyweed Tumbleweed Mullein, common Geranium, wild Prickly sida Velvet leaf Goathead Mustard Primrose Veronica Goatsbeard Mustard, black Puncture vine Vervain Goldenrod Mustard, garlic Purslane Vervain, blue Mustard, hedge Ground-ivy Radish, wild Vervain, hoary Groundsel Mustard, tansy Vervain, prostrate Ragweed Mustard, tumble Vervain, tall Gumweed Rape, wild Hawkweed Mustard, white Rose, Southern wild Vetch Hawkweed, orange Mustard, wild Vetch, bird Rush Healall Nettle Salsify, Western Vetch, milk Heartleaf drymary Vetch, narrowleaf Nettle, buil Shepherdspurse Heathaster Nettle, burning Smartweed Vetch, wild Hemlock, poison Nettle, dead Virginia buttonweed Sneezeweed Nettle, stinging Hemp Sorrel, red Virginia creeper Henbit Nettle, tall Wavyleaf bullthistle Sorrel, sheep Hoary cress Oak, poison Sowthistle Willow Honeysuckle Onion, wild Sowthistle, common Witchweed Horsenettle Oxalis Spanishneedle Woodsorrel Horsetail Pansy, field Spatterdock Wormseed ironweed Parsley-piert Speedwell Yarrow Ivy, Poison Parsnip Spiny amaranth Yellow rocket Jewelweed Parsnip, wild Yellowflower pepperweed Spurge

Pearlwort

Pennycress

Pennywort

Jimsonweed

Knawel

Knapweed, spotted

Some of these species may require repeat spot applications and/or use of higher rate recommended on this product label even under ideal conditions for application.

Spurge prostrate

Spurge, spotted

Spurweed

and other broadleaf weeds

### ORNAMENTAL LAWNS, TURF, AND SOD FARMS

	MILLENNIUM ULTRA <sup>2</sup> USE RATE		SPRAY VOLUME IN GALLONS		APPLICATION METHODS		
SITE	FI. Oz. Per 1,000 Sq. Ft.	Pints per Acre	Per 1,000 Sq. Ft.	Per Acre	AND INSTRUCTIONS		
Cool Season Grasses	.73 to 1.10	2 to 3	0.02	1 to 4	Controlled Droplet and Spinning Disc Applicators		
(Bluegrass Fescue spp. Ryegrass)			up to 0.11	up to 5	Low Volume Ride On, Walk Behind and Towable Applicators (such as Perma Green™ and Turf Tamer®)		
}				<u> </u>	Note: Use only equipment that is capable of delivering uniform coverage.		
ł	{	!	0.11	5 to 10	Low Volume Boom Sprayers		
f 1					Limited coverage of the target leaf surface may reduce performance.		
ł	1		0.23 to 1.84	10 to 80	Conventional Boom Sprayers		
ļ			0.46 to 4.6	20 to 200	Professional Lawn Care Hand Held Gun		
					Note: For optimum results use spray volumes between 85 and 135 gallons per acre.		
			per Gallon per 1,000 Sq. Ft.	] <del></del>	Portable Sprayers (such as back packs, hand- held cans and atv units)		
Bentgrass	0.55 to 0.73	1.5 to 2	.023 to 5.5	20 to 240	Conventional Boom Sprayers		
(excluding Golf Course Greens and Tees)					Avoid applications during periods when turf is under stress due to high heat, humidity and reduced moisture. Slight yellowing may occur but will disappear in about seven days.		
Warm Season Grasses: (Bahiagrass, Common Bermudagrass, and Zoysiagrass)	0.73 to 1.10	2 to 3	0.02 to 0.09	1 to 4	Controlled Droplet and Spinning Disc Applicators		
			up to 0.11	up to 5	Low Volume Ride On, Walk Behind and Towable Applicators (such as Perma Green™ and Turf Tamer*) Note: Use only equipment that is capable of delivering uniform coverage.		
		l	0.23 to 0.55	5 to 10	Low Volume Boom Sprayers Limited coverage of the target leaf surface may reduce performance.		
	1		0.23 to 1.84	10 to 80	Conventional Boom Sprayers		
	{		0.46 to 4.6	20 to 200	Professional Lawn Care Hand Held Gun		
					Precautions: Use lower rates when turf is under drought stress. Applications made during transition from dormancy to green up may delay green up or cause temporary discoloration.		
	.73 to 1.10		per Gallon per 1,000 Sq. Ft.		Portable Sprayers (such as back packs, hand- held cans and atv units)		

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label.

(RV022706N)

<sup>1</sup>The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of New York, California, Oregon and/or Washington. The label prohibitions are optional for all other states.

Millennium Ultra is a Trademark of Nularm Americas Inc.