



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Registration Division (R9588E)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
228-335

Date of Issuance:  
MAR 4 1998

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Riverdale Horsepower™  
8000 Lawn Weed Killer

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Company  
425 West 194th Street  
Glenwood, Illinois 60425-1584

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

- a. Revise the EPA Registration Number to read, "EPA Reg. No.228-335".
- b. On the Non-Commercial label, add the following statement in the paragraph under "Hazards To Humans And Domestic Animals":

"If using the disposable sprayer and bottle, wear eye protection when refilling the bottle or when connecting or disconnecting the bottle from the sprayer."

Signature of Approving Official:

Date:  
MAR 4 1998

c. On the Non-Commercial label, please delete the directions for "Spot Treatment". These directions instruct users to put the subject product into "an empty quart container with trigger sprayer." These directions are too vague and could present a health and safety hazard if the product is put into inappropriate containers or stored in improperly labeling bottles. EPA would prefer that you provide the trigger sprayer for use in spot treatments. If you wish to do so, you may submit an amendment to reinstate the spot treatment directions for EPA's review.

3. Submit one (1) copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Copies of EPA's reviews of the product-specific data you submitted in support of this application are also enclosed.

Sincerely,

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Attachment

3?>3EBB5>35C

SYMBOL ▶	7505C						
SURNAME ▶	MHOWARD						
DATE ▶	Mar 3, 1998						

(Non-Commercial Label)

RIVERDALE

HORSEPOWER™ 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL  
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, TRICLOPYR AND DICAMBA

KILLS (OR CONTROLS): Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis,  
Clover, Cocklebur, Thistle and many other species of  
broadleaf weeds; some of which are listed on this label.  
One quart covers 8,000 sq. ft.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	10.97%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	1.25%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	1.08%
INERT INGREDIENTS:	86.70%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	8.960%	0.770 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.896%	0.077 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	0.896%	0.077 lbs./gal.

HORSEPOWER™ - Is A Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que  
se la explique a usted en detalle.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND  
STATEMENT OF PRACTICAL TREATMENT

PA REG. NO. 228-

NET CONTENTS GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY


RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

NOTE: Spanish language is optional

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

MAR 4 1998  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended for the pesticide  
registered under EPA Reg. No.  
228-335



## (Side Panels)

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles, face shield or safety glasses when mixing and loading this product. After product has been diluted in accordance with Directions for Use, eye protection is not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**STATEMENT OF PRACTICAL TREATMENT**

*Add: If using the disposable sprayer & bottle, wear eye protection when refilling the bottle or when connecting or disconnecting the bottle from*

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HORSEPOWER™ 8000 LAWN WEED KILLER is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs), and similar non-crop areas.

**USE PRECAUTIONS**

Spray when air is calm to avoid spray drift. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum kill (or control) of weeds will be obtained from Spring or early fall applications when weeds are actively growing. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede, or St. Augustine grasses unless turf injury can be tolerated. The suitable use of Horsepower Lawn Weed Killer on non-recommended turf species

may be determined by treating a small area at any rate/sq. ft. which does not exceed 3 pints/9,600 sq. ft. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

#### WEEDS CONTROLLED BY HORSEPOWER LAWN WEED KILLER

Alfalfa, Amaranth, Aster, Beadstraw, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter  
 Creeperweed, Black medic, Bladder campion, Bracken fern, Broomweed, Buckhorn, Buckwheat,  
 Buffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle,  
 Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover,  
 Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog  
 fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley,  
 Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground ivy,  
 Halfshrub sundrop, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle,  
 Henbit, Horsenettle, Horseweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed,  
 Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort,  
 Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard,  
 Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert,  
 Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison  
 oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed,  
 Red sorrel, Red clover, Redvine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse,  
 Sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny,  
 Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sumpweed,  
 Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Thistle, Toadflax,  
 Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild  
 violet, Wild radish, Wild onion, Wild aster, Wild carrot, Wild lettuce, Wild garlic, Wild  
 geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket,  
 and many other broadleaf weeds.

#### ORNAMENTAL LAWNS AND TURF

d.k. Apply Horsepower 8000 Lawn Weed Killer at the rate of 2 pints in 4 to 50 gallons of water  
 per 8,000 sq. ft. (4 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to  
 control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf  
 course greens and tees), Bahia, Bermuda and Zoysia.

**Disposable Sprayer and Bottle Instructions:** Determine area to be sprayed. Apply at a rate  
 of 32 oz. (1 quart) per 8,000 sq. ft. Connect spray nozzle to hose. Remove bottle cap and  
 screw bottle into nozzle. While holding sprayer at waist level, point in direction away  
 from face and body. Turn on the water at the tap. Place finger over hole on right side  
 of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix  
 automatically at the proper ratio. Walk back and forth at steady pace.

2,000 Sq. Ft.

4,000 Sq. Ft. (Optional marks on left for 1 qt. disposable bottle.  
 Not to scale.)

6,000 Sq. Ft.

8,000 Sq. Ft.

Review

**For Spot Treatment:** Put 2.1 oz. of Horsepower 8000 Lawn Weed Killer into an empty quart container with trigger sprayer. Then add 30 oz. of tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds.

**STORAGE AND DISPOSAL**

**STORAGE:** To be stored in original container and placed in an area inaccessible to children.

**PESTICIDE (OR PRODUCT) DISPOSAL:** Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Rinse thoroughly before discarding in trash. Do not reuse empty

**NOTICE:** Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (PR938 042194)

(Commercial Label)

RIVERDALE

HORSEPOWER™ 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL  
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, TRICLOPYR AND DICAMBA

KILLS (OR CONTROLS): Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis,  
Clover, Cocklebur, Thistle and many other species of  
broadleaf weeds; some of which are listed on this label.  
One quart covers 8,000 sq. ft.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	10.97%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	1.25%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	1.08%
INERT INGREDIENTS:	86.70%
TOTAL	100.00%

Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	8.860%	0.770 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.896%	0.077 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	0.896%	0.077 lbs./gal.

HORSEPOWER™ - Is A Trademark of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que  
se la explique a usted en detalle.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND  
STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228- NET CONTENTS GALS. EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584


NOTE: Spanish language is optional

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

MAR 4 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-335



(Side Panels)

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles, face shield or safety glasses when mixing and loading this product. After product has been diluted in accordance with Directions for Use, eye protection is not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

**HORSEPOWER™ 8000 LAWN WEED KILLER** is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs), and similar non-crop areas.

**USE PRECAUTIONS**

/ Spray when air is calm to avoid spray drift. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum kill (or control) of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede, or St. Augustine grasses unless turf injury can be tolerated. The suitable use of Horsepower Lawn Weed Killer on non-recommended turf species may be determined by treating a small area at any rate/sq. ft. which does not exceed 3



9 of 10

pints/9,600 sq.ft. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

#### WEEDS CONTROLLED BY HORSEPOWER LAWN WEED KILLER

Alfalfa, Amaranth, Aster, Beadstraw, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campton, Bracken fern, Broomweed, Buckhorn, Buckwheat, Ruffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gronwell, Ground ivy, Halfshrub sundrop, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle, Henbit, Horsenettle, Horsetweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red sorrel, Red clover, Redevine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny, Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sumpweed, Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Thistle, Toadflax, Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild violet, Wild radish, Wild onion, Wild aster, Wild carrot, Wild lettuce, Wild garlic, Wild geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket, and many other broadleaf weeds.

#### ORNAMENTAL LAWNS AND TURF

Apply Horsepower Lawn Weed Killer at the rate of 2 pints in 4 to 50 gallons of water per 8,000 sq. ft. (4 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

*Retal* **For Spot Treatment:** Put 2.1 oz of Horsepower 8000 Lawn Weed Killer into an empty quart container with trigger sprayer. Then add 30 oz. of tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds.

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds or other pesticides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**NOTICE:** Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (PR938 042194)