

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

228-335

01/30/2006

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 30 2006

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

George Meindl
Nufarm Americas, Inc.
1333 Burr Ridge Parkway
Suite 125A
Burr Ridge, IL 60527-0866

Dear Mr. Meindl:

Subject: Correction Needed to Label
Riverdale HORSEPOWER 8000 LAWN WEED KILLER (Residential Label)
EPA Registration No. 228 - 335
Application Date: October 10, 2005
Accepted/Stamped Date: November 18, 2005

The *correction* is required in reference to the above-mentioned Label Amendment, which was previously accepted on November 18, 2005. Please note the following *change*:

1. ENVIRONMENTAL HAZARDS -

i). Entry Restrictions:

Change "..... until dusts have settled." to " until sprays have dried."

Please submit one (1) copy of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records. **This amended labeling supersedes all previously accepted ones.**

Sincerely yours,

A handwritten signature in black ink that reads "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

2/6

Riverdale®

(Residential label)

HORSEPOWER® 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN RESIDENTIAL TURF
CONTAINS MCPA, TRICLOPYR AND DICAMBA

KILLS (OR CONTROLS): Clover, Cocklebur, Dandelion, Chickweed, Black medic, Ground Ivy, Knotweed, Plantain, Oxalis, Spurge, Thistle, Wild Violet and many other species of broadleaf weeds; some of which are listed on this label.

One Quart Covers 8,000 Square Feet

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	10.97%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	1.25%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	1.08%
OTHER INGREDIENTS	<u>86.70%</u>
TOTAL:	100.00%

By Isomer Specific AOAC Method, Equivalent To:

*2-Methyl-4-Chlorophenoxyacetic Acid	8.960%, 0.770 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.896%, 0.077 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	0.896%, 0.077 lbs./gal.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-335
EPA EST. NO. 228-IL-1

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN 30 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GA.,S.

000228-00335 00350215 RED

228-335

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Entry Restrictions: Do not allow people or pets to enter the treated area until dusts have settled.

Application Restrictions: Avoid contact with eyes, skin, or clothing during and after application. Do not apply this product in a way that will contact person or pet either directly or through drift. Keep people and pets out of the area during application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is for use on residential turf.

USE PRECAUTIONS

Spray when air is calm to avoid spray drift. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable Clovers are present. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum kill (or control) of weeds will be obtained from spring or early fall applications when weeds are actively growing. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede, or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/square feet which does not exceed 3 pints/15,000 square feet. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall

4/6

is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

Alder	Common chickweed	Horsetail	Prostrate spurge	Thistle
Annual yellow sweet clover	Common mullein	Indiana mallow	Prostrate vervain	Tick trefoil
Artichoke	Common sowthistle	Ironweed	Puncture vine	Toadflax
Aster	Corn Chamomile	Jewelweed	Purslane	Trailing Crownvetch
Austrian fieldcress	Creeping jenny	Jimsonweed	Ragweed	Tumble mustard
Bedstraw	Crimson clover	Kochia	Red clover	Tumble pigweed
Beggartick	Croton	Knawel	Redroot pigweed	Tumbleweed
Biden	Cudweed	Knotweed	Red sorrel	Velvet leaf
Bindweed	Curly dock	Lambsquarter	Redstem filaree	Venice mallow
Bird vetch	Curly indigo	Lespedeza	Rough cinquefoil	Veronica
Bitter wintercress	Dandelion	Locoweed	Rough fleabane	Vervain
Bitterweed	Dead nettle	Lupine	Roundleafed marigold	Vetch
Black-eyed Susan	Dock	Mallow	Rush	Virginia buttonweed
Black medic	Dogbane	Marshelder	Russian pigweed	Virginia creeper
Black mustard	Dogfennel	Matchweed	Russian thistle	Virginia pepperweed
Black-seed plantain	Elderberry	Mexicanweed	St. Johnswort	Wavyleaf bullthistle
Blessed thistle	English daisy	Milk vetch	Scarlet pimpernel	Western clematis
Blue lettuce	Fall dandelion	Milkweed bloodflower	Scotch thistle	Western salsily
Blue vervain	False dandelion	Morningglory	Sheep sorrel	White clover
Box elder	False flax	Mouseear chickweed	Shepherdspurge	White mustard
Bracted plantain	False sunflower	Mugwort	Slender plantain	Wild mustard
Brassbuttons	Fiddleneck	Musk thistle	Smallflower galinsoga	Wild aster
Bristly oxtongue	Field bindweed	Mustard	Smartweed	Wild buckwheat
Broadleaf dock	Field pansy	Narrowleaf plantain	Smooth dock	Wild carrot
Broadleaf plantain	Flea bane (daisy)	Narrowleaf vetch	Smooth pigweed	Wild four-o'clock
Broomweed	Flixweed	Nettle	Sneezeweed	Wild garlic
Buckhorn	Florida betony	Orange hawkweed	Southern wild rose	Wild geranium
Buckhorn plantain	Florida pusley	Oxalis	Sowthistle	Wild lettuce
Bulbous buttercup	Frenchweed	Oxeye daisy	Spanishneedle	Wild marigold
Bull nettle	Galinsoga	Parsley-piert	Spatterdock	Wild onion
Bull thistle	Garlic mustard	Parsnip	Speedwell	Wild parsnip
Burdock	Goathead	Pearlwort	Spiny Amaranth	Wild radish
Burning nettle	Goatsbeard	Pennycress	Spiny cocklebur	Wild rape
Bur ragweed	Goldenrod	Pennywort	Spotted catsear	Wild strawberry
Burweed	Ground ivy	Peppergrass	Spotted knapweed	Wild sweet potato
Buttercup	Gumweed	Pepperweed	Spotted spurge	Wild vetch
Canada thistle	Hairy bittercress	Pigweed	Spurge	Willow
Carolina geranium	Hairy fleabane	Pineywoods bedstraw	Spurweed	Witchweed
Carpetweed	Hawkweed	Plains coreopsis	Stinging nettle	Woolly morningglory
Catchweed bedstraw	Healall	Plantain	Stinkweed	Woodsorrel
Catsear	Heartleaf drymary	Poison hemlock	Stitchwort	Woolly croton
Catnip	Hedge bindweed	Poison ivy	Strawberry clover	Woolly plantain
Chickweed	Hedge mustard	Poison oak	Sumac	Warmseed
Chicory	Hemp	Pokeweed	Sunflower	Yarrow
Cinquefoil	Henbit	Poorjoe	Sweet clover	Yellow rocket
Clover	Hoary cress	Povertyweed	Tall nettle	Yellowflower
Cockle	Hoary plantain	Prickly lettuce	Tall vervain	pepperweed
Cocklebur	Hoary vervain	Prickly sida	Tansy mustard	and other broadleaf
Coffeebean	Honeysuckle	Primrose	Tansy ragwort	weeds.
Coffeeweed	Hop clover	Prostrate knotweed	Tanweed	
	Horsenettle	Prostrate pigweed	Tarweed	

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 pints in 4 to 50 gallons of water per 8,000 square feet (4 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

5/6

Disposable Sprayer and Bottle Instructions: Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 8,000 sq.ft. Remove bottle cap and screw bottle into nozzle. Connect spray nozzle to hose while holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.


- 2,000 square feet
- 4,000 square feet
- 6,000 square feet
- 8,000 square feet

(Optional marks on left for 1 quart disposable bottle. Not to scale.)

OR

HOSE-END SPRAYER INSTRUCTIONS:

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 8,000 square feet. Before attaching sprayer to garden hose, make sure that the blue flow switch is in the "OFF" position; red flow switch in the up, "CLOSED" position. Connect spray nozzle to garden hose while holding sprayer at waist level, point in direction away from face and body. Insure sprayer is securely tightened to bottle. Turn on water at the tap; move blue flow switch to "ON" position; and, turn red flow switch to the forward, "OPEN" position. This product (or product name) will mix automatically at the proper ratio. When spraying, walk back and forth at a steady pace. When finished spraying, turn blue flow switch to "OFF" position and red flow switch to "CLOSED" position.



STORAGE AND DISPOSAL

PESTICIDE STORAGE: To be stored in original container and placed in an area inaccessible to children.

PESTICIDE (OR PRODUCT) DISPOSAL: If empty - Do not reuse this container except as specified in directions to prepare refill. Place in trash or offer for recycling if available. If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith.

(RV091505)

Riverdale and Horsepower are registered trademarks of Nufarm Americas Inc.

Optional Marketing Statements
 For Both Cool Season & Warm Season Grasses
 (Product) is for use on Ornamental Turf

Alternate Statements
 OTHER

*Please note, some states still may require the term INERT.

Editor's Note: Subregistrant will choose either Disposable or Hose-end instructions.

NEXT

LABEL

228-335

01/30/2006

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 30 2006

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

George Meindl
Nufarm Americas, Inc.
1333 Burr Ridge Parkway
Suite 125A
Burr Ridge, IL 60527-0866

Dear Mr. Meindl:

Subject: LABEL AMENDMENT - TRICLOPYR
Riverdale HORSEPOWER 8000 LAWN WEED KILLER (Commercial Label)
EPA Registration No. 228 - 335
Application Date: January 20, 2006

The amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling change below before you release the product for shipment bearing the amended labeling:
 - a. ENVIRONMENTAL HAZARDS -
 - i). Entry Restrictions:

Change "..... until dusts have settled." to " until sprays have dried."
2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

This amended labeling supersedes all previously accepted ones. A stamped copy of the labeling is enclosed for your records.

Sincerely yours,
Joanne I. Miller
Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

2/5

Riverdale®

(Commercial label)

HORSEPOWER® 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN
ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, TRICLOPYR AND DICAMBA

KILLS (OR CONTROLS): Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis,
Clover, Cocklebur, Thistle, and many other species of broadleaf weeds; some of which
are listed on this label.

One Quart Covers 8,000 Square Feet

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	10.97%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	1.25%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	1.08%
OTHER INGREDIENTS	<u>86.70%</u>
TOTAL:	100.00%

By Isomer Specific AOAC Method, Equivalent To:

*2-Methyl-4-Chlorophenoxyacetic Acid	8.960%, 0.770 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.896%, 0.077 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	0.896%, 0.077 lbs./gal.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:
JAN 30 2006

EPA REG. NO. 228-335
EPA EST. NO. 228-IL-1

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

228-335

000228-00335.20050915.RED

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and all other handlers must wear: Long-sleeved shirt and long pants, socks and shoes, and chemical resistant gloves made of any waterproof material.

Follow manufacturer's instructions for cleaning/maintaining (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN
Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated areas until dusts have settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

USE PRECAUTIONS

Spray when air is calm to avoid spray drift. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other

4/5

desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable Clovers are present. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum kill (or control) of weeds will be obtained from spring or early fall applications when weeds are actively growing. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede, or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/square feet which does not exceed 3 pints/15,000 square feet. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

Alder	Chicory	Hairy fleabane	Parsnip	Smooth dock
Annual yellow sweet clover	Cinquefoil	Hawkweed	Pearlwort	Smooth pigweed
Artichoke	Clover	Healall	Pennycress	Sneezeweed
Aster	Cockle	Heartleaf drymary	Pennywort	Southern wild rose
Austrian fieldcress	Cocklebur	Hedge bindweed	Peppergrass	Sowthistle
Bedstraw	Coffebean	Hedge mustard	Pepperweed	Spanishneedle
Beggartick	Coffeeweed	Hemp	Pigweed	Spatterdock
Biden	Common chickweed	Henbit	Pineywoods bedstraw	Speedwell
Bindweed	Common mullein	Hoary cress	Plains coreopsis	Spiny Amaranth
Bird vetch	Common sowthistle	Hoary plantain	Plantain	Spiny cocklebur
Bitter wintercress	Corn Chamomile	Hoary vervain	Poison hemlock	Spotted catsear
Bitterweed	Creeping jenny	Honeysuckle	Poison ivy	Spotted knapweed
Black-eyed Susan	Crimson clover	Hop clover	Poison oak	Spotted spurge
Black medic	Croton	Horsenettle	Pokeweed	Spurge
Black mustard	Cudweed	Horsetail	Poorjoe	Spurweed
Black-seed plantain	Curly dock	Indiana mallow	Povertyweed	Stinging nettle
Blessed thistle	Curly indigo	Ironweed	Prickly lettuce	Stinkweed
Blue lettuce	Dandelion	Jewelweed	Prickly sida	Stitchwort
Blue vervain	Dead nettle	Jimsonweed	Primrose	Strawberry clover
Box elder	Dock	Kochia	Prostrate knotweed	Sumac
Bracted plantain	Dogbane	Knawal	Prostrate pigweed	Sunflower
Brassbuttons	Dogfennel	Knotweed	Prostrate spurge	Sweet clover
Bristly oxtongue	Elderberry	Lambsquarter	Prostrate vervain	Tall nettle
Broadleaf dock	English daisy	Lespedeza	Puncture vine	Tall vervain
Broadleaf plantain	Fall dandelion	Locoweed	Purslane	Tansy mustard
Broomweed	False dandelion	Lupine	Ragweed	Tansy ragwort
Buckhorn	False flax	Mallow	Red clover	Tanweed
Buckhorn plantain	False sunflower	Marshelder	Redroot pigweed	Tarweed
Bulbous buttercup	Fiddleneck	Matchweed	Red sorrel	Thistle
Bull nettle	Field bindweed	Mexicanweed	Redstem filaree	Tick trefoil
Bull thistle	Field pansy	Milk vetch	Rough cinquefoil	Toadflax
Burdock	Flea bane (daisy)	Milkweed bloodflower	Rough fleabane	Trailing Crownvetch
Burning nettle	Flixweed	Morningglory	Roundleafed marigold	Tumble mustard
Bur ragweed	Florida betony	Mouseear chickweed	Rush	Tumble pigweed
Burweed	Florida pusley	Mugwort	Russian pigweed	Tumbleweed
Buttercup	Frenchweed	Musk thistle	Russian thistle	Velvet leaf
Canada thistle	Galinsoga	Mustard	St. Johnswort	Venice mallow
Carolina geranium	Garlic mustard	Narrowleaf plantain	Scarlet pimpernel	Veronica
Carpetweed	Goathead	Narrowleaf vetch	Scotch thistle	Vervain
Catchweed bedstraw	Goatsbeard	Nettle	Sheep sorrel	Vetch
Catsear	Goldenrod	Orange hawkweed	Shepherdspurse	Virginia buttonweed
Catnip	Ground ivy	Oxalis	Slender plantain	Virginia creeper
Chickweed	Gumweed	Oxeye daisy	Smallflower galinsoga	Virginia pepperweed
	Hairy bittercress	Parsley-piert	Smartweed	Wavyleaf bullthistle

575

Western clematis	Wild carrot	Wild parsnip	Witchweed	Yellow rocket
Western salsify	Wild four-o'clock	Wild radish	Woolly morningglory	Yellowflower
White clover	Wild garlic	Wild rape	Woodsorrel	pepperweed
White mustard	Wild geranium	Wild strawberry	Woolly croton	and other broadleaf
Wild mustard	Wild lettuce	Wild sweet potato	Woolly plantain	weeds.
Wild aster	Wild marigold	Wild vetch	Wormseed	
Wild buckwheat	Wild onion	Willow	Yarrow	

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 pints in 4 to 50 gallons of water per 8,000 square feet (4 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith.

(RV091505)

Riverdale and Horsepower are registered trademarks of Nufarm Americas Inc.