

228-336

9/27/2002

*EPHA copy
with highlighted
changes 1/3*

RIVERDALE®

HORSEPOWER® SPOT WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, TRICLOPYR AND DICAMBA

KILLS (OR CONTROLS): Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis,
Clover, Cocklebur, Thistle and many other species of
broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	0.740%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	0.084%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.072%
INERT INGREDIENTS:	99.104%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.604%, 0.057 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.060%, 0.006 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	0.060%, 0.006 lbs./gal.

Riverdale and Horsepower Are Registered Trademarks of Riverdale Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-336

NET CONTENTS 1 qt. (32 Fl. Oz.)

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

BURR RIDGE, ILLINOIS 60527-0866

ACCEPTED

SEP 27 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 228-336

Revised 9/6/02
JB

Revised Precautionary Statements and First Aid for oral exposure per
Agency's letter of 8/9/02; also First Aid per PR Notice 2001-1 and
Disposal per PR Notice 2001-6 all via Amendment.

(Side Panels)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Avoid contact with eyes. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HORSEPOWER® SPOT WEED KILLER is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs), and similar non-crop areas.

USE PRECAUTIONS

Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray roots of desirable trees and plants. Maximum kill of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Do not use of Carpetgrass, Dichodra, nor on lawns or turf where desirable clovers are present. Avoid application to Centipede, or St. Augustine grasses unless turf injury can be tolerated. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

WEEDS CONTROLLED BY HORSEPOWER® SPOT WEED KILLER

Alfalfa, Amaranth, Aster, Beadstraw, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campton, Bracken fern, Broomweed, Buckhorn, Buckwheat, Buffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground ivy, Halfshrub sundrop, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle, Henbit, Horsenettle, Horseweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort,

Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red sorrel, Red clover, Redvine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny, Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sumpweed, Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Thistle, Toadflax, Trumpetcreeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild violet, Wild radish, Wild onion, Wild aster, Wild carrot, Wild lettuce, Wild garlic, Wild geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket, and many other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. One application should be sufficient. Some hard-to-kill (or control) broadleaf weeds may require a second (or an additional) treatment in 3 to 4 weeks.

HOW TO USE: Trigger Spray - Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage, raise container.

TO PREPARE REFILL: Add 1.7 ozs. of Riverdale Horsepower Lawn Weed Killer or 2.1 ozs. of Horsepower 8000 Lawn Weed Killer to this empty container. Then add 30 oz. tap water and use according to directions in this label.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in an area inaccessible to children.

PESTICIDE (OR PRODUCT) DISPOSAL: **If empty** - Do not reuse this container except as specified in directions to prepare refill. Place in trash or offer for recycling if available. **If partly filled** - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (RV 090602)

Additional Statement for Sub-Registrants not using the word "Horsepower®" in the product name: "Contains Horsepower® Selective Herbicide."