

228-337

05/26/2000

1/10

(Non-Commercial)

RIVERDALE

XRM-5202 PREMIUM LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN
ORNAMENTAL LAWNS AND TURF GRASSES
And Other Listed Sites

For Both Northern and Southern Grasses
Kills (or Controls) Weeds - Not lawn grasses

CONTAINS 2,4-D, TRICLOPYR, AND CLOPYRALID
One Quart Covers Up to 10,000 Square Feet
(See Directions for Different Grasses.)

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-D*	14.19%
Triethylamine Salt of Triclopyr**	1.06%
Clopyralid***	0.36%
INERT ¹ INGREDIENTS	84.39%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-D	7.61%, 0.660 lbs./gal.
**Triclopyr	0.76%, 0.066 lbs./gal.
***Clopyralid	0.36%, 0.031 lbs./gal.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT²

EPA REG. NO. 228-337

NET CONTENTS


GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



(Side Panels)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, or using the toilet. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried. The maximum number of broadcast applications per treatment site is 2 per year.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

STATEMENT OF PRACTICAL TREATMENT²

- IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if symptoms persist.
- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Most cases of ground water contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

XRM-5202 PREMIUM LAWN WEED KILLER is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf such as Lawns, Parks, Cemeteries, Roadsides, Golf Courses (Fairways, Aprons, and Roughs), and similar non-crop areas. Do not use on Golf Greens nor Tees.

3/10

XRM-5202 may be applied to the following established turfgrass species:

Established Turfgrasses (Common Names)

Northern Grasses (Cool Season)

bentgrass†
bluegrass, Kentucky
fescue, chewing
fescue, creeping red
fescue, sheeps
fescue, tall
ryegrass, perennial

Southern Grasses†† (Warm Season††)

bahiagrass†
bermudagrass†
buffalograss
centipedegrass†
zoysiagrass†
zoysiagrass†
fescue, tall (growing in
warm season areas)

†Do not apply to these turfgrass species unless potential turfgrass injury can be tolerated. Also, when treating these turfgrass species, do not apply more than 4 ozs. per 1000 sq. ft. To minimize grass injury, a second application should be made at least four weeks later. Avoid swath overlaps.

††Do not treat warm season turfgrass with XRM-5202 Premium Lawn Weed Killer when mowing height is less than 1/2 inch.

Note: XRM-5202 Premium Lawn Weed Killer may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, well-established turf that has a well-anchored root system. Do not apply this product to grass species not listed on this label unless potential turf injury can be tolerated.

USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact with cotton, citrus, grapes, tobacco, vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind speed is low and keeping spray boom as low as possible. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Do not pour spray solutions near desirable plants. Do not apply to exposed roots of shallow rooted trees and shrubs. Do not use grass clippings from turf treated with this product for mulch or compost. Mow newly seeded turf at least twice before treatment with XRM-5202. Maximum kill (or control) of weeds will be obtained from Spring or early fall applications when weeds are actively growing. Unless noted elsewhere on this label, do not apply more than 5.25 ozs. per 1,000 sq. ft. per application. Do not exceed recommended rates because damage to turf may occur. Do not apply when temperature exceeds 90°F.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 18 hours, nor should lawns be irrigated (watered) for 18 hours following application period. Watering will wash off weed killing material; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product.

WEEDS CONTROLLED BY XRM-5202 PREMIUM LAWN WEED KILLER

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Acacias, Alder, Artichoke, Aster, Austrian fieldcress, Bachelor button, Beggartick, Bidden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Buckwheat, Buffalobur, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chamomile, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Cornflower, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dog fennel, Elderberry, Fanweed, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Groundsel, Gumweed, Hawksbeard, Hawkweed, Healall, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Horsetweed, Indiana mallow, Ironweed, Jewelweed,

4/10

Jimsonweed, Kochia, Knapweed, Knotweed, Kudzu, Ladysthumb, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Meadow salsify, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nightshade,, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Shepherdspurse, Sicklepod Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Starthistle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansy Ragwort, Tarweed, Teasel, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper, Wild aster, Wild carrot, Wild geranium, Wild lettuce, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill(or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event more than two broadcast applications per year are necessary, use as a spot treatment. Spot treat with ready-to-use XRM-5202 Premium Spot Weed Killer or make up solution as listed below.

TURF APPLICATION DIRECTIONS

Make application using equipment that will ensure uniform coverage such as hose-end sprayer, tank-type pump-up pressure sprayer, or sprinkling can. Broadleaf weeds germinate at different times of the year. Only emerged weeds present at the time of application will be killed (or controlled) by XRM-5202 Premium Lawn Weed Killer. Spray entire lawn, wetting weeds and lawn grasses. Be sure the lawn is well watered. If dry, water 1 to 2 days before applying this product. Application under drought conditions may provide less than desirable results. When hard-to-kill(or control) species are prevalent, when applications are made in late Summer, when weeds are mature, or during drought conditions, use the higher rates. Measure the area to be treated. Multiply the length by the width to calculate the square feet.

Cool Season Grasses - Listed above: use 3.2 to 5.25 ounces of this product in 5 gallons of water to 1,000 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 100 square feet.

Warm Season Grasses - Listed above: use 3.2 to 4 ounces in 5 gallons of water to treat 1,000 square feet (3 teaspoonfuls in 4 quarts of water applied to 150 square feet). If slight turf yellowing occurs, it should disappear after about 1 week.

Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service Weed Control Specialist.

For Refill of Riverdale's Spot Weed Killer: Put 2.8 ounces of XRM-5202 Premium Lawn Weed Killer into the labeled empty quart container. Then add 30 ounces of tap water. Trigger Sprayer - adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. Repeat as necessary.

DISPOSAL SPRAYER AND BOTTLE INSTRUCTIONS

Determine area to be sprayed. Apply at a rate of 32 ounces (1 quart) per 6,100 (5.25 ozs./m sq.ft.) to 10,000 square feet (3.2 ozs./1,000 sq.ft.) for Cool Season grasses or 8,000 (4.0 ozs./1,000 sq.ft.) to 10,000 square feet(3.2 oz./m sq.ft.) for Warm Season grasses. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever tot he "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2,000	Square Feet	4,000
4,000	Square Feet	8,000

(Marks on left for 1 quart disposable bottle. Not to scale.)

6,000	Square Feet	12,000
8,000	Square Feet	16,000
10,000	Square Feet	20,000
Cool Season		Warm Season

STORAGE AND DISPOSAL

STORAGE: To be stored in original container and placed in area inaccessible to children.

PESTICIDE DISPOSAL: Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Rinse thoroughly before discarding in trash. Do not reuse empty container.

NOTICE

Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith.

(ERP 061594) (PR938 042194/RV 021999)

Optional Marketing Statements

- Water Based
- Outdoor Use Only
- For Both Cool Season & Warm Season Grasses

Alternate Statements

- ¹OTHER*
- ²FIRST AID

*Please note, some states still require the term INERT.

6/10

(Commercial)

RIVERDALE

XRM-5202 PREMIUM LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN
ORNAMENTAL LAWNS AND TURF GRASSES
And Other Listed Sites

For Both Northern and Southern Grasses
Kills (or Controls) Weeds - Not lawn grasses

CONTAINS 2,4-D, TRICLOPYR, AND CLOPYRALID
One Quart Covers Up to 10,000 Square Feet
(See Directions for Different Grasses.)

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	14.19%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	1.06%
3,6-Dichloro-2-Pyridinecarboxylic Acid***	0.36%
INERT INGREDIENTS	84.39%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	7.61%	0.660 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.76%	0.066 lbs./gal.
***3,6-Dichloro-2-Pyridinecarboxylic Acid	0.36%	0.031 lbs./gal.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT²

EPA REG. NO. 228-337

NET CONTENTS

GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED

MAY 26 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 228-337

7/10

(Side Panels)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried. The maximum number of broadcast applications per treatment site is 2 per year.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if symptoms persist.
- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Most cases of ground-water contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

XRM-5202 PREMIUM LAWN WEED KILLER is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf such as Lawns, Parks, Cemeteries, Roadsides, Golf Courses (Fairways, Aprons, and Roughs), and similar non-crop areas. Do not use on Golf Greens nor Tees.

8/10

XRM-5202 may be applied to the following established turfgrass species:

Established Turfgrasses (Common Names)

Northern Grasses (Cool Season)

bentgrass†
bluegrass, Kentucky
fescue, chewing
fescue, creeping red
fescue, sheeps
fescue, tall
ryegrass, perennial

Southern Grasses†† (Warm Season††)

bahiagrass†
bermudagrass†
buffalograss
centipedegrass†
zoysiagrass†
zoysiagrass†
fescue, tall (growing in
warm season areas)

†Do not apply to these turfgrass species unless potential turfgrass injury can be tolerated. Also, when treating these turfgrass species, do not apply more than 4 ozs. per 1000 sq. ft. To minimize grass injury, a second application should be made at least four weeks later. Avoid swath overlaps.

††Do not treat warm season turfgrass with XRM-5202 Premium Lawn Weed Killer when mowing height is less than 1/2 inch.

Note: XRM-5202 Premium Lawn Weed Killer may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, well-established turf that has a well-anchored root system. Do not apply this product to grass species not listed on this label unless potential turf injury can be tolerated.

USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact with cotton, citrus, grapes, tobacco, vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind speed is low and keeping spray boom as low as possible. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Do not pour spray solutions near desirable plants. Do not apply to exposed roots of shallow rooted trees and shrubs. Do not use grass clippings from turf treated with this product for mulch or compost. Mow newly seeded turf at least twice before treatment with XRM-5202. Maximum kill (or control) of weeds will be obtained from Spring or early all applications when weeds are actively growing. Unless noted elsewhere on this label, do not apply more than 5.25 ozs. per 1,000 sq. ft. per application. Do not exceed recommended rates because damage to turf may occur. Do not apply when temperature exceeds 90°F.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 18 hours, nor should lawns be irrigated (watered) for 18 hours following application period. Watering will wash off weed killing material; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product.

WEEDS CONTROLLED BY XRM-5202 PREMIUM LAWN WEED KILLER

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Acacias, Alder, Artichoke, Aster, Austrian fieldcress, Bachelor button, Beggartick, Bidden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Buckwheat, Buffalobur, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chamomile, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Cornflower, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dog fennel, Elderberry, Fanweed, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Groundsel, Gumweed, Hawksbeard, Hawkweed, Healall, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Horseweed, Indiana mallow, Ironweed, Jewelweed,

Jimsonweed, Kochia, Knapweed, Knotweed, Kudzu, Ladysthumb, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Meadow salsify, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nightshade,, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Shepherdspurse, Sicklepod Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Starthistle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansy Ragwort, Tarweed, Teasel, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper, Wild aster, Wild carrot, Wild geranium, Wild lettuce, , Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill(or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event more than two broadcast applications per year are necessary, use as a spot treatment. Spot treat with ready-to-use XRM-5202 Premium Spot Weed Killer or make up solution as listed below.

TURF APPLICATION DIRECTIONS

Make application using equipment that will ensure uniform coverage such as hose-end sprayer, tank-type pump-up pressure sprayer, or sprinkling can. Broadleaf weeds germinate at different times of the year. Only emerged weeds present at the time of application will be killed (or controlled) by XRM-5202 Premium Lawn Weed Killer. Spray entire lawn, wetting weeds and lawn grasses. Be sure the lawn is well watered. If dry, water 1 to 2 days before applying this product. Application under drought conditions may provide less than desirable results. When hard-to-kill(or control) species are prevalent, when applications are made in late Summer, when weeds are mature, or during drought conditions, use the higher rates. Measure the area to be treated. Multiply the length by the width to calculate the square feet.

Cool Season Grasses - Listed above: use 3.2 to 5.25 ounces of this product in 5 gallons of water to 1,000 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 100 square feet.

Warm Season Grasses - Listed above: use 3.2 to 4 ounces in 5 gallons of water to treat 1,000 square feet (3 teaspoonfuls in 4 quarts of water applied to 150 square feet). If slight turf yellowing occurs, it should disappear after about 1 week.

Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service Weed Control Specialist.

For Refill of Riverdale's Spot Weed Killer: Put 2.8 ounces of XRM-5202 Premium Lawn Weed Killer into the labeled empty quart container. Then add 30 ounces of tap water. Trigger Sprayer - adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. Repeat as necessary.

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds or other pesticides. Store at temperatures above 32°F. If allowed to freeze, shake well before using. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label.

(ERP 061594) (PR938 042194/RV 021999)

Optional Marketing Statements

- Water Based
- Outdoor Use Only
- For Both Cool Season & Warm Season Grasses