228-340 Please read instructions on re	verse before completing	4-)-	-2-004	Form Approved	1. OMB Nc. 2070	-0060)7
\$epa	Environmenta	nited States I Protectio ngton, DC 204		X	Registration Amendmen Other	OPP identifier Number	
Application for Pesticide - Section I							
1. Company/Product Number 228-340			2. EPA I	Product Manager J. I. Miller		3. Proposed Classification	
4. Company/Product (Name) Riverdale XRM-5202 Premium Weed & Feed			PM#	23		None Resu-icte	nd
5. Name and Address of App Nufarm Americas, Inc Riverdale - A Nufarm 1333 Burr Ridge Parky Burr Ridge, IL 60527 Check if t	Company way, Suite 125A		(b)(i), r to: EPA f		nilar or identical	with FIFRA Section 3(c)(3) in composition and labeling FICATION - 1 2004	_
	· · · · · · · · · · · · · · · · · · ·		Section - I	1			
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.				Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.			
that if this notification is not cons action and penalties under section	la of this product. I under istent with the terms of P is 12 and 14 of FIFRA.	rstand that it is a	violation of 18 U.S.C	. Sec. 1001 to willfu is product may be in	lly make any false st	atement to EPA. I further understan and I may be subject to enforcemen	
1. Material This Product Will Child-Resistant Packaging							-1
* Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt	No. per . container	Yes No If "Yes" Package wgt	No Plastic Glass Paper			_
3. Location of Net Contents Information 4. Size(s) Ret			tail Container 5 Location of Label Directions On Label On Label On Labeling accompanying product				
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Other							
Section - IV							
1. Contact Point (Complete It	ems directly below for	<u> </u>	in the second se	contacted, if neo	cessary, to proces	s this application.)	
Name Russell F. Sawyer			Title Telephone Nc. (Include Area Code) Regulatory Affairs Manager 708/754-3330				
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and completc. I acknowledge that any knowingly faise or misleading statement may be punishable by fine or imprisonment or the complete of							
2. Signature RA Acury			3. Title				
4. Typed Name		5	. Date				
Russell F. Sawyer				February 25, 20	04		
			LAJL 14	PA File Coov (orig	inal) Valley, Applicant Co		

EPA Form 8570-1 IRev. 8-94) Previous editions are obsolete.

PLEASE SEE SPECIAL EDITOR'S NOTE ON LAST PAGE

(Residential Label)

RIVERDALE®



XRM-5202™ PREMIUM WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, TRICLOPYR, AND CLOPYRALID.

Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS: 1.069% Triethylamine Salt of Triclopyr** 0.079% Clopyralid Acid*** 0.028% INERT ¹ INGREDIENTS: 98.824% TOTAL 100.000%
By Isomer Specific AOAC Method, Equivalent to: *2,4-D Acid
GUARANTEED ANALYSIS:° Total Nitrogen (N)
Available Phosphate (P ₂ O ₅)
Chlorine (Cl) Maximum

²IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN NASSAU AND SUFFOLK COUNTIES. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY. IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL*, FOREST*, RIGHT-OF-WAY*, GOLF COURSE OR CEMETERY SITES. *This product cannot be used on Agricultural, Forest sites or Rights-of-Way.

KEEP OUT OF REACH OF CHILDREN CAUTION	7
SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS	

EPA REG. NO. 228-340

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 10/7/03 Added more State Restrictions for California and Oregon via Notification. 2/25/04 Corrected per EPA's e-mail dated 1/21/04 regarding Oregon restrictions via Notification. (Residential)

Page 2

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils that would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

XRM-5202TM PREMIUM WEED AND FEED is a stable mixture of three broad-spectrum herbicides. It is for use on Residential Turf.

This product may be used on established Bahia*, Bermuda*, Blue Grass, Buffalo, Centipede*, Fescue, Rye, and Zoysia* grass. DO NOT USE on Centipede, Dichondra, St. Augustine, and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

*Do not apply to these turfgrasses unless potential turf injury can be tolerated. To minimize grass injury, an additional application should be made no sooner than four weeks later. Avoid overlapping. DO NOT USE GRASS CLIPPINGS FROM TURF TREATED WITH THIS PRODUCT FOR COMPOSTING AND/OR MULCH.

This product is designed to promote lush green growth in lawns and such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black

Riverdale XRM-5202^m Weed and Feed

(Residential)

mustard, Blackseed pantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild fouro'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 (or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 4 (or 3.6 for 18 lb. bag) pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 (.9 for 18 lb. Bag) pound per 250 square feet.

417

(Residential)

SPREADER SETTINGS ONE PASS APPLICATION: °

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing <u>*(No. of lbs.)</u> pounds of nitrogen per 1,000 square feet (optional P_2O_5 and K_2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. *Based on Inert fertilizer claimed.



Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RV 062503B)

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

(Residential)

Page

OPTIONAL MARKETING STATEMENTS

Contains XRM-5202™ - A Premium Selective Herbicide

The following note will not be part of the actual label copy.

"Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

¹OTHER*

*Please note, some states may still require the term INERT and/or the full chemical name of the Active Ingredients.

Editor's Note:

²The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The label prohibitions are optional for all other states.

Special Editor's Note

Not to appear on final printed label. The approved use of this label will automatically be terminated once the Clopyralid Technical, (Labeling approved by The Office of Pesticide Programs (OPP) prior to September 27, 2002) is depleted. Please Note; Clopyralid Technical marketed with labeling approved by OPP on September 27, 2002 (or later) prohibits formulating that Clopyralid Technical for use on residential turf. We will provide the Agency with updated labeling based on the new technical label before packaging. Any questions regarding these statements must be directed to Riverdale's Regulatory Affairs Manager at 1-800/345-3330.



Nufarm Americas Inc. 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527-0866 Phone: 708.754.3330 Fax: 708.754.0314 www.us.nufarm.com

UPS - Overnight February 25, 2004

Document Processing Desk (NOTIF) Registration Division (7505C) Office of Pesticide Programs U.S. Environmental Protection Agency Crystal Mall Building, #2 1921 Jefferson Davis Arlington, VA 22202

Dear Sir/Madam:

Subject: Riverdale XRM-5202 Premium Weed And Feed EPA Registration No. 228-340 Label Revisions Application For Pesticide: Other OPP Identifier No. 292604

We are resubmitting corrected labels. We followed the guidance fact sheet in your January 21, 2004 e-mail covering an identical Oregon label problem.

We have enclosed an extra copy of the label highlighted to show just the changes contained in this Notification.

If there is anything else we should do at this time please contact us immediately.

Thank you for your continuing assistance.

Sincerely yours,

NUFARM AMERICAS INC.

By Russell F. Sawyer Regulatory Affairs Manager

RFS:jb

Enclosures