

PM 23

228-341

10-16-98

10-16



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
228-341

Date of Issuance:  
OCT 16 1998

NOTICE OF PESTICIDE:  
  X   Registration  
       Reregistration

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Riverdale XRM-5202  
Premium Granular Weed  
Killer

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Company  
425 West 194th Street  
Glenwood, IL 60425-1584

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right of exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes before you release the product for shipment [Note: The label changes apply to both the commercial and non-commercial labels except as noted]:

- a. Revise the EPA Registration Number to read "EPA Reg. No. 228-341".
- b. Revise the beginning of the "Hazards To Humans And Domestic Animals" section to read as follows:

"Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes."

The rest of the paragraph, beginning with "After using this product...", is acceptable.

Signature of Approving Official:

Date:  
OCT 16 1998

20710

page 2  
EPA Reg. No. 228-341

- c. Add a First Aid section containing the following ocular first aid statement:

"If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists."

- d. On the non-commercial label only, add the groundwater advisory that is required for all products containing clopyralid. Non-commercial (homeowner) clopyralid products have not been exempted from this requirement:

"Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils that would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater."

- 3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Stamped copies of the labels (Commercial and Non-commercial) are enclosed for your records. Copies of EPA's product chemistry and acute toxicity reviews for this product are also enclosed.

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Enclosures

RD:STANTON:PM Team 23:Rm. 237:CM-2:305-5218:Disk #9:228-GUN.REG

CONCURRENCES

SYMBOL *	7505C							
SURNAME *	S. Stanton							
DATE *	Oct 15, 1998							

(Commercial Label)

30410

RIVERDALE

XRM-5202™ PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH  
KILLS (OR CONTROLS) A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, TRICLOPYR, AND CLOPYRALID.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	2.139%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	0.159%
3,6-Dichloro-2-Pyridinecarboxylic Acid***	0.057%
INERT INGREDIENTS:	97.645%
TOTAL	100.000%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	1.147%
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.114%
***3,6-Dichloro-2-Pyridinecarboxylic Acid	0.057%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.

EPA REG. NO. 228-GUR

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

OCT 16 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-341

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this product will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

XRM-5202™ PREMIUM GRANULAR WEED KILLER is a stable mixture of three broad-spectrum herbicides. It is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, and Golf Courses (Fairways, Aprons, and Roughs).

This product may be used on established Bahia, Bent\* (excluding golf course greens and tees), Bermuda\*, Blue Grass, Buffalo, Centipede\*, Fescue, Rye, and Zoysia\* grass. DO NOT USE on Centipede, Dichondra, St. Augustine, and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

\*Do not apply to these turfgrasses unless potential turf injury can be tolerated. To minimize grass injury, an additional application should be made no sooner than four weeks later. Avoid overlapping.

This product is designed to kill (or control) many broadleaf weeds such as: Acacia, Alder, Artichoke, Aster, Austrian fieldcress, Bachelor button, Beggartick, Bident, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Buckwheat, Buffalobur, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chamomile, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Cornflower, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dog fennel, Elderberry, Fanweed, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Groundsel, Gumweed, Hawksbeard, Hawkweed, Healall, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Horseweed, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knapweed, Knotweed, Kudzu, Ladysthumb, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Meadow salsify, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nightshade, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Shepherdspurse, Sicklepod Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Starthistle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansy Ragwort, Tarweed, Teasel, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper,

Wild aster, Wild carrot, Wild geranium, Wild lettuce, , Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 lb. per 250 square feet.

**SPREADER SETTINGS**

ONE PASS APPLICATION:°

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers or fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains XRM-5202™ - A Premium Selective Herbicide

## Environmental Benefits

(Grasscycling  
Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

For Both Cool Season & Warm Season Grasses

The following note will not be part of the actual label copy.

\*Note: Spreader Settings will vary among various commercial spreader manufacturers.

70410

(Non-Commercial Label)

RIVERDALE

XRM-5202 PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH KILLS (OR CONTROLS) A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, TRICLOPYR, AND CLOPYRALID.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-D*	2.139%
Triethylamine Salt of Triclopyr**	0.159%
Clopyralid Acid***	0.057%
INERT <sup>1</sup> INGREDIENTS	97.645%
TOTAL	100.000%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-D Acid	1.147%
**Triclopyr Acid	0.114%
***Clopyralid Acid	0.057%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.

EPA REG. NO. 228-GUR

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

OCT 16 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-341

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

XRM-5202 PREMIUM GRANULAR WEED KILLER is a stable mixture of three broad-spectrum herbicides. It is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, and Golf Courses (Fairways, Aprons, and Roughs).

This product may be used on established Bahia, Bent\* (excluding golf course greens and tees), common Bermuda\*, Blue Grass, Buffalo, Centipede\*, Fescue, Rye, and Zoysia\* grass. DO NOT USE on Centipede, Dichondra, St. Augustine, and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

\*Do not apply to these turfgrasses unless potential turf injury can be tolerated. To minimize grass injury, an additional application should be made no sooner than four weeks later. Avoid overlapping.

This product is designed to kill (or control) many broadleaf weeds such as: Acacia, Alder, Artichoke, Aster, Austrian fieldcress, Bachelor button, Beggartick, Bidden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Buckwheat, Buffalobur, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chamomile, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Cornflower, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dog fennel, Elderberry, Fanweed, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Groundsel, Gumweed, Hawksbeard, Hawkweed, Healall, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Horseweed, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knapweed, Knotweed, Kudzu, Ladysthumb, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Meadow salsify, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nightshade, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Shepherdspurse, Sicklepod Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Starthistle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansy Ragwort, Tarweed, Teasel, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper, Wild aster, Wild carrot, Wild geranium, Wild lettuce, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after



application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 lb. per 250 square feet.

#### SPREADER SETTINGS

ONE PASS APPLICATION:

#### MAKE

#### SETTING

Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

#### STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides.

**PESTICIDE DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(ERP 061594) (PR938 042194)

#### OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

#### OPTIONAL MARKETING STATEMENTS

Contains XRM-5202™ - A Premium Selective Herbicide

## Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

For Both Cool Season & Warm Season Grasses

Alternate Statements

\*OTHER\*

\*Please note, some states still require the term INERT and/or the full chemical name of the Active Ingredients.

The following note will not be part of the actual label copy.

\*Note: Spreader Settings will vary among various commercial spreader manufacturers.