

228-343

7-12-2002

1/12

Rease read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other

OPP Identifier Number 268495

Application for Pesticide - Section I

1. Company/Product Number 228-343
2. EPA Product Manager J. Miller
3. Proposed Classification None Restricted
4. Company/Product (Name) Riverdale Millennium Ultra™ Weed And Feed
5. Name and Address of Applicant (Include ZIP Code) Riverdale Chemical Company
6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to:

Section - II

Amendment - Explain below. Resubmission in response to Agency letter dated. Notification - Explain below. Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below. NOTIFICATION JUL 12 2002

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Adding the optional statement "In Washington, this product may not be used on residential lawns. In Massachusetts, do not use grass clippings from turf treated with this product for composting and/or mulch in the growing season of application."

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging, 2. Type of Container, 3. Location of Net Contents Information, 4. Size(s) Retail Container, 5. Location of Label Directions, 6. Manner in Which Label is Affixed to Product

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name: Russell F. Sawyer, Title: Regulatory Affairs Manager, Telephone No. (Include Area Code): 708/754-3330

Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. 2. Signature: Russell F. Sawyer, 3. Title: Regulatory Affairs Manager, 4. Typed Name: Russell F. Sawyer, 5. Date: July 3, 2002, 6. Date Application Received (Stamped)

2/12

NOTIFICATION
JUL 12 2002

(Non-Commercial Label)

RIVERDALE

MILLENNIUM ULTRA™ WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-D*	0.690%
Monoethanolamine Salt of Clopyralid**	0.094%
Dimethylamine Salt of Dicamba***	0.086%
INERT ¹ INGREDIENTS:	<u>99.130%</u>
TOTAL	100.000%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-D Acid	0.573%
**Clopyralid Acid	0.071%
***Dicamba Acid	0.071%

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	

Fertilizer guaranteed by (Name of fertilizer registrant)

Riverdale is a Registered Trademark of Riverdale Chemical Company

† Millennium Ultra™ - A Trademark of Riverdale Chemical Company

Patent No. 5,266,553

¹IN WASHINGTON, THIS PRODUCT MAY NOT BE USED ON RESIDENTIAL LAWNS.
IN MASSACHUSETTS, DO NOT USE GRASS CLIPPINGS FROM TURF TREATED WITH THIS PRODUCT FOR COMPOSTING AND/OR MULCH IN THE GROWING SEASON OF APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-343

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
BURR RIDGE, ILLINOIS 60527-0866

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

STATEMENT OF PRACTICAL TREATMENT²

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRA™ WEED AND FEED is for use on Ornamental Turf such as Lawns, Golf Courses (Fairways, Aprons, Tees* and Roughs), Parks, Cemeteries and similar non-crop areas.

*Excluding Bentgrass Tees.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Alfalfa, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitter wintercress, Bitterweed, Black medic, Blackeyed Susan, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckbrush, Buckhorn, Buffalobur, Bull thistle, Bur ragweed, Burdock, Burhead, Buttercup, Canada thistle, Carpetweed, Catnip, Chamise, Cherokee rose, Chickweed, Chicory, Cinquefoil, Clover, Coastal redstem sage, Cockle, Cocklebur, Coffee bean, Coffeeweed, Common groundsel, Common sowthistle, Common sunflower, Cornflower (bachelor button), Corn speedwell, Coyotebrush, Creeping jenny, Croton, Curly Indigo, Dandelion, Devil's claw, Dock, Dogbane, Dogfennel, Elderberry, English daisy, False Chamomile (scentless), Fanweed, Fiddle neck, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Goosefoot, Ground ivy, Gumweed, Halogeton, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horestail, Horsetweed, Indiana mallow, Indigo, Ironweed, Jerusalem artichoke, Jewelweed, Jimsonweed, Klamathweed, Knapweed (diffuse, Russian, & spotted), Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Locoweed, Lupine, Mallow, Many

flowered aster, Manzanita, Marijuana, Marshelder, Matchweed, Meadow salsify (goatsbeard), Mexican weed, Milkvetch, Morningglory, Musk thistle, Mustard, Narrowleaf hawksbeard, Nettle, Nightshade (cutleaf & hairy), Orange hawkweed, Oxalis (stricta and corniculata), Parrotfeather, Parsley-piert, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed,

Pigweed, Pineappleweed, Pinnate tansymustard, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Rabbitbrush, Ragweed, Red clover, Red sorrel, Redroot pigweed, Redstem, Rush, Russian thistle, Salsify, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiderwort, Spurge, Spotted catsear, Spurweed, St. Johnswort, Starthistle, Starthistle, Stinging nettle, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tansymustard, Tansyragwort, Tarweed, Texas blueweed, Thistle, Toadflax, Tumbleweed, Velvetleaf, Veronica, Vervain, Vetch, Virginia creeper, White clover, Wild aster, Wild buckwheat, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild violet, Witchweed, Wormseed, Wormwood, Woodsorrel, Yarrow, Yellow rocket, Yellow starthistle, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 (or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 (or 3.6 for 18 lb. bag) pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 (.9 for 18 lb. bag) pound per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION: °

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 110398C)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra™ - A Premium Selective Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

¹Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

- ¹OTHER*
- ²FIRST AID

*Please note, some states may still require the term INERT and/or the full chemical name of the Active Ingredients.

Editor's Note:

³The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of New York, Washington and/or Massachusetts. The label prohibitions are optional for all other states.

Additional marketing and fertilizer hype:

- **CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS**
- **TIME-RELEASED NITROGEN PLUS WEED CONTROL FOR A GREEN, WEED-FREE LAWN**
- **CONTAINS IRON & SULFUR FOR EXTRA DEEP GREENING**
- **FOR FALL APPLICATION - CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS**
- **LAWN FOOD PROMOTES QUICK GREENING IN THE SPRING**

**CONTAINS
MILLENNIUM ULTRA™ HERBICIDE**

<p>X NITROGEN: Controlled Slow Release for growing a dense, green lawn</p>	<p>X PHOSPHORUS: Improves root growth & plant food storage capabilities</p>	<p>X POTASSIUM: Vital for growth & improves hardiness & drought resistance</p>
--	---	--

START YOUR YEAR ROUND LAWN PROGRAM NOW! See back for details

SATISFACTION
GUARANTEED

(SEAL)

If for any reason, you are not completely satisfied with the results after applying this product, you are entitled to product replacement or your money back simply by presenting evidence of purchase to your local (Seller Location)

Early Spring

Crabgrass
Preventer
plus Lawn
Food

Late Spring

Lawn Food or
Weed & Feed

(Seller's Name)

**YEAR ROUND
LAWN CARE
PROGRAM**

Fall

Winter Lawn
Food or
Winter Weed &
Feed

Summer

Super Fortifier
Plus
and/or
Insect Control

Alternate Spreader Information

Do not use your hands to spread this product.

Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.

BROADCAST SPREADER SETTINGS (One Pass Application)

For best results, apply in a circular path starting on the outside and working toward the center(see Path A). A little overlap of product will ensure complete coverage.

	3.6# Rate	
Earl May Broadcast.....	6	<hr style="border: 1px solid black;"/> <p>(Illustration)</p> <hr style="border: 1px solid black;"/>
Earl May (SB4000P).....	6	
Precision Tractor TBS-90.....	6	
Republic EZ 540/75450..... 3-4		
Scotts EasyGreen..... 26½ - 27½		
Scotts SpeedyGreen.....	6	

DROP SPREADER SETTINGS (One Pass Application)

First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

	3.6# Rate	
Earl May Drop.....	6½ - 7	<hr style="border: 1px solid black;"/> <p>(Illustration)</p> <hr style="border: 1px solid black;"/>
Republic EZ 530/75350.....	7 - 8	
Scotts Accugreen.....	6½	
Scotts PF4 Metal.....	5 - 6½	

Visit our web site at: www.xxx.com

A fertilizer.....formulated to the highest lawn care standards.
Attacks weeds in the fall when they are most vulnerable.

(Commercial Label)

RIVERDALE[®]

MILLENNIUM ULTRA™ WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.
Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.690%
Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid**	0.094%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.086%
INERT INGREDIENTS:	99.130%
TOTAL	100.000%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	0.573%
**3,6-Dichloro-2-Pyridinecarboxylic Acid	0.071%
***3,6-Dichloro-o-Anisic Acid	0.071%

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	
Fertilizer guaranteed by (Name of fertilizer registrant)	

Riverdale is a Registered Trademark of Riverdale Chemical Company

† Millennium Ultra™ is a Trademark of Riverdale Chemical Company Patent No. 5,266,553

¹IN WASHINGTON, THIS PRODUCT MAY NOT BE USED ON RESIDENTIAL LAWNS.
IN MASSACHUSETTS, DO NOT USE GRASS CLIPPINGS FROM TURF TREATED WITH THIS PRODUCT FOR COMPOSTING AND/OR MULCH IN THE GROWING SEASON OF APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-343

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
BURR RIDGE, ILLINOIS 60527-0866

8/12
NOTIFICATION
JUL 12 2002

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

MILLENNIUM ULTRA™ WEED AND FEED is for use on Ornamental Turf such as Lawns, Golf Courses (Fairways, Aprons, Tees* and Roughs), Parks, Cemeteries and similar non-crop areas.
*Excluding Bentgrass Tees.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Alfalfa, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitter wintercress, Bitterweed, Black medic, Blackeyed Susan, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckbrush, Buckhorn, Buffalobur, Bull thistle, Bur ragweed, Burdock, Burhead, Buttercup, Canada thistle, Carpetweed, Catnip, Chamise, Cherokee rose, Chickweed, Chicory, Cinquefoil, Clover, Coastal redstem sage, Cockle, Cocklebur, Coffee bean, Coffeeweed, Common groundsel, Common sowthistle, Common sunflower, Cornflower (bachelor button), Corn speedwell, Coyotebrush, Creeping jenny, Croton, Curly Indigo, Dandelion, Devil's claw, Dock, Dogbane, Dogfennel, Elderberry, English daisy, False Chamomile (scentless), Fanweed, Fiddle neck, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Goosefoot, Ground ivy, Gumweed,

Halogeton, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horestail, Horsetweed, Indiana mallow, Indigo, Ironweed, Jerusalem artichoke, Jewelweed, Jimsonweed, Klamathweed, Knapweed (diffuse, Russian, & spotted), Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Locoweed, Lupine, Mallow, Many flowered aster, Manzanita, Marijuana, Marshelder, Matchweed, Meadow salsify (goatsbeard), Mexican weed, Milkvetch, Morningglory, Musk thistle, Mustard, Narrowleaf hawksbeard, Nettle, Nightshade (cutleaf & hairy), Orange hawkweed, Oxalis (stricta and corniculata), Parrotfeather, Parsley-piert, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineappleweed, Pinnate tansymustard, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Rabbitbrush, Ragweed, Red clover, Red sorrel, Redroot pigweed, Redstem, Rush, Russian thistle, Salsify, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiderwort, Spurge, Spotted catsear, Spurweed, St. Johnswort, Starthistle, Stinging nettle, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tansymustard, Tansyragwort, Tarweed, Texas blueweed, Thistle, Toadflax, Tumbleweed, Velvetleaf, Veronica, Vervain, Vetch, Virginia creeper, White clover, Wild aster, Wild buckwheat, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild violet, Witchweed, Wormseed, Wormwood, Woodsorrel, Yarrow, Yellow rocket, Yellow starthistle, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18)pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag)pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:°

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not

provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 10038C)

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra™ - A Premium Selective Herbicide

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

*Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Editor's Note:

*The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states Washington and/or Massachusetts. The label prohibitions are optional for all other states.