Please read instructions on reverse before completing form. Form Approved, OMB No. 26 OPP Identifier Number Registration **United States SEPA Environmental Protection Agency Amendment** 292562 Washington, DC 20460 Other Application for Pesticide - Section I 2. EPA Product Manager 3. Proposed Classific tion 1. Company/Product Number 228-343 J. I. Miller Restric None 4. Company/Product (Name) PM# Riverdale Millennium Ultra Weed And Feed 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) Nufarm Americas, Inc. (b)(i), my product is similar or identical in composition and labeling Riverdale - A Nufarm Company 1333 Burr Ridge Parkway, Suite 125A EPA File Symbol Burr Ridge, IL 60527-0866 Check if this is a new address **Product Name** Section - II Amendment - Explain below. Final printed labels in response to Agency letter dated Resubmission in response to Agency letter dated \_\_\_ "Me Too" Application. DCT 2 8 2003 Other - Explain below. Notification - Explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) Added more state restrictions (California and Oregon) and added more weeds. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec, 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Section - III 1. Material This Product Will Be Packaged In: Child-Resistant Packaging **Unit Packaging** 2. Type of Container Water Soluble Packaging Yes ' Metal Yes Yes **Plastic** No No No Glass If "Yes" Paper If "Yes" No. per No. per \* Certification must Unit Packaging wgt. container Other (Specify) Package wgt container be submitted 3. Location of Net Contents Information Location of Label Directions 4. Size(s) Retail Container On Label Container On Labeling accompanying product Lithograph 6. Manner in Which Label is Affixed to Product Other Paper glued Stenciled Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name Telephone Nc. (Include Area Code) 708/754-3330 Russell F. Sawyer Regulatory Affairs Manager 6. Date Application Certification Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable low. 3. Title 2. Signature

Russell F. Sawyer

4. Typed Name

Regulatory Affairs Manager

October 7, 2003

5. Date

# PLEASE SEE SPECIAL EDITOR'S NOTE ON LAST PAGE

## (Residential Label)

# RIVERDALE® MILLENNIUM ULTRA™ WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

# CONTAINS 2,4-D, CLOPYRALID AND DICAMBA Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-D* Monoethanolamine Salt of Clopyralid** Dimethylamine Salt of Dicamba***  INERT INGREDIENTS:  TOTAL  0.690% 0.094% 0.086% 100.000%
By Isomer Specific AOAC Method, Equivalent to:
GUARANTEED ANALYSIS: Total Nitrogen (N)
X% Urea Nitrogen Available Phosphate $(P_2O_5)$
Chlorine (Cl) Maximum
Fertilizer guaranteed by (Name of fertilizer registrant) Riverdale is a Registered Trademark and Millennium Ultra is a Trademark of Riverdale - A Nufarm Company

IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN NASSAU AND SUFFOLK COUNTIES.

IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL, FOREST, RIGHT-OF-WAY, GOLF COURSE OR CEMETERY SITES. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY.

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-343

NET WEIGHT LBS.

EPA EST. NO. 228-11-1

# MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

- Revised 4/4/03 Changed First Aid Statement per PR 2001-1 and the Non-Commercial Disposal Statement per PR 2001-6 via Amendment.
  - 6/25/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting. Also added special Editor's Note.
  - 10/7/03 Added more State Restrictions for California and Oregon and added more weeds via Notification.

3/4

Page 2

of v

## PRECAUTIONARY STATEMENTS

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

# CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

# FIRST AID STATEMENT AND A SECOND SECOND OF THE STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN . OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

## ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

# DIRECTIONS FOR USE STATEGIA BY THE ACC

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRAM WEED AND FEED is for use on Residential Turf, except as noted on the front cover.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperatura exceeds 90°F. Avoid overlapping. Do not use grass clippings from turf treated with this product for composting and/or mulch.

Whis product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyeld Susan, Black medic, Black mustard, Blackseed pantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Puttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean,

Page 3 V

Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag) pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.

# Riverdale Millennium Ultram Weed and Feed (Residential)

Page 4 of

SPREAD	ER	SE	ľľ	ING	S		
ONE	PAS	S	λP	PLI	CAT	ION	: 0

<u>MAKE</u>	<u>SETTING</u>
	and the state of t
Central Calibrated	
Cyclone Broadcast	3-3/4
Jackson	<u></u>
Sears Broadcast	ar Tay County (1986)
Sears	Name (All Cartholic 1995) (1997)
Scotts Calibrated	- hr 150 s 4

The Land Control of the Control of the

THE SHARE SWEETER OF STORY

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing  $\frac{*(No.\ of\ lbs.)}{*(No.\ of\ lbs.)}$  pounds of nitrogen per 1,000 square feet (optional  $P_2O_5$  and  $K_2O$  levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

# Environmental Benefits (crosscycling)

Logo)

Besides increasing properly values, did you know a well-maintained lawn provides the following benefits to your home and environment:

April 1985 Annie

in the production of the indicate of the contract of the contr

- Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.
- ✓ A 50' x 50' lawn produces enough axygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifles Water & Modifles Temperature
- ✓ Improves Soil Condition & Controls Erosion

# STORAGE AND DISPOSAL COMPANY OF THE

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

**DISPOSAL:** If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Saller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 100703)

# 'Riverdale Millennium Ultram Weed and Feed (Residential)

age 5

# OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

#### OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra™ - A Premium Selective Herbicide

The following note will not be part of the actual label copy.

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

¹OTHER\*

\*Please note, some states may still require the term INERT and/or the full chemical name of the Active Ingredients.

### Editor's Note:

'The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The state label prohibition is optional for all other states.

Additional marketing and fertilizer hype:

- CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS
- TIME-RELEASED NITROGEN PLUS WEED CONTROL FOR A GREEN, WEED-FREE LAWN
- CONTAINS IRON & SULFUR FOR EXTRA DEEP GREENING
- FOR FALL APPLICATION CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS
- LAWN FOOD PROMOTES QUICK GREENING IN THE SPRING

CONTAINS
MILLENNIUM ULTRAPH HERBICIDE

Riverdale Millennium Ultra™ Weed and Feed (Residential)

x NITROGEN: Controlled Slow Release for growing a dense, green lawn

x PHOSPHORUS: Improves root growth & plant food storage capabilities

x POTASSIUM: Vital for growth & improves hardiness & drought resistance

· 2 大概是實施 (新聞) (新聞) (新聞) (新聞)

START YOUR YEAR ROUND LAWN PROGRAM NOW! See back for details

> it could be interpreted to the same of the i milita example

> > aldeen tel

'n

SATISFACTION **GUARANTEED** 

(SEAL)

If for any reason, you are not completely " " satisfied with the results after applying this product, you are entitled to product replacement or your money back simply by presenting evidence of purchase to your local (Seller Location)

and the state of t

Early Spring Crabgrass Preventer plus Lawn Food

Late Spring Lawn Food or Weed & Feed

(Seller's Name)

YEAR ROUND LAWN CARE **PROGRAM** 

Fall Winter Lawn Food or Winter Weed & Feed

Super Fortifier Plus and/or Insect Control

Alternate Spreader Information

Do not use your hands to spread this product.

Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage or the lawn.

BROADCAST SPREADER SETTINGS (One Pass Application) For best results, apply in a circular path starting on the outside and working toward the

# 'Riverdale Millennium Ultra Weed and Feed (Residential)

Page 7 or

center(see Path A). A little overlap of product will ensure complete coverage.

	3.6# Rate	
Earl May Broadcast	6	
Earl May (SB4000P)	6	
Precision Tractor TBS-90	(Illustration)	
Republic EZ 540/75450	3-4	(IIIustration)
Scotts EasyGreen	26 <del>¼</del> - 27 <del>¼</del>	
Scotts SpeedyGreen	6	

DROP SPREADER SETTINGS (One Pass Application)

First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

	3.6# Rate	
Earl May Drop	6½ - 7	
Republic EZ 530/75350	7 - 8	(Illustration)
Scotts Accugreen	6 <del>½</del>	
Scotts PF4 Metal	5 - 61/2	

Visit our web site at: www.xxx.com

A fertilizer.....formulated to the highest lawn care standards. Attacks weeds in the fall when they are most vulnerable.

Special Editor's Note

Not to appear on final printed label.

The approved use of this label will automatically be terminated once the Clopyralid Technical, (Labeling approved by The Office of Pesticide Programs (OPP) prior to September 27, 2002) is depleted. Please Note; Clopyralid Technical marketed with labeling approved by OPP on September 27, 2002 (or later) prohibits formulating that Clopyralid Technical for use on residential turf. We will provide the Agency with updated labeling based on the new technical label before packaging. Any questions regarding these statements must be directed to Riverdale's Regulatory Affairs Manager at 1-800/345-3330.

# PLEASE SEE SPECIAL EDITOR'S NOTE ON LAST PAGE

(Commercial Label)

RIVERDALE\* MILLENNIUM ULTRA™ WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

# CONTAINS 2,4-D, CLOPYRALID AND DICAMBA Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Square Feet (50' x 100')

CTIVE INGREDIENTS:	
	90€
Monoethanolamine Salt of Clopyralid**	948
Dimethylamine Salt of Dicamba***	868
NERT INGREDIENTS:	
TOTAL	
By Isomer Specific AOAC Method, Equivalent to:	
*2,4-D Acid	73%
**Clopyralid Acid	71%
	71%
GUARANTEED ANALYSIS:	
Total Nitrogen (N)	X₹
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P2O5)	Хθ
Soluble Potash (K <sub>2</sub> O)	
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Trip	
Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	<b>7</b> %
The urea nitrogen in this product has been coated to provide X% coated slow release	
urea nitrogen.	=
Fertilizer guaranteed by (Name of fertilizer registrant)	
Millennium Ultra is a Trademark and Riverdale is a Registered Trademark of Riverdale -	A
Nufarm Company	

2 IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN NASSAU AND SUFFOLK COUNTIES. IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL, FOREST, RIGHT-OF-WAY, GOLF COURSE OR CEMETERY SITES. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**KPA REG. NO. 228-343** 

NET WEIGHT LBS. EPA EST. NO. 228-IL-1

# MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 4/1/03 Changed First Aid Statement per PR 2001-1 via Amendment.

6/25/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting. Also added Special Editor's Note.

10/7/C3 Added more State Restrictions for California and Oregon and added more weeds via Notification.

278 od 12/14

Riverdale Millennium Ultra Weed and Feed (Commercial)

## PRECAUTIONARY STATEMENTS

Page 2 14

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

### FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of ground water contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRAM WEED AND FEED is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees\* and Roughs) and similar turf areas, except as noted on front cover. \*Excluding Bentgrass Tees.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, when air composed temperature exceeds 90°F. Avoid overlapping. Do not use grass clippings from turn treated with this product for composting and/or mulch.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyed Susan, Black medic, Black mustard, Blackseed pantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly

Page 3 10

oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

when to APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: Whell applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested mate of 4(or 3.6 for 18 lb. bag) pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18.1b. bag) pound per 250 square feet.

# Riverdale Millennium Ultra™ Weed and Feed (Commercial)

Į	J	-114
Page	4	all
	P	2 2

SPREADER SETTINGS ONE PASS APPLICATION: O	MAKE	SETTING
	Central Calibrated	5
	Cyclone Broadcast	3-3/4
	Jackson	5
	Sears Broadcast	3 .
	Sears	5
	Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing  $\frac{*(No.\ of\ lbs.)}{}$  pounds of nitrogen per 1,000 square feet (optional  $P_2O_5$  and  $K_2O$  levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. \*Based on Inert fertilizer claimed.

# Environmental Benefits

(Grasscycling Logo) Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

- Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lown. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.
- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✔ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

# STORAGE AND DISPOSAL

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

**PRODUCT DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 100703)

# Riverdale Millennium Ultra™ Weed and Feed (Commercial)



Ç.

# OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

#### OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultram - A Premium Selective Herbicide

The following note will not be part of the actual label copy.

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

1OTHER\*

\*Please note, some states may still require the term INERT and/or the full chemical name of the Active Ingredients.

# Editor's Note:

The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The state label prohibition is optional for all other states.

# Special Editor's Note

Not to appear on final printed label. The approved use of this label will automatically be terminated once the Clopyralid Technical, (Labeling approved by The Office of Pesticide Programs (OPP) prior to September 27, 2002) is depleted. Please Note; Clopyralid Technical marketed with labeling approved by OPP on September 27, 2002 (or later) prohibits formulating that Clopyralid Technical for use on residential turf. We will provide the Agency with updated labeling based on the new technical label before packaging. Any questions regarding these statements must be directed to Riverdale's Regulatory Affairs Manager at 1-800/345-3330.

estrected to



14/14

Regulatory Affairs Department 1333 Burr Ridge Parkway, Suite 125A

Burr Ridge IL 60527-0866 Office Ph: 708 754 3330 Facsimile: 708 754 0314

UPS - Overnight October 7, 2003

Document Processing Desk (NOTIF) Registration Division (7505C) Office of Pesticide Programs U.S. Environmental Protection Agency Crystal Mall Building, #2 1921 Jefferson Davis Arlington, VA 22202

Dear Sir/Madam:

Subject: Riverdale Millennium Ultra Weed And Feed

EPA Registration No. 228-343 Application For Pesticide: Other OPP Identifier No. 292562

The states of California and Oregon have requested we modify our labels to include their restrictions regarding the use of Clopyralid. We also added more weed names.

We have also enclosed an extra copy of each label highlighted to show just the changes contained in this submission.

If there is anything else we should do at this time please contact us immediately.

Thank you for your continuing assistance.

Sincerely yours,

RIVERDALE - A NUFARM COMPANY -

By Russell F. Sawyer ()
Regulatory Affairs Manager

RFS:jb

**Enclosures**