UNITED STATES	Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W.	Number: 228-348	Date of Issuance: JUN 30 1998
A GERIC	Washington, D.C. 20460	Term of Issuance:	
THUNAL PROTECTOR NOTI	CE OF PESTICIDE:	Conditiona	L
	<u>x</u> Registration Reregistration	Name of Pessicide	Froduct:
under FIFRA, as amended;		Riverdale I Weed and Fe	Dissolve LBN eed
ame and Address of Regist Riverdale Chemic			
25 West 194 <sup>th</sup> Si Slenwood, IL 604	t.		
e submitted to and accept	differing in substance from that accepted by the Registration Division prior t duct alws, a refer to the above ERA regi	: use of the label in comme	
	n furnissed by the registrant, the above der the Federal Insecticide, Fungicide		
n order to protect nealth ancel the registration of with the registration of a	to be construed as an endorsement of ro- and the evolutient, the Administrator a pesticle in accordance with the Art sproduct other this Act is not to be pro- or to its use of it has been covered of	in his motich, may at any The acceptance of any ha estrued as giving the regis	time suspend in re in connection
	ct is conditionally regi (7)(A) provided that you		ance with
	and/or cite all data re der FIFRA sec. 3(c)(5) wi		
registrants of s acceptable respo	similar products to subm onses required for rereg	it such data; and	d submit
registrants of s acceptable respo ander FIFRA sect	similar products to subm onses required for rereg	it such data; and istration of you:	d submit
registrants of s acceptable respo ander FIFRA sect 2. Make th a. Revis	similar products to subm onses required for rereg tion 4.	it such data; and istration of you: es:	d submit r product
registrants of s acceptable respo under FIFRA sect 2. Make th a. Revis No. 2 3. Add the fresh a	similar products to submonses required for reregation 4. The following label changes the EPA Registration 1228-345". The following inhalation states of the second states of the	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial	d submit r product "EPA Reg. e victim tc
registrants of s acceptable responder FIFRA sect 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti	similar products to submonses required for reregion 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, and a station, preferably mouth-form.	it such data; and istration of you es: Number to read, tatement: "Remove give artificial to-mouth. Get me	d submit r product "EPA Reg. e victim tc edical
registrants of s acceptable responder FIFRA sect 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for reregion 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, and a station, preferably mouth-form.	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for
registrants of s acceptable responder 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for reregation 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, water in the set of the set of the revised of the	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for
registrants of s acceptable responder FIFRA sect 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for reregation 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, water in the set of the set of the revised of the	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for
registrants of s acceptable responder 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for reregation 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, water in the set of the set of the revised of the	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for
registrants of s acceptable responder FIFRA sect 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for reregation 4. The following label changes the EPA Registration 1228-345". The following inhalation stair. If not breathing, water in the set of the set of the revised of the	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for
registrants of s acceptable responder 2. Make th a. Revis No. 2 3. Add the fresh a respira attenti 4. Submit the rec	similar products to submonses required for rereg- tion 4. The following label chang- se the EPA Registration 1 228-345". The following inhalation states and the second states of the revis- tion, preferably mouth- ton. two copies of the revis- cord.	it such data; and istration of you: es: Number to read, tatement: "Remove give artificial to-mouth. Get me ed final printed	d submit r product "EPA Reg. e victim tc edical label for

• ,

1

page 2 EPA Reg. No.228-348

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

s4.

Joanne I. Miller Product Manger (23) Herbicide Branch Registration Division (7505C)

Enclosure

ĺ

Į

1

Ξ.

# RIVERDALE

# DISSOLVE® LBN WEED AND FEED

· .

# THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION. Contains 2,4-D, Mecoprop and Dichlorprop

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

	ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid INERT INGREDIENTS:	c Acid** d***	0.327% 0.322% 
	<pre>ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO: *2,4-Dichlorophenoxyacetic Acid</pre>	• • • • • • • • • • •	0.52% 0.27% C.27%
(	UARANTEED ANALYSIS:° Total Nitrogen (N)		
	Available Phosphate( $P_2O_5$ )		X% X%
	Primary Plant Foods derived from Ammonium Phosphate, Sul Super Phosphate, and Muriate of Potash.	fur Coated Urea,	Urea, Triple
	Chlorine (Cl) Maximum		X%
	The urea nitrogen in this product has been coated to pro urea nitrogen.	vide X% coated sl	ow release "
	Fertilizer guaranteed by (Name of fertilizer registrant).		
	<code>†DISSOLVE®</code> - A Registered Trademark of Riverdale Chemical Com	mpany Patent	No. 5,266,553
/	KEEP OUT OF REACH OF CHILDREN		
l	CAUTION		4
	SEE BACK PANEL FOR ADDITIONAL PRECAUTIONAR AND STATEMENT OF PRACTICAL TREATM		
	NET WEIGHT 16 LBS.		••••
	EPA REG. NO. 228-GUT	EPA EST.	NO
	MANUFACTURED BY	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	
	RIVERDALE CHEMICAL COMPANY	4 4 4 7 1 4 4 4	• • · · • • • •
	GLENWOOD, ILLINOIS 60425-1584		•
		ACCEPTED with COMMENTS In EPA Letter Date	di yang kanalari yang kana Al
		JUN 30 1998	<b>C</b> -
		Under the Federal Inse Fundicide, and Rodensic as amended, for the p registered under EPA R 2.28 348	esticide

.

.

Riverdale Dissolve® LBN Weed and Feed (Commercial)



### (Back Panel)

4

#### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dist and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks, shoes and gloves when loading and applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention. IF ON SKIN: Wash skin with soap and water. Get medical attention.

### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater in the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP, and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervicus pad to contain spills will help prevent groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH (LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DISSOLVE<sup>®</sup> LBN WEED AND FEED is a stable mixture of three herbicides. It is for use ca Ornamental Turf, such as Lawns, Parks, Cemeteries, Golf Courses (Fairway, Aprons, Tees and Roughs).

This product ray be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Biden, Bindweed, Bitterweed, Bitter, Wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elferberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian

. . . . . .

Riverdale Dissolve<sup>®</sup> LBN Weed and Feed (Commercial)

thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

100 and 10

100 100 - **1**4

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high remperature and high humidity, over-application may cause minor transitory discoloration n Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader cn 250 square feet and adjust to apply at the rate of .8 pound per 250 square feet.

SPREADER SETTINGS CNE PASS APPLICATION:°

ĺ

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	· 4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement you regular fertilizer program by providing  $\underline{*}(No. of lbs.)$  pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O, and K<sub>2</sub>O\*levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. \*Based on Inert Fertilizer claimed.

#### STORAGE AND DISPOSAL

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or

....



 $\mathcal{U}\mathcal{U}^{\prime}$ 

feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (EEP 061594) (PR938 042194)

# OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and (shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS:

Contains Dissolve® A Premium Broadleaf Herbicide



. . . .

 Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose mare readily so nutrients go back into your lown.

 A 50° x 50° lawn produces enough oxygen for a family of 4 every day

- Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Poüutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

# RIVERDALE

# DISSOLVE® LBN WEED AND FEED

# THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION. CONTAINS 2,4-D, MECOPROP AND DICHLORPROP

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

	ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-D*		0.626%
	Dimethylamine Salt of Mecoprop**		0.327%
	Dimethylamine Salt of Dichlorprop***		0.322%
	INERT' INGREDIENTS:		<u>98.725%</u>
	TCTAL	• •	100.000%
	ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:		
	*2,4-D Azid		
	**Mecoprop Acid		
			. 0.276
l,	JARANTEED ANALYSIS:		
	Total Nitrogen (N)		X%
	X% Ammoniacal Nitrogen X% Urea Nitrogen		
	Available Phosphate (P <sub>2</sub> O <sub>5</sub> )		X%
	Soluble Potash(K,O)		
	Primary Flant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Super Phosphate, and Muriate of Potash.	Urea	a, Triple
	•	4 / 1 1	,
	Chlorine (Cl) Maximum	<b>.</b>	
	The urea nitrogen in this product has been coated to provide X% coated s urea nitrogen.	low	release
	Fertilizer guaranteed by (Name of fertilizer registrant)	i	
	tDISSOLVE® - A Registered Trademark of Riverdale Chemical Company	No,	5,266,553
	·	د . <del>د</del> د	(
Ę	KEEP OUT OF REACH OF CHILDREN		
	CAUTION		
	SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT		
	AND STATEMENT OF PRACTICAL TREATMENT NET WEIGHT 16 LBS.	NO.	228-IL-1
	AND STATEMENT OF PRACTICAL TREATMENT NET WEIGHT 16 LBS.	NO.	228-IL-1

# MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

ACCEPTED whb COMMENTS In EPA Letter Dated JUN 30 1998

4

-

14

٦

GLENWOOD, ILLINOIS 60425-1584 

.

MENTS ... c.'s Letter Dated

Under the Federal Insecticide, Fundicide, and Rodenticide Act as anended, for the posticide registered ander EPA Reg. No. 228 - 348

Under the Federal Insecticide, Sundicide and Rodenticide Act ∿q.

ເປ

0//0

Riverdale Dissolve® LBN Weed and Feed (Non-Commercial)

# Page 2

「日本の日本の

# (Back Panel)

#### PRECAUTIONARY STATEMENTS

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye injury. Harmful if absorbed through skin. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

#### STATEMENT OF PRACTICAL TREATMENT<sup>2</sup>

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
Wash skin with soap and water. Get medical attention.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

•••••

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DISSOLVE® 'L'EN'WEED AND FEED is a stable mixture of three herbicides. It is for use on Ornamental' 'Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees, and Roughs)' and similar non-crop areas.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this ( product where desirable clovers are present or allow the product to "wind drift" onto flowers, Vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Biden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Carpetweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds. WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

<sup>1</sup> The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .8 pound per 250 square feet.

SPREADER SETTINGS CNE PASS APPLICATION:

<u>Make</u>

		•
Central Calibrated	5	
Cyclone Broadcast	3-3/4	****
Jackson	5	FECE
Sears Broadcast	5°, 5°,	5
Sears	5	1997 - A.
Scotts Calibrated	<u>i</u>	1 4 t
	``	

SETTING

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Without spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas ith oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing  $\star$  (No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. \*Eased in Inert Fertilizer claimed.

### STORAGE AND DISPOSAL

**STORAGE:** Store in criginal container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed, or other pesticides.

**PRODUCT DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (ERP 061594) (PR938 042194)

1

# **OPTIONAL SPREADER INSTRUCTIONS:**

(Illustration)	For Drop Spreaders: Firs	t apply two	strips at end	s. Then apply
	product along the length	of the lawn	, overlapping	very slightly and
	shutting off spreader whe	n turning.		

For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly. (Illustration)

# OPTIONAL MARKETING STATEMENTS:

Contains Dissolve® A Premium Broadleaf Herbicide

Environmental Benefits (Grasseyeling Logo)	Besides increasing property Values, did you know benefits to your home and environment: Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.	<ul> <li>A 50' x 50' lawn provides the following</li> <li>A 50' x 50' lawn produces enough oxygen for a family of 4 every day,</li> <li>Reduces Glare &amp; Reduces Noise</li> <li>Traps Dust, Smoke Particles &amp; other Follutants</li> <li>Purifies Water &amp; Modifies Temperature</li> <li>Improves Soil Condition &amp; Controls Erosion</li> </ul>
The following note will	not be part of the actual label	copy.
<pre>commercial fe guarantee are guarantee are List Editor's note List Use Commercial fe guarantee are List Commercial fe guarantee are List Commercial fe Guarantee are List Commercial fe Commercial fe Guarantee are List Commercial fe Commercial fe Guarantee are List Commercial fe Comme</pre>	fertilizer) Analysis and Spreader ertilizer manufacturers. Chlorine e optional except where mandated b gistrant Warranty Statements prev n their own registered products a es still require the term INERT	e declaration and fertilizer by State fertilizer regulation. iously accepted by U.S.EPA for re <u>not</u> included in this label.

10/10