

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number: 228-358

Date of Issuance: NOV 17 1998

Term of Issuance: Conditional

Name of Pesticide Product: Riverdale Esteron(R) 99(R) C

NOTICE OF PESTICIDE:
[X] Registration
Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Co.
425 West 194th St.
Glenwood, IL 60425

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce.

*On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes:
a. Revise the EPA Registration Number to read, "EPA Reg. No. 278-358".
3. Under the "Statement of Practical Treatment", revise the dermal statement to read as follows: "Wash with plenty of soap and water. Get medical attention."

Signature of Approving Official:

Date:

NOV 17 1998

Handwritten notes: 7505C, 11/17/98

4. Under the "Crop Use" section of the labeling, page 7, in the "Caution" area for "Woody weeds in Low-Brush Blueberry Fields in Maine", second sentence, replace the word "of" with the word "or" so that the sentence reads "...treatment year or until a..."

5. Submit two copies of the revised final printed label for the record.

A stamped copy of the label is enclosed for your records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller
Product Manger (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

RIVERDALE

ESTERON® 99® C

A SELECTIVE HERBICIDE

FOR THE CONTROL OF MANY BROADLEAF WEEDS, HERBACEOUS PERENNIALS,
AND WOODY PLANTS SUSCEPTIBLE TO 2,4-D IN GRASS PASTURES,
CERTAIN CROPS AND NON-CROP AREAS.

ACTIVE INGREDIENT:

2-Ethylhexyl Ester of 2,4-Dichlorophenoxyacetic Acid* 65.9%

INERT INGREDIENTS: 34.1%

TOTAL 100.0%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid 43.7%, 3.8lbs./gal.

Esteron® & 99® are trademarks of Dow AgroSciences LLC.

KEEP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

(If you do not understand the label, find someone
to explain it to you in detail.)

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-GLI

NET CONTENTS

GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY


RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED
with COMMENTS
in EPA Letter Dated
NOV 17 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-358

NOTE: Spanish Language is optional.



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION - CAUCION**

Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Avoid breathing vapors or mists. Avoid contact with skin, eyes, or clothing. For containers of over 1 gallon, but less than 5 gallons: Mixers and loaders who do not use a mechanical system (such as probe and pump or spigot) to transfer the contents of this container must wear coveralls or chemical-resistant apron in addition to other required PPE.

Personal Protective Equipment (PPE)- Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton, shoes plus socks and protective eyewear. Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selections chart.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements - For containers of 5 gallons or more: Do not open pour product from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Immediately flush skin with plenty of water. Get medical attention if irritation persists.

IF SWALLOWED: Do NOT induce vomiting. Call a physician. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: May cause chemical pneumonitis if aspirated. If lavage is performed, suggest endotracheal and/or esophagosopic control.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not cut or weld container.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils and Viton \geq 14 mils; shoes plus socks, and protective eyewear.

GENERAL INFORMATION

Esteron® 99® C Herbicide is intended for the control of many broadleaf weeds, herbaceous perennials and woody plants susceptible to 2,4-D in grass pastures, certain crops and non-crop areas.

Apply Esteron 99 C Herbicide as water or oil spray during warm weather when weeds or brush are actively growing. Application under drought conditions often will give poor results. Use low spray pressure to minimize drift. On cropland and along roadsides, do not exceed 20 psi pressure. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons per acre by ground equipment and 3 to 5 gallons by aircraft. Higher gallonage may be used if desired to improve spray coverage. Generally, the lower dosages recommended on this label will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. For crop uses, do not mix

with oil or other adjuvants unless specifically recommended on this label. Deep-rooted perennial weeds such as Canada thistle and Field Bindweed and many woody plants usually require repeated applications for maximum control. Do not apply Esteron 99 C where spray drift may contact nearby susceptible crops or other desirable plants or may contaminate water for irrigation or domestic use. Read and follow all Use Precautions given on this label.

Note: If there are uncertainties concerning special local use situations or specific crop variety tolerances to 2,4-D, consult your State Agricultural Experiment Station or local Extension Service Weed Specialists for advice.

GENERAL USE PRECAUTIONS

Chemigation: Do not apply this product through any type of irrigation system.

Avoid Contact With 2,4-D Susceptible Crops and Other Desirable Broadleaf Plants: Do not apply directly to or otherwise permit even minute amounts to contact cotton, grapes, tobacco, fruit trees, vegetables, flowers, ornamentals, or other desirable plants susceptible to 2,4-D. Do not use in or near a greenhouse.

DO NOT APPLY IN THE VICINITY OF COTTON, GRAPES, TOBACCO, TOMATOES, OR OTHER DESIRABLE 2,4-SUSCEPTIBLE CROPS OR PLANTS. DO NOT SPRAY WHEN WIND IS BLOWING TOWARDS SUSCEPTIBLE CROPS OR ORNAMENTAL PLANTS.

Avoid Spray Drift: Applications should be made only when there is no hazard from spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during both growing and dormant periods. Use coarse sprays to minimize drift since, under adverse weather conditions, fine spray droplets may drift a mile or more. A spray thickening agent such as Nalco-Trol, may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Equipment: With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using no more than 20 pounds spraying pressure at large droplet producing nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray. Determine air movement and directions before foliar application. Use a smoke generator or other means at or near the application site for the detection of air movement, air stability or temperature inversions. Such a condition exists when there is little or no wind and air temperature is lower near the ground than at higher levels. Use appropriate drift control measures or avoid application when smoke is moving toward nearby desirable susceptible plants or sensitive areas.

Aerial Application: With aircraft, drift can be lessened by applying a coarse spray; by using no more than 20 pounds spray pressure at the nozzles; by using straight stream nozzles directed straight back; by using a spray boom no longer than 3/4 the wing or rotor span of the aircraft; and by spraying only when wind velocity is less than 6 mph.

Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination or plant growth. Violent wind storms may move soil particles. If 2,4-D is on soil particles and they are blown onto the susceptible plants, visible symptoms may appear. Serious injury is unlikely. The hazard of movement of 2,4-D on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.

At high temperatures, vapors from this product may injure susceptible plants growing nearby. To avoid injury to desirable plants, do not handle or apply other agricultural chemicals with the same equipment used for Esteron 99 C unless appropriately cleaned first. Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station or Extension Service weed specialists for cleaning methods which are in compliance with local regulations and for advice in selecting treatments from this label to best fit local conditions. Be sure that use of this product conforms to all applicable regulations.

Weeds Controlled

Esteron 99 C is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. Species controlled

include the following, plus many others: Beggarticks, Bitterweed, Texas blueweed, Broomweed, Buckbrush, Wild buckwheat, Burdock, Burhead, Carpetweed, Catnip, Chamise, Chicory, Cocklebur, Coffeeweed, Cornflower, Coyotebrush, Croton, Dandelion, Docks, Dogfennel, Elderberry, Galinsoga, Wild garlic, Goatsbeard, Wild hemp, Jewelweed, Jimsonweed, Ladysthumb, Lambsquarter, Bigbend loco, Venice mallow, Manzanita, Marshelder, Milkvetch, Annual Morningglory, Nettles, Wild onion, Pennycress (Fanweed), Field pepperweed, Pigweed†, Plantains, Poorjoe, Rabbitbrush, Wild radish, Ragweed, Tansy ragwort, Wild rape, Redstem, Coastal sage, Big sagebrush, Sand sagebrush, Salsify, Sand shinnery oak, Shepherdspurse, Sicklepod, Annual smartweed, Bitter sneezeweed, Annual Sowthistle, Spanishneedles, Sumac, Sunflower, Sweetclover, Bull thistle, Musk thistle, Russian thistle, Tumbleweed, Velvetleaf, Vervains, Vetch, Water plantain, Wild mustard, Willow, Witchweed, Wormweed, Yellow rocket, Yellow starthistle.

†The control of "hybrid" Pigweeds appears to be less satisfactory from 2,4-D products than formerly experienced on "non-hybrid" varieties. Since 2,4-D herbicides are not as effective on the "hybrid" Pigweeds, it is necessary to apply higher rates of 2,4-D for control, especially later in the growing season. Higher rates injure some crops, so less than satisfactory Pigweed control may be experienced by the highest tolerated crop dosage.

Therefore, Dow AgroSciences no longer includes Pigweed among the species covered by the performance guarantee statements on the labels for Esteron 99 C. At this time, this disclaimer applies only to the High Plains of Texas and western Oklahoma, including the Panhandles. All other guarantees on these product labels are unchanged by this disclaimer.

TO PREPARE THE SPRAY: (1) Fill the spray tank about half full with water, then add the required amount of Esteron 99 C with agitation, and finally the rest of the water. Note: Esteron 99 C in water forms an emulsion which tends to separate unless the mixture is kept agitated. (2) If oil is added, first mix Esteron 99 C and the oil and then add this mixture to the water. However, with adequate agitation, the oil can be added after Esteron 99 C is mixed with water. (3) If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil-herbicide mixture to avoid formation of an invert emulsion.

SELECTIVE WEEDING IN CROPS

USE IN LIQUID NITROGEN FERTILIZER

This product may be combined with liquid nitrogen fertilizer suitable for foliar application to accomplish broadleaf weed control and fertilization of corn, small grains or pastures in a single operation. Use Esteron 99 C in accordance with recommendations for these crops provided in this label. Use liquid fertilizer at rates recommended by the supplier or Extension Service Specialist. Test for mixing compatibility by mixing spray ingredients in correct proportions in a clean glass jar before mixing in spray tank. A compatibility aid such as Unite or Compex may be needed in some situations. Compatibility is best with liquid fertilizer solutions containing only nitrogen. Mixing with N-P-K solutions may not be satisfactory, even with the addition of a compatibility aid. Pre-mixing Esteron 99 C with 1 to 4 parts water may help in situations when mixing difficulty occurs.

Fill the tank about half full with the liquid fertilizer, then add the required amount of Esteron 99 C with agitation. Maintain agitation and complete filling the tank with liquid fertilizer. Apply immediately and continue agitation in spray tank during application. Do not store the spray mixture. Application during very cold weather (near freezing) is not advisable.

APPROVED USES

Crop and Forestry Uses

Agricultural Use Requirements for Crops and Forestry: For the following crop and forestry uses, follow PPE and Reentry instructions in the "Agricultural Use Requirements" section of this label.

Weed Control in Small Grains Not Underseeded With A Legume

Note: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment.

Spring and Winter Wheat, Barley, and Rye

Apply ½ to 1 pint per acre. Spray when grain is in full tiller-stage (usually 4 to 8 inches tall) but before the boot stage and boot to dough stage. For improved control of difficult weeds including wild garlic and wild onion or under dry or cool conditions, apply up to 2 pints per acre. Wild garlic and wild onion may not be killed but dockage should be reduced. Do not use higher rates unless possible crop injury will be acceptable. Consult State Agricultural Experiment Station or Extension Service Weed Specialists for recommendations or suggestions to fit local conditions.

Spring Seeded Oats

Apply ½ pint per acre at the full tiller stage but before the early boot stage. Oats are less tolerant to 2,4-D than Wheat or Barley and are more likely to suffer some injury.

Fall Seeded Oats (Southern) Grown for Grain

Apply ¾ to 1 ¼ pints per acre after full tillering but before the early boot stage. Some difficult weeds may require higher rates for maximum control but crop injury may result. Do not spray during or immediately following cold weather.

Preharvest Treatment: Apply 1 to 2 pints per acre when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth.

Note: Do not feed treated straw to livestock.

Weed Control in Corn

(Use one of the following programs)

Preemergence: Apply 1 to 2 quarts per acre to soil anytime after planting but before corn emerges. Only emerged broadleaf weeds are likely to be controlled. Do not apply more than 1 quart per acre unless the increased risk of crop injury can be tolerated. Do not use on light sandy soil.

Emergence: Apply 1 pint per acre just as corn plants are breaking ground.

Postemergence: After emergence of corn use ½ pint per acre. Application of ¾ to 1 pint per acre may be needed for maximum control of some weeds but such rates are more likely to injure the corn. If corn is over 8 inches tall, use drop nozzles to keep the spray off the corn foliage as much as possible. Do not apply from the tasseling to dough stage. Do not use with oil, atrazine or other adjuvants. Crop injury is more likely to occur if corn is growing rapidly under high temperature and high soil moisture conditions. To reduce breakage of stalks from temporary brittleness caused by 2,4-D, delay cultivation for 8 to 10 days after treatment. Do not forage or feed corn fodder for 7 days following application.

Note: Hybrids vary in response to 2,4-D and some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Contact seed company or your Agricultural Experiment Station or Extension Service weed specialists for this information.

Preharvest Corn Treatment: After the hard dough or denting stage, apply 1 to 2 pints per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as Bindweed, Cocklebur, Dogbane, Jimsonweed, Ragweed, Sunflower, Velvetleaf and vines that interfere with harvesting. Do not forage or feed corn fodder for 7 days following application.

Weed Control in Sorghum (Milo)

Apply ½ pint per acre when sorghum is 5 to 15 inches tall. A higher rate of ¾ to 1 pint per acre may be needed to control some weeds but the chance for crop injury is likewise increased. Do not use with oil. Do not treat before the sorghum is 5 inches tall nor during the boot, flowering or early dough stages. If sorghum is taller than 8 inches, use drop nozzles to keep the spray off the foliage as much as possible. Temporary crop injury may occur under conditions of high soil moisture and high air temperatures. Varieties vary in tolerance to 2,4-D and some hybrids are quite sensitive. Spray only varieties known to be tolerant to 2,4-D. Contact seed company or your Agricultural Experiment Station or Extension Service weed specialists for this information.

Weed Control in Fallow Land And Crop Stubble

Apply 1 to 2 pints per acre for control of annual and biennial broadleaf weeds. Use the higher rate on older drought stressed plants or hard to kill species. Apply 2 to 6 pints per acre for control of perennial broadleaf weeds. Spray weeds in the bud to bloom stage

or in good vegetative growth. Do not plant treated fallow land for three months or until chemical has disappeared from soil.

Control of Wild Garlic and Wild Onion: Following harvest of small grains, Soybeans, Corn or grain Sorghum, Wild garlic and Wild onion often produce new fall growth. This new growth may be treated with Esteron 99 C at a rate of 2 to 3 quarts per acre. This treatment is useful as part of an overall control program.

Control of Woody Weeds in Low-Brush Blueberry Fields in Maine

How To Use: Mount a drum 8 to 10 feet long or some other suitable length, and 1 ½ to 2 feet in diameter on an axle such as an old hay rake frame. Cover the drum with water-absorbent yet tough cloth which will resist rapid wear and tear. Draw the cloth-covered drum across the blueberry field and at the same time spray evenly onto the full length of the top of the cloth covered drum a spray mixture made by diluting 1 quart of Esteron 99 C in 50 gallons of water per acre. Have the drum mounted so that as it revolves on its axis, it is high enough to miss most of the low brush blueberry stems, yet low enough to forcibly brush the spray-saturated cloth-covered drum against the higher woody weeds, principally Sweet fern, Wild cherry and Poplar. Keep the cloth wet enough to provide top coverage of the weeds, yet not so wet as to allow runoff of the liquid which could cause injury to the Blueberry plants.

When to Use: Apply during June and July when weed tops have emerged sufficiently above the blueberry stems to allow treatment of the weeds and not the Blueberry plants. Apply only during the year before the first burn. To use this method of weed control, two-year burns should be extended to three years.

Caution: Do not allow the spray being applied to the cloth-covered drum to be directed onto the Blueberries. Do not harvest-rake field during the herbicide treatment year or until a two-year interval thereafter.

Forest Site Preparation

For control of susceptible broadleaf weeds and brush on sites to be planted in forests, use 1.5 to 4 quarts per acre of Esteron 99 C in sufficient spray volume for good plant coverage, usually 6 to 25 gallons. Applications can be made by air or ground (hand gun, boom, or powered knapsack sprayer). Two to eight quarts of diesel oil per acre or a suitable surfactant or penetrant may be added to improve brush control.

Forest Conifer Release

For applications in late winter or spring to control susceptible deciduous brush species, such as Alder, Willow, Poplars, Cascara, Cherry, service Cherry and Vine maple during early growth and before Conifer budbreak, use Esteron 99 C at rates up to 3 quarts per acre in diesel or stove oil by air or ground in sufficient spray volume for good plant coverage, usually 6 to 25 gallons. Do not use in plantations where Pine or Larch are among the desired species.

For treatment before Conifer budbreak to control susceptible Evergreen brush species, such as Tanoak, Madrone, Chinquapin, ceanothus spp. and Manzanita or deciduous brush after leafout or broadleaf weeds, use Esteron 99 C at rates up to 3 quarts per acre alone or with 0.5 to 2.0 gallons per acre of diesel or similar oil or suggested rates of suitable surfactants or penetrants. After Conifer budbreak, Esteron 99 C without oil, surfactant or penetrant can be used at rates up to 2 quarts per acre but may cause injury or suppression of the Conifer growth. Use sufficient volume of spray for good coverage of brush, usually 6 to 25 gallons. Some species of Pine may be seriously injured by treatment of these growth stages.

After Conifer species such as White pine, Ponderosa pine, Jack pine, Red pine, Black spruce, White spruce, Red spruce, and Balsam fir cease growth and harden off and brush is actively growing in late summer, 1.5 to 3.0 quarts of Esteron 99 C per acre in enough water to obtain good plant coverage may be applied by air or ground to control certain competing hardwood species such as Alder, Aspen, Birch, Hazel, and Willow. Since this treatment may cause occasional Conifer injury, do not use if such injury cannot be tolerated.

Directed Sprays In Conifer Plantations (including pine)

Apply Esteron 99 C at any time brush or broadleaf weeds are susceptible by directing spray around the Conifers to avoid contact of needles with injurious amounts of spray. Rates of Esteron 99 C are not to exceed 4 quarts per acre in oil, oil-water, or water carrier at 10 to 100 gallons per acre.

Rangeland, Pasture, and Non-Crop Uses

Use Requirements for Rangeland, Pasture and Non-cropland Areas: No worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is applied to rangeland, pasture or non-cropland areas.

Weed And Brush Control in Rangeland, Conservation Reserve, and Grass Pastures

Note: Do not use on Bent grass, Alfalfa, Clover, or other Legumes.
Do not use on newly seeded areas until grass is well established.
Do not use from early boot to milk stage where grass seed production is desired.

For Conservation Reserve Land, follow all applicable state and Federal regulations. Follow the most severe grazing restrictions imposed either by the pesticide label or by the USDA Acreage Conservation Reserve Program, whichever is longest.

Grazing Restrictions: Do not graze lactating dairy animals on treated areas within 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days. Withdraw meat animals from treated forage at least 3 days before slaughter.

Control of Bitterweed, Broomweed, Croton, Docks, Kochia, Marshelder, Muskthistle, and other Broadleaf Weeds

Use 2 quarts of Esteron 99 C per acre in the amount of water needed for uniform application. If the weeds are young and growing actively, 1 quart per acre will provide control of some species. Deep-rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

Control of Wild Garlic and Wild Onion

Apply 2 to 3 quarts per acre, making three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring. For rangeland and pasture, the maximum rate is 2 quarts per acre.

Control of Weed Control in Newly Sprigged Coastal Bermudagrass

Apply 1 to 2 quarts per acre preemergence and/or postemergence.

Control of Sand Shinnery Oak and Sand Sagebrush

On the Oak, use 1 quart in 5 gallons of oil or in 4 gallons of water plus 1 gallon of oil per acre. Apply by aircraft between May 15 and June 15. On the Sagebrush, use 1 quart in 3 gallons of oil per acre and apply by aircraft when foliage is fully expanded and the brush is actively growing.

Control of Big Sagebrush and Rabbitbrush

Use 2 to 3 quarts per acre in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. For Rabbitbrush, the 3 quart rate is usually required. Brush should be leafed out and growing actively when treated. Retreatment may be needed. For rangeland and pasture, the maximum application rate is 2 quarts per acre.

Control of Chamise, Manzanita, Buckbrush, Coastal Sage, Coyotebrush, and Certain Other Chaparral Species

Use 2 quarts per acre in 5 to 10 gallons of water. One gallon of fuel may be included in the spray mixture for added effectiveness. Make applications by aircraft or ground equipment to obtain uniform spray coverage. For effective control, the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed. For rangeland and pasture, the maximum application rate is 2 quarts per acre.

Woody Plant Control in Non-Crop Areas

To control species susceptible to 2,4-D in rights-of-way, fencerows, roadsides, and along drainage ditch banks, spray brush up to 5 to 8 feet tall after spring foliage is well developed, using 3 to 4 quarts of Esteron 99 C in 100 gallons of water and wetting all parts of the brush foliage, stems and bark. This may require up to 400 gallons of spray per acre for adequate coverage of solid stand of brush. Make application in such a way as to prevent drift of the spray off the area being treated. Spraying can be effective at any time up to 3 weeks before frost as long as the soil moisture is sufficient for active growth of the brush. Control will be less effective in midsummer during hot, dry weather when soil moisture is deficient and plants are not actively growing. Oil or wetting agent may be added to the spray if needed for increased effectiveness.

Weed Control in Non-Crop Areas Airfields, Roadsides, Vacant Lots

Apply 1 to 3 quarts of Esteron 99 C per acre in the amount of water needed for uniform application. Usually 2 quarts per acre provides good weed control under average

conditions. Treat when weeds are young and growing well. Do not use on Dichondra or other broadleaf herbaceous ground covers. Do not use on creeping grasses such as Bent and St. Augustine except for spot treating, nor on newly seeded turf until grass is well established. Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed so do not treat where Legumes are desired. Deep-rooted perennial weeds may require repeated treatments in the same season or in subsequent years.

Tule (Bulrush) And Other Rushes

Mix 2 quarts of Esteron 99 C and 1 gallon of diesel oil or kerosene, then add this mixture to 100 gallons of water. Spray to wet all foliage (400-800 gallons per acre). Addition of a wetting agent may be advisable. Apply in the spring during flower head emergence. Respray if needed when regrowth is 3 to 5 feet tall.

Spot Treatment

To control broadleaf weeds in small non-cropland areas with a hand sprayer, use 1/4 pint of Esteron 99 C in 3 gallons of water and spray to thoroughly wet all weed foliage. Keep spray mixture agitated to prevent separation.

Turf Uses

Use Requirements for Turf Including Grass Seed Crops

Restricted Entry Interval: When used on grass seed crops, follow PPE and reentry instructions in the "Agricultural Use Requirements" section of this label. For use on other turf areas, do not allow people (other than applicator) or pets on treatment during application. Do not enter into treated areas until sprays have dried.


Restrictions on Retreatment: Do not apply more than 2 broadcast applications per year per treatment site.

Weed Control in Grass Seed Crops

Use 1 to 1 1/2 pints per acre in the amount of water required for uniform application by air or ground equipment. Apply to established stands in spring from the tiller to early boot stage. Do not spray in boot stage. New spring seedlings may be treated with the lower rate after the grasses have at least five leaves. Perennial weed regrowth may be treated in the fall.

Weed Control In Turf Areas Such As Lawn, Golf Courses, Cemeteries, and Parks

Apply 1 to 2 quarts of Weed Killer 4D per acre in the amount of water needed for uniform application. Usually 2 quarts per acre provides good weed control under average conditions. Treat when weeds are young and growing well. Do not use on golf greens nor on Dichondra or other broadleaf herbaceous ground covers. Do not use on creeping grasses such as Bent and St. Augustine except for spot treating, nor on newly seeded turf until grass is well established. Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed so do not treat areas where Legumes are desired. Deep-rooted perennial weeds may require repeated treatments in the same season or in subsequent years.



STORAGE AND DISPOSAL

STORAGE: Keep container tightly closed when not in use. This product can be stored in an unheated building. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL (Metal): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL (Plastic): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL: Consult federal state, or local disposal authorities for approved alternate procedures. Be sure that use of this product conforms to all application regulations.

WARRANTY

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith. (ERP 102393) (PR933&11 042194)

