UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

THITED STATE	_
thu _n	AGE NO
ENTAL SECTION	•

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

_ Reregistration

x Registration

NOTICE OF PESTICIDE:

EPA Reg. Number:

PEBOF Issuance:

228-371

Term of Issuance:

Conditional

Name of Pesticide Product:

Riverdale MTCA Premium Selective Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Co. 425 West 194th St.

Glenwood, IL 60425-1584

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the rasis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA sec. 3(c)(5) or 4 of 3(g) when the Agency requires all registrants of similar products to submit such data.
 - 2. Make the following label changes:
- a. Revise the EPA Registration Number to read, "EPA Reg. No. 228-371."
- 3. Under the "Storage and Disposal" section of the labeling, in the "Storage" statement, relocate the last sentence, that reads, "Do not contaminate...", to appear immediately under the heading "Storage and Disposal".

Signature of Approving Official:

Date

FEB | 2000

EPA Form 8570-5

1505t Myryan 1/31/00 page 2 EPA Reg. No.228-371

- 4. Submit two copies of the revised final printed label for the record.
- 5. Submit a corrosion characteristics study (GRN 830-6320) within one (1) year of the date of this notice.

A stamped copy of the label is enclosed for your records. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller Product Manger (23) Herbicide Branch Registration Division (7505C)

Enclosure

RIVERDALE

MTCA PREMIUM SELECTIVE HERBICIDE

FOR THE CONTROL OF AMNUAL AND PERENNIAL BROADLEAF WEEDS IN ORNAMENTAL TURF INCLUDING SOD FARMS

CONTAINS MCPA, TRICLOPYR, AND CLOPYRALID

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS: Dimethylamine Salt of 2-Methyl-4-Chlorophene Triethylamine Salt of 3,5,5-Trichloro-2-Pyr:	idiny	ylox	yace	etio	c A	cid	* *		-			3.8%
3,6-Dichloro-2-Pyridinecarboxylic Acid***												1.3%
INERT INGREDIENTS:												
							TOTA	L		-		100.0%
By Isomer Specific AOAC Method, Equivalent to: *2-Methyl-4-Chlorophenoxyacetic Acid **3,6-Dichloro-2-Pyridinecarboxylic Acid ***3,6-Dichloro-o-Anisic Acid								2.7%,	0	.27	lbs	s./gal.

For Use By Professional Turf Maintenance Personnel, Landscaping or Commercial Applicators Only.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-

The Telescope of the State of t

NRŢ CONTENTS

GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

with COMMENTS
In EPA Letter Dated
FEB | 2000

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 228-371

(Inside Booklet)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed, or absorbed through skin. Do not get in eyes or clothing. Wash thoroughly with soap and water after handling.

NON-WPS TURF USE: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general only agricultural plant uses are covered by the WPS--must wear eye protection (goggles, face shield, or safety glasses) when using this product. It is recommended that safety glasses include front, brow, and temple protection.

WPS SOD FARM USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170)—in general, agricultural plant uses are covered—must wear: Long—sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment(PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS: If the mechanical system is used in a manner that meets the requirements listed in the worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands, face, and arms with soap and water before eating, smoking, drinking, or using the toilet. Wash the outside of gloves before removing.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES:

IF ON SKIN: IF SWALLOWED:

, , '

The state of the s

Hold eyelids open, and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

Wash with plenty of soap and water. Get medical attention.
Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg white, gelatine solution, or if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use .,, of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this product will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

"se this product only in accordance with its labeling and with the Worker Protection bandard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI of 48 hours.

PPE required for early entry to treated areas that is permitted under the worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application Do not enter treatment areas until spray has dried.

MTCA PREMIUM SELECTIVE HERBICIDE is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf such as Lawns, Parks, Cemeteries, Roadsides, Golf Courses (Fairways, Aprons, and Roughs), and similar non-crop areas; also for use on sod farms. Do not use on Golf Greens or Tees.

MTCA may be applied to the following established turfgrass species:

Established Cool Season Turfgrasses

Common Name Scientific Name bentgrass Agrostis species bluegrass, Kentucky Poa pratensis

fescue, chewing Festuca rubravar. commutata

fescue, creeping red Festuca rubra fescue, sheeps fescue, tall Festuca ovina

Festuca arundinaceae

ryegrass, perennial . Lolium perenne

Established Warm Season Turfgrasses†

Common Name Scientific Name bahiagrass Paspalum notatum bermudagrass Cynodon dactylon buffalograss Buchloe dactyloides centipedegrass Eremochloa cphiuroides zoysiagrass Zoysia japonica zoysiagrass Zoysia tenuifolia fescue, tall (growing in Festuca arundinacea warm season areas)

†Do not treat warm season turfgrass with MTCA when mowing height is less than 1/2 inch.

Note: MTCA may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, wellestablished turf that has a well-anchored root system. Do not apply this product to grass species not listed on this label unless potential turf injury can be tolerated.

USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact: with cotton, citrus, grapes, tobacco, vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solutions near desirable plants. Ayoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind velocity is low, keeping spray boom as low as possible, using nozzles that deliver larger spray droplets and minimize the production of fine spray particles, and by using higher spray volumes and lower pressures. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants

near enough to be injured. Do not apply to exposed roots of shallow rooted trees and shrubs. Do not use grass clippings from turf, treated with this product for mulch or compost. Maximum kill of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Do not apply more than 4 pints per acre per application. Mow newly seeded turf at least twice before treatment with MTCA.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is usei; (2) turf should not be moved 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product.

WEEDS CONTROLLED BY MTCA PREMIUM SELECTIVE HERBICIDE

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Acacia, Alder, Alfalfa, Arrowhead, Artichoke, Aster, Austrian fieldcress, Bachelor button, Beggarweed, Beggartick, Biden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Buckwheat, Buffalobur, Eull thistle, Burcucumber, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamomile, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Cornflower, Creeping jenny, Croton, Curly indigo, Daisy, Dandelion, Dock, Dogbane, Dog fennel, Dragonhead Mint, Elderberry, Fanweed, Fat hen, Flea bane, Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goatweed, Caldenrod, Ground ivy, Groundsel, Gumweed, Halbertleaved saltbush, Hawksbeard, Hawkweed, Alall, Hemp, Henbit, Hoary cress, Honeysuckle, Hop Clover, Horsetail, Horseweed, Indiana Mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knapweed, Knotweed, Kudzu, Ladysthumb, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Meadow salsify, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nightshade, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Queen Annes lace, Ragweed, Red clover, Red root, Red stem, Red sorrel, Rush, Russian thistle, St. Johnswort, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdck, Speedwell, Spurge, Spurwood, Starthistle, Stinging nettle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansy Ragwort, Tarweed, Teasel, Thistle, Thornapple, Toadflax, Tree-of-heaven Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper, Virginia pepperweed, Whitebrush, White clover, Whitetop, Wild aster, Wild carrot, Wild geranium, Wild gosseberry, Wild j

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second 'roadcast treatment in 3 to 4 weeks. In the event more than two broadcast applications repear are necessary, use as a spot treatment. To spot treat, make up solution as listed below.

TURF APPLICATION DIRECTIONS

Apply 3 to \pm pints of MTCA per acre (1.1 to 1.5 ozs./1,000 sq.ft.) to control above listed broadleaf weeds.

The maximum use rate for most warm season grasses is 3 pints per acre.

Make application using equipment that will ensure uniform coverage (see specifized application directions below). Foliar sprays should be applied during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. When hard-to-control species are prevalent, when applications, are made in late Summer, when weeds are mature, or during drought conditions, use the higher rates.

On closely moved golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.73 fluid ounces in 0.55 to 5.5 gallons of water per 1000 square feet). During conditions which promote turf stress use lower rates, 1 to 1-1/2 pints in 20 to 240 gallons of water, per acre (0.36 ro 0.55 fluid dunces in 0.3 to 5.5 gallons per 1000 square feet). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course greens or tees.

Note: Broadleaf weeds germinate at different times of the year. Only emerged weeds

present at the time of application will be controlled by MTCA.

Standard Volume Broadcast Applications: Apply 3 to 4 pints of MTCA in enough water to deliver 20 to 200 gallons of total spray mix per acre (0.5 to 5 gallons spray per 1,000 square feet). Higher application volumes may be used when this product is tank mixed with liquid fertilizers.

Low Volume Broadcast Applications: Apply 3 to 4 pints of MTCA in enough water to deliver from 5 to 40 gallons of total spray mixture per acre. Use low pressures and application equipment capable of delivering a uniform droplet size that can wet the weed leaf surface.

Targeted Zone Applications or Spot Treatment Using Portable Sprayers: Mix 0.5 to 1.0 ounces of MTCA in enough water to make one gallon of spray and apply at any time broadleaf weeds are actively growing. Target plants should be sprayed to wet but not beyond point of runoff.

Herbi™ Controlled Droplet Applicator: Add 2-1/4 to 3 pints of MTCA to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution.

While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of MTCA must be applied when grass is stressed from heat, drought, etc.

 $\mathsf{Herbi^{TM}}$ is a trademark of North American Micron.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use MTCA at the rate of 3 to 4 pints per acre (1.1 to 1.5 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Note: Follow CDA equipment spray instructions.

Bertrandistante washington while acceptance of

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near open containers of herbicides and other pesticides, fertilizer and seed. Store at temperatures above 23°F or agitate before using. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR NON-REFILLABLE CONTAINERS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by barring. If burned, stay out of smoke.

CONTAINER DESPOSAL FOR REFILLABLE CONTAINERS: Close all openings which have been opened during use and replace all caps. Contact Riverdale Chemical's Customer Service Department, at 1-708/754-3330, to arrange for return of the empty refillable container.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label.

(PR938&11 042194)

`\

a built