

228-372

2-28-2000

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

228-372

Date of Issuance:

FEB 28 2000

NOTICE OF PESTICIDE:

- Registration
- Reregistration

Term of Issuance:

Conditional

Name of Pesticide Product:

Riverdale Eclipse
Selective Herbicide

Under FIFRA, as amended:

Name and Address of Registrant (include ZIP Code):

Riverdale Chemical Company
425 West 194th Street
Glenwood, IL 60425-1584

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA section 3(c)(5) or 4 or 3(g) when the Agency requires all registrants of similar products to submit such data.

2. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read "EPA Reg. No. 228-372".

Signature of Approving Official:

Date:

FEB 28 2000

M. C. Ke
2-28-00

AT YOUR DISCRETION: You may revise the First Aid statements in accordance with recent Agency guidance. The new statements (below) were developed as part of the Consumer Labeling Initiative in close cooperation with poison control center personnel and other medical experts. While it is not mandatory that you revise your label at this time, you are strongly encouraged to substitute the revised statements for those statements currently on the label:

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

3. Submit one copy of the revised final printed label of the record before you release the product for shipment.

Page 3
EPA Reg. No. 228-372

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

4/8

RIVERDALE

ECLIPSE™ SELECTIVE HERBICIDE

A PREMIUM THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE CONTAINING THE AMINES OF MCPA, CLOPYRALID AND DICHLORPROP-p. FOR USE ON GOLF COURSES, PARKS, ORNAMENTAL TURF LAWNS AND SIMILAR NON-CROP AREAS. CONTROLS DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	47.77%
Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid**	2.53%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†	9.54%
INERT INGREDIENTS:	40.16%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	39.00%	3.74 lbs./gal.
**3,6-Dichloro-2-Pyridinecarboxylic Acid	1.92%	0.18 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	8.00%	0.76 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p.

ECLIPSE™ - Is A Trademark of Riverdale Chemical Company

For Use By Professional Turf Maintenance Personnel,
Landscaping or Commercial Applicators Only.
Not For Sale to or Use By Homeowners.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-GTE

NET CONTENTS

GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

ACCEPTED
with COMMENTS
In EPA Letter Dated
FEB 28 2000

NOTE: Spanish language is optional

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-372

5/8

(Inside Booklet)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER - PELIGRO**

Corrosive, causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, eggwhites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this product will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Eclipse™ SELECTIVE HERBICIDE is for use on Ornamental Turf such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees* and Roughs), and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.*Excluding Bentgrass Tees.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Carpetgrass, Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline

of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

The suitable use of Eclipse on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED BY ECLIPSE™ SELECTIVE HERBICIDE

Alfalfa, Amaranth, Aster, Bachelor button, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campion, Bracken fern, Broomweed, Uckhorn, Buckwheat, Buffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground ivy, Groundsel, Halfshrub sundrop, Hawksbeard, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle, Henbit, Horsenettle, Horsetweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Locoweed, Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red sorrel, Red clover, Redvine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse, sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny, Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sumpweed, Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Toadflax, Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild radish, Wild onion, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket, and many other broadleaf weeds.

ORNAMENTAL LAWNS AND TURF

Apply Eclipse Selective Herbicide at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

Herbi Controlled Droplet Applicator: Add 1-1/2 to 2-1/4 pints of Eclipse Selective Herbicide to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution.

While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Reduce application rates to stressed turf unless temporary injury can be tolerated.

Herbi™ is a trademark of North American Micron.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use Eclipse Selective Herbicide at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) Avoid double coverage resulting from overlap unless temporary turf injury can be tolerated. (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

7/8

COMPATIBILITY

Eclipse Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of Eclipse Selective Herbicide.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding the Eclipse Selective Herbicide.)
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds or other pesticides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings and replace all caps. Contact Riverdale Chemical's Customer Service Department at 1-708/754-3330, to arrange for return of the empty refillable container.

8/8

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label. (PR938&11 042194)

