



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505C)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

228-375

Date of Issuance:

JUL - 1 2009

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Consair Selective Herbicide

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc
 150 Harvester Drive, Suite 200
 Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above provided you make the following changes to your label.

1. On page 1, revise the first paragraph to read "For selective control of grassy and broadleaf weeds in turfgrass, **on lawns, commercial areas, and industrial sites**, and sodfarms.
2. Base on the acute toxicology review, the Hazards to Humans and Domestic Animals must be revised to read:

"Caution
 Harmful if Absorbed through skin. Avoid contact with skin, eyes or clothing."

Signature of Approving Official:

James A. Tompkins, Product Manager (25)
 Herbicide Branch, Registration Division (7505P)

Date:

JUL - 1 2009

3. The handler PPE section must be revised to read:

“Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear;

Long-sleeved shirt and long pants

Shoes and socks, and

Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride.”

4. Revise the second sentence of the Environmental Hazards section to read “Do not contaminate water when disposing of equipment washwaters or **rinsate.**”

5. Delete the text “For Turf Use (except Sod Farms) must be deleted from the entry restriction text in the Non-Agricultural Use Requirements box.

6. The Statement Aerial application is prohibited,” must be added to the label.

7. On page 4, under Selection of Use Rates, revise the second paragraph to read “Apply this product at the rate of 1.0 to **2.6** ounces per acre or **0.0029** lbs a.i./1000 sq. ft.” Also, all reference to rates above 2.6 ounces per acre must be deleted from the label. This includes the rate listings and Weeds Controlled in this section.

Enclosed is a copy of your label stamped “Accepted With Comments”. This action is taken under the authority of section 4(g)(2)© of the Federal Insecticide, Fungicide, and Rodenticide Act as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

CORSAIR®

Selective Herbicide

A Water Dispersible Granule

For selective control of grassy and broadleaf weeds in turfgrass, including Sod Farms. Controls tall fescue, ryegrass, Virginia buttonweed, wild violets, chickweed, purslane, wild onion, wild garlic, and many other weeds.

ACTIVE INGREDIENT:

Chlorsulfuron: 2-Chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) aminocarbonyl]benzenesulfonamide 75%

OTHER INGREDIENTS:

25%

TOTAL: 100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JUL 1 - 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
228-375

EPA REG. NO. 228-375
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS 2 OZ.

000228-00375.20080213.EPA.Pending
chlorsulfuron DCI 8 month response

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May irritate eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Mixers, loaders, applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment washwaters.

There are no grazing restrictions on this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
<p>Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.</p> <p>Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.</p> <p>PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, shoes plus socks, chemical resistant gloves made of any waterproof material.</p>

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For Turf use (except Sod Farms), do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

DIRECTIONS FOR USE

This product can be applied on all turfgrass species except tall fescue, ryegrass, and dichondra. Some discoloration or delayed greenup may occur on St. Augustinegrass, Buffalograss, Centipedegrass, or Zoysiagrass, or Bahiagrass when this product is applied while grass is under environmental stress or during the spring transition period.

This product is a dispersible granule to be mixed in water and applied as a spot or broadcast spray treatment to certain species of established grasses. It is noncorrosive, nonflammable nonvolatile and does not freeze. This product rapidly inhibits growth of susceptible weeds. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 3 weeks after application depending on growing conditions and weed susceptibility. Warm moist conditions following treatment promote the activity of this product while cold, dry conditions delay the activity of this product. Weeds hardened off by cold weather or drought stress may not be fully controlled. Degree of control and duration of effect depend on: a) rate used; b) weed spectrum; c) weed size; d) growing conditions at and following time of treatment; e) soil pH; f) precipitation; and g) soil organic matter content.

Avoid drift at the application site. This product should be applied only when potential drift to adjacent sensitive area (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops, native plant communities) is minimal (e.g. when wind is blowing away from sensitive areas). Avoid application under conditions that may allow spray drift since very small quantities of spray may seriously injure susceptible crops during either active growth periods or dormancy. Follow the additional precautions below to minimize the potential for spray drift.

The interaction of many equipment and weather-related factors determines the potential for spray drift. The user is responsible for considering all these factors when making applications decisions.

Where states have more stringent regulations, they must be observed. The applicator should be familiar and take into account the information covered in the following:

Drift Control Adjuvants

A drift control adjuvant may be used to further reduce the potential for drift. If a drift control adjuvant is used, follow the use directions and precautions on the manufacturer's label. Do not use an adjuvant which increases viscosity with Microfoil, Thru-Value booms, or other systems that cannot accommodate viscous sprays.

Controlling Droplet Size:

Nozzle Type

Use a nozzle type according to manufacturer's specifications that is designed for the intended application and produces a Coarse to Very Coarse droplet size spectrum (ASAE S572) under application conditions. Applicators must consider nozzle orientation, nozzle pressure, and flight speed in determining droplet size. Nozzles should always be oriented in the manner that minimizes the effects of air shear. Solid stream nozzles oriented straight back produce the largest droplets and lowers drift.

Pressure

Do not exceed the nozzle manufacturer's recommended pressures. When higher flow rates are needed, use a higher capacity nozzle instead of increasing pressure.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore on up and downward edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment upwind. Swath adjustment distance should increase with increasing drift potential.

Wind

Drift potential is lowest with a sustained wind between 2 - 10 mph. However many factors including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided during gusty conditions and when winds are below 2 mph due to variable wind direction and high potential for temperature inversion. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are not hot and dry.

Surface Temperature Inversion

Applications must not occur during a local, surface temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds which are common during inversions. Temperature inversions are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog, however if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Shield Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

Boom Length/Height

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce. Limit nozzle to no greater than 4 feet above the top of the largest crops.

IMPORTANT: Do not use on food or feed crops. Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not apply where runoff water may flow onto agricultural land as injury to crops may result. Do not contaminate any body of water, including irrigation water. Do not use on walks, driveways, tennis courts, or similar areas. Prevent drift of spray to cropland and other desirable plants. Do not use on dichondra lawns or where violets or ryegrass are the desired species. Keep from contact with fertilizers, insecticides, fungicides and seeds. Following applications, the spray tanks should not be used for other than non-crop application. This is extremely important, as low rates of this product can kill or severely injure most crops. See SPRAYER CLEANUP section.

This product is to be applied as a post-emergence treatment with a hand held or a boom sprayer to control perennial "bunch or clump" grasses such as Tall fescue or Ryegrass and many other weeds in actively growing established Kentucky Bluegrass, Fine Fescue, Bentgrass (when mowed over 1/2 inch in height), Bahiagrass, and Bermudagrass For best results, apply this product when the desirable turf is not under any stress(i.e.,heat or drought). Not for use on golf tees or greens. Do not apply to desirable Ryegrass turf. Adequate moisture by rainfall or irrigation is needed to activate the herbicide. Do not apply to soil saturated with moisture or during periods of intense heat. Use a non-ionic surfactant at a rate of 1/4 to 1/3 oz. per gallon of spray solution.

Chemigation: Do not apply this product through any type of irrigation system.

RATES OF APPLICATION: This product is to be applied as a spot application with a hand held-spray or boom type sprayer. When spraying, only spray to wet the leaf blades. Over-application may result in temporary chlorosis of desirable grasses. This product may also suppress desirable grass growth and inhibit grass seedhead formation.

SELECTION OF USE RATES: Rate selection is bases on weed species, weed size , and soil texture. Use the higher rates on established plants and on the fine textured soil and lower rates on smaller weeds and coarse textured soils. For low volume or controlled droplet application, mix. This product according to application rates per acre. A repeat application may be necessary after 60 days.

Apply this product at a rate of 1.0 to 5.33 ounces per acre, or 0.0057 lbs. a.i./1,000 sq. ft. depending upon weed species. Use the low rate on soils with pH of 7.0 and above.

Weeds Controlled at 1 oz./Acre:

Annual ryegrass	Henbit	Shepherdspurse
Annual sowthistle	Pineappleweed	Smooth pigweed
Common chickweed	Prostrate pigweed	Wild mustard
Field pennygrass	Redroot pigweed	Yellow rocket

Weeds Controlled at 1 to 2 ounces /Acre:

Buttercup	Filaree	Perennial ryegrass
Common speedwell*	Goldenrod	Purslane
Common lambsquarter	Groundsel	Sweet clover*
Dandelion*	Mouseear chickweed	Wild parsnip

Weeds Controlled at 2 to 2.76 ounces /Acre:

Bedstraw	Common yarrow	Red clover
Burclover	Common mallow	Russian Thistle
Canada thistle*	Curly dock	Velvetgrass
Common mullein*	Horsetail	White clover
Common cinquefoil	Prostrate knotweed	Wild carrot

Weeds Controlled at 2.76 to 5.33 ounces /Acre:

Broadleaf Plantain*	Smutgrass+	Wild Onion
Pennsylvania Smartweed	Tall Fescue+	Wild Violet+
Ragweed*	Virginia Buttonweed*+	Wild Garlic

* To increase activity and spectrum of broadleaf weed control, this product can be tank mixed with 2,4-D, MCPP, DICAMBA, MCPA or TRICLOPYR at their lowest labeled rate for turfgrass. Follow all label precautions for use on turf. Apply with a non-ionic surfactant at a rate of 1/4 to 1/3 oz. per gallon of spray solution.

+ A second application 60 days after the first may be necessary.

APPLICATION RESTRICTIONS

Industrial Use sites (including rights-of-ways):

Application is restricted to ground application only. Do not apply more than three times per year. Do not apply more than 0.125 lbs ai./A per year.

Fine turf and ornamentals:

Broadcast application to lawns is prohibited. Application is limited to spot treatments of 240 square yards or less per acre.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pests to enter treated area until sprays have dried.

HOW TO USE

SPRAY PREPARATION: Add this product to water and agitate or shake thoroughly to uniformly disperse the water dispersible granules. Agitate the tank before each application. Use the mixed spray within 72 hours if the tank pH is below 5.0. This product is more stable in high pH mixtures. Avoid application in low pH conditions. Do not mix with acid forming compounds.

APPLICATION TECHNIQUE: Apply this product with a sprayer until leaf blades are wet. DO NOT OVER APPLY. Do not allow spray to drift into adjacent plants or trees, as injury may occur. Extreme care must be taken to prevent drift.

SPRAY VOLUME: For hand held or backpack sprayers, mix the correct amount of this product to spray 1 gallon per 1,000 sq. ft. For larger tank and boom-type sprayers, mix the recommended amount of this product in a minimum of 20 gallons to cover one acre. Do not over apply. Use the low rate on soils with pH of 7.0 and above.

AGITATION: Thoroughly mix and agitate spray solution. An anti-foam agent may be used to avoid excessive foaming.

SPRAYER CLEANUP: This product is a very active compound at low rates. To thoroughly remove this product from the spray tank, drain and flush with water a minimum of 10 minutes including all hoses, booms, and nozzles. Fill tank half full of water and add 1 to 2 gallons of ammonia or 4 lbs. of "Nutra-Sol" per 100 gallons of water. Fill tank and circulate for 15 minutes. Repeat procedure and rinse a final time with water. Clean nozzles and screens separately. Do not use the sprayer to treat any trees, shrubs, or ornamental plants after use with this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: This product should be stored in its original container in a cool, dry locked place out of the reach of children and out of direct sunlight. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds, after the flow begins to drip. Fill the container half full with water and soap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH

8/8

RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV021308R)

Corsair is a registered trademark of Nufarm Americas Inc.