

228-376

03-31-2004

114

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-3

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration	OPP Identifier Number 292605
		<input checked="" type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I

1. Company/Product Number 228-376	2. EPA Product Manager J. I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Riverdale Millennium Ultra Acid Weed & Feed	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. Riverdale - A Nufarm Company 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527-0866 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 3 1 2004
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Revised State use limitations for Oregon.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes * <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Russell F. Sawyer	Title Regulatory Affairs Manager	Telephone No. (include Area Code) 708/754-3330
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Affairs Manager	
4. Typed Name Russell F. Sawyer	5. Date February 25, 2004	

NOTIFICATION
MAR 31 2004
of 12
2/14

PLEASE SEE SPECIAL EDITOR'S NOTE ON LAST PAGE

(Residential Label)

RIVERDALE® MILLENNIUM ULTRA™ ACID WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.
Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:*

2,4-D Acid	0.861%
Clopyralid Acid	0.037%
Dicamba Acid	0.072%
INERT ¹ INGREDIENTS:	99.030%
TOTAL	100.000%

*Isomer Specific AOAC Method

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of fertilizer registrant)

Millennium Ultra is a Trademark and Riverdale is a Registered Trademark of Nufarm Americas Inc. Patent No. 5,266,553

²IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN NASSAU AND SUFFOLK COUNTIES. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY. IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL*, FOREST*, RIGHT-OF-WAY*, GOLF COURSE OR CEMETERY SITES.

*This product cannot be used on Agricultural, Forest Sites or Rights-of-Way.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-376

NET WEIGHT LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

- Revised 6/25/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting. Also added Special Editor's Note.
- 10/7/03 Added more State Restrictions for California and Oregon and added more weeds via Notification.
- 2/25/04 Corrected per EPA's e-mail dated 1/21/04 regarding Oregon restrictions via Notification.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse. Promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

MILLENNIUM ULTRA™ ACID WEED AND FEED is for use on Residential Turf.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping. **DO NOT USE GRASS CLIPPINGS FROM TURF TREATED WITH THIS PRODUCT FOR COMPOSTING AND/OR MULCH.**

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bident, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana

- 4/14

mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurge, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18)pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag)pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:°

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader

5/14

after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing **(No. of lbs.)** pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling
Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled** - If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 062503B)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra - A Premium Selective Herbicide

6/14

The following note will not be part of the actual label copy.

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

OTHER

*Please note, some states will require the term INERT and/or the full chemical name of the Active Ingredients.

Editor's Note:

The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The label prohibitions are optional for all other states.

Additional marketing and fertilizer hype:

- **CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS**
- **TIME-RELEASED NITROGEN PLUS WEED CONTROL FOR A GREEN, WEED-FREE LAWN**
- **CONTAINS IRON & SULFUR FOR EXTRA DEEP GREENING**
- **FOR FALL APPLICATION - CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS**
- **LAWN FOOD PROMOTES QUICK GREENING IN THE SPRING**

CONTAINS MILLENNIUM ULTRA™ HERBICIDE

<p>X NITROGEN: Controlled Slow Release for growing a dense, green lawn</p>	<p>X PHOSPHORUS: Improves root growth & plant food storage capabilities</p>	<p>X POTASSIUM: Vital for growth & improves hardiness & drought resistance</p>
--	---	--

7/14

**START YOUR YEAR ROUND
LAWN PROGRAM NOW!** See back for details

SATISFACTION
GUARANTEED

(SEAL)

If for any reason, you are not completely satisfied with the results after applying this product, you are entitled to product replacement or your money back simply by presenting evidence of purchase to your local (Seller Location)

Early Spring

Crabgrass
Preventer
plus Lawn
Food

Late Spring

Lawn Food or
Weed & Feed

(Seller's Name)

**YEAR ROUND
LAWN CARE
PROGRAM**

Fall

Winter Lawn
Food or
Winter Weed &
Feed

Summer

Super Fortifier
Plus
and/or
Insect Control

Alternate Spreader Information

Do not use your hands to spread this product.

Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.

BROADCAST SPREADER SETTINGS (One Pass Application)

For best results, apply in a circular path starting on the outside and working toward the center(see Path A). A little overlap of product will ensure complete coverage.

	3.6# Rate
Earl May Broadcast.....	6
Earl May (SB4000P).....	6
Precision Tractor TBS-90.....	6
Republic EZ 540/75450.....	3-4
Scotts EasyGreen.....	26% - 27%
Scotts SpeedyGreen.....	6

(Illustration)

8/14

DROP SPREADER SETTINGS (One Pass Application)

First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

	3.6# Rate
Earl May Drop.....	6½ - 7
Republic EZ 530/75350.....	7 - 8
Scotts Accugreen.....	6½
Scotts PF4 Metal.....	5 - 6½



Visit our web site at: ww.xxx.com

Special Editor's Note

Not to appear on final printed label.

The approved use of this label will automatically be terminated once the Clopyralid Technical, (Labeling approved by The Office of Pesticide Programs (OPP) prior to September 27, 2002) is depleted. Please Note; Clopyralid Technical marketed with labeling approved by OPP on September 27, 2002 (or later) prohibits formulating that Clopyralid Technical for use on residential turf. We will provide the Agency with updated labeling based on the new technical label before packaging. Any questions regarding these statements must be directed to Riverdale's Regulatory Affairs Manager at 1-800/345-3330.

PLEASE SEE SPECIAL EDITOR'S NOTE ON LAST PAGE

9/14

(Commercial Label)

RIVERDALE®

MILLENNIUM™ ULTRA ACID™ WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS:*

2,4-D Acid	0.861%
Clopyralid Acid	0.037%
Dicamba Acid	0.072%
INERT ¹ INGREDIENTS:	99.030%
TOTAL	100.000%

*Isomer Specific AOAC Method

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	

Fertilizer guaranteed by (Name of fertilizer registrant)

Millennium Ultra is a Trademark and Riverdale is a Registered Trademark of Nufarm Americas Inc. Patent No. 5,266,553

²IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN NASSAU AND SUFFOLK COUNTIES. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY. IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL*, FOREST*, RIGHT-OF-WAY*, GOLF COURSE OR CEMETERY SITES.

*This product cannot be used on Agricultural, Forest Sites or Rights-of-Way.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-376 NET WEIGHT LBS. EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

- Revised 6/25/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting. Also added Special Editor's Note.
- 10/7/03 Added more State Restrictions for California and Oregon and added more weeds via Notification.
- 2/25/04 Corrected per EPA's e-mail dated 1/21/04 regarding Oregon restrictions via Notification.

10/14

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse. Promptly and thoroughly wash hands and exposed skin with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRA™ ACID WEED AND FEED is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turf areas.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping. DO NOT USE GRASS CLIPPINGS FROM TURF TREATED WITH THIS PRODUCT FOR COMPOSTING AND/OR MULCH.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bidden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping

jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18)pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag)pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION: °

MAKE

SETTING

Central Calibrated
Cyclone Broadcast

5
3-3/4

12/14

Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.)* pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(RV 062503B)

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

12
13/14

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra - A Premium Selective Herbicide

The following note will not be part of the actual label copy.

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

OTHER

*Please note, some states will require the term INERT and/or the full chemical name of the Active Ingredients.

Editor's Note:

²The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The label prohibitions are optional for all other states.

Special Editor's Note

Not to appear on final printed label.

The approved use of this label will automatically be terminated once the Clopyralid Technical, (Labeling approved by The Office of Pesticide Programs (OPP) prior to September 27, 2002) is depleted. Please Note; Clopyralid Technical marketed with labeling approved by OPP on September 27, 2002 (or later) prohibits formulating that Clopyralid Technical for use on residential turf. We will provide the Agency with updated labeling based on the new technical label before packaging. Any questions regarding these statements must be directed to Riverdale's Regulatory Affairs Manager at 1-800/345-3330.



14/14
Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866
Phone: 708.754.3330 Fax: 708.754.0314
www.us.nufarm.com

UPS
February 25, 2004

Document Processing Desk (NOTIF)
Registration Division (7505C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Crystal Mall Building, #2
1921 Jefferson Davis
Arlington, VA 22202

Dear Sir/Madam:

Subject: Riverdale Millennium Ultra Acid Weed And Feed
EPA Registration No. 228-376
Label Revisions
Application For Pesticide: Other
OPP Identifier No. 292605

We are resubmitting corrected labels. We followed the guidance fact sheet in your January 21, 2004 e-mail covering an identical Oregon label problem.


We have enclosed an extra copy of the label highlighted to show **just the changes contained in this Notification.**

If there is anything else we should do at this time please contact us immediately.

Thank you for your continuing assistance.

Sincerely yours,

NUFARM AMERICAS, INC.


By Russell F. Sawyer
Regulatory Affairs Manager

RFS:jb

Enclosures