Rease read	28-3	377	7		05/	'15/	2001	an in part in a difficient star and a difficient start and a diffici				- <u> </u>
	one on re	verse	before complet Ui ronmental	nited S Prot	tates	-	ency	Form Approx		OMB No. Registra Amendi Other	ntion	OPP Identifier Number
1. Company/Product	Number			Appli	catio	n for	Pestici	de - Secti	on l	,	÷т	
4. Company/Produc Riverdale Millenr	t (Name) uium Ultr	228 га ^{тм} N				-	PM#	Product Manag	-	i Karana ang karana		None Restricted
5. Name and Address Riverdale Cher 1333 Burr Ridg Burr Ridge, IL Che	nical Co ge Parkv 60521-	ompan vay, Si 0866	у	ode)	-107		6. Exp (b)(i), n to: FPA I	Reg. No.	iw. simi	In accorda lar or iden	ance with tical in co	FIFRA Section 3(c)(3) mposition and labeling
						Se	ction -	ict Name	ા સ્ટ્ર	_ ##*, 48° 4		
Amendment Resubmissio	n in resp	onse t	o Agency letter	dated.		Final						
Adding the option use in Nassau and This notification is labeling or the Con statement to EPA. in violation of FIF	Suffolk (consisten fidential I further	countie ent with I Staten r under	s." 1 the provisions nent of Formula stand that if this	of PR	Notice 9 s produc cation is	98-10 an t. I und not con n and p	d EPA reg erstand tha isistent with	ulations at 40 C t it is a violatio h the terms of F ler sections 12	CFR 1 on of 1 PR No	.52.46, and 1 18 U.S.C. So ptice 98-10 a	no other cha ec. 1001 to and 40 CFR	willfun
1. Material This Pro		Be Pa	ckaged in:							r		
Certification n be submitted			Packaging Yes No s" Packaging wg	No. t. cont	per ainer	If "Y	r Soluble F Yes No es" tage wgt	Packaging No. per container		2. Type of	Container Metal Plastic Glass Paper Other (S	
S. Location of Net C	ontents i	nforma	ition	4. Size	e(s) Reta	ail Cont	ainer	- <u> </u>	5 Log	cation of La IOn Labe		ns
Label 6. Manner in Which		ontaine Affixed			Lithogr Paper g Stencil	glued	and the state of the	Other				panying product
					Staticity		ction - I	v			·	
1. Contact Point (Co	mpiete it	tems di	rectly below fo	r identi	fication				f nece	ssary, to pi	rocess this	
Name Russell F. Sawyer				1				Telephone	e No, (include Area Code) 4-3330			
l certify that the second seco	that any	y knowi	have made on ingly false or m	this fo	ing state	all attac	chments th nay be puni	ereto are true, ishable by fine	accu or im	prisonment	spolate. t dr	6. Date Application Recoved (Stamped)
	£1	Sa	γ				Regulato	ry Affairs Ma	inage	ar .		••••
4. Typed Name			0			5. Date						••••••
Russell F. Sawyer						May 4, 20	001					

EPA Form 8570-1 IRev. 8-94) Previous editions are obsolete.

т



Regulatory Affairs Department 1333 Burr Ridge Parkway, Suite 125A Burr Ridge IL 60521-0866 Office Ph: 708 754 3330 Facsimile: 708 754 0314

May 4, 2001

Document Processing Desk (NOTIF) Office of Pesticide Programs(7504C) U.S. Environmental Protection Agency Ariel Rios Building 1200 Pennsylvania Avenue, NW Washington, DC 20460-0001

Dear Sir/Madam:

Subject: Riverdale Millennium Ultra[™] N.Y. Weed And Feed EPA Reg. No. 228-377 Application for Pesticide: Other OPP Identifier No. 264005

Based on our June 1999 conversation with Ms. Joanne I. Miller, please find enclosed what we believe to be the necessary paperwork to notify the Agency of a labeling change. We have added to the front panel of our label the optional statement "In New York State this product may only be sold to and used by professional applicators and is prohibited from sale and use in Nassau and Suffolk counties." This is being done at the request of the state of New York as part of groundwater product stewardship.

If there are any questions regarding this, please feel free to contact us.

Sincerely yours,

RIVERDALE CHEMICAL COMPANY

By Russell F. Sawver

Regulatory Affairs Manager

RFS:jb

Enclosures

• • •

,

ł

İ

.

NOTIFICATION

RIVERDALE

MAY 1 5 2001

MILLENNIUM ULTRA™ N.Y. WEED AND FEED

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn. Treats (or covers) 5,000 Square Feet (50' x 100')

	ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid Dicamba (3,6-Dichloro-o-Anisic Acid)	d**	· · · ·	<u>99.145%</u>
(<pre>3y Isomer Specific AOAC Method, Equivalent to: *2,4-Dichlorophenoxyacetic Acid</pre>			0.573% 0.071%
	GUARANTEED ANALYSIS: [°] Total Nitrogen (N)			X8
	Available Phosphate (P ₂ O ₅)			X%
	Chlorine (Cl) Maximum			X% Lease
	Fertilizer guaranteed by (Name of fertilizer registrant)			
•	† Millennium Ultra™ - A Trademark of Riverdale Chemical Company			•
	In New York State this product may only be sold to and used applicators and is prohibited from sale and use in Nassau and a			
	KEEP OUT OF REACH OF CHILDREN			
	CAUTION			
	SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STA AND FIRST AID STATEMENT	ATEMENTS		
	NET WEIGHT LBS. EPA REG. NO. 228-377	EPA EST	. NO. 2	28-IL-1
	MANUFACTURED BY RIVERDALE CHEMICAL COMPANY BURR RIDGE, ILLINOIS 60521-0866		• • • • • • •	
	Revised 5/4/01 Added optional N.Y. County and Pro Applicator jb via notification.		restric	ctions,

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Commercial)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if (available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving herbicides such as 2,4-D and Clopyralid have been associated with mixing/loading and disposal sites. Additionally, using Clopyralid where soils are permeable, particularly where the water table is shallow, may also result in leaching to groundwater. Caution should be exercised when handling this (product at mixing, loading and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRATH N.Y. WEED AND FEED is for use on Ornamental Turf such as Lawns, Golf Courses ((Fairways, Aprons, Tees* and Roughs), Parks, Cemeteries, Athletic fields and similar non-crop areas. *Excluding Bentgrass Tees.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada

· 4/14 Page 2 thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Nutgrass, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild fouro'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

5/14 Page 3

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18)pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag)pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.

SPREADER SETTINGS ONE PASS APPLICATION:°

MAKE

Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

SETTING

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing <u>*(No. of lbs.)</u> pounds of nitrogen per 1,000 square feet (optional P,O, and K,O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. *Based on Inert fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

alisia Katisa Katisa 1.23

Page 4

f

(RV 061200A)

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Commercial)

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra™ - A Premium Selective Herbicide



Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

The following note will not be part of the actual label copy.

"Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

RIVERDALE

NOTIFICATION

8/14

MILLENNIUM ULTRA™ N.Y. WEED AND FEED

MAY 1 5 20C1

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

ACTIVE INGREDIENTS: †	
Dimethylamine Salt of $2,4-D^*$	690%
Monoethanolamine Salt of Clopyralid**	094%
Dicamba Acid	071%
NERT ¹ INGREDIENTS:	<u>145%</u>
TOTAL	0008
By Isomer Specific AOAC Method, Equivalent to:	
*2,4-D Acid	573%
**Clopyralid Acid	071%
GUARANTEED ANALYSIS:°	
Total Nitrogen (N)	. X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate $(P_{2}O_{5})$	
Soluble Potash (K ₂ O)	
Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Tri	ple
Super Phosphate, and Muriate of Potash.	
Chlorine (Cl) Maximum	
The urea nitrogen in this product has been coated to provide X% coated slow releas	e
urea nitrogen.	

Fertilizer guaranteed by (Name of fertilizer registrant)

† Millennium Ultra™ - A Trademark of Riverdale Chemical Company

In New york State this product may only be sold to and used by professional applicators and is prohibited from sale and use in Nassau an Suffolk counties.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FIRST AID STATEMENT

WEIGHT LBS.

EPA REG. NO. 228-377

EPA EST. NO. 228-IL-1

MANUFACTURED BY RIVERDALE CHEMICAL COMPANY BURR RIDGE, ILLINOIS 60521-0866

Revised 5/4/01 Added optional N.Y. County and Pro Appicator Sale & Use restrictions, jb via notification.

Page 2

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Non-Commercial)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syprup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

MILLENNIUM ULTRATM N.Y. WEED AND FEED is for use on Ornamental Turf such as Lawns, Golf Courses (Fairways, Aprons, Tees* and Roughs), Parks, Cemeteries, Athletic Fields and similar non-crop areas. *Excluding Bentgrass Tees.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Blackeyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervsin, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadlerf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistlc, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefcil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Cormon sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida gusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Redge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed,

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Non-Commercial)

Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Nutgrass, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild fouro'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Wooly morningglory, Woodsorrel, Wooly croton, Wooly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20(or 18) pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 4(or 3.6 for 18 lb. bag)pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1(.9 for 18 lb. bag) pound per 250 square feet.



10/14

· 11/14

Page 4

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Non-Commercial)

SPREADER SETTINGS ONE PASS APPLICATION:°

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing <u>*(No. of lbs.)</u> pounds of nitrogen per 1,000 square feet (optional P,O, and K,O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. *Based on Inert fertilizer claimed.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizers, seed or other pesticides.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RV 061200A)

. 4 x 1 5 x ες. ₁ 1-1-1-1

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

Contains Millennium Ultra™ - A Premium Selective Herbicide



benefits to your home and environment:
Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to oxygen for a family of 4 every day

- oxygen for a family of 4 every day ✓ Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & other Pollutants
- Purifies Water & Modifies Temperature

- Improves Soil Condition & Controls Erosion

Besides increasing property values, did you know a well-maintained lawn provides the following

Note: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer

reduce this amount, leave grass clippings on your lawn.

Increased mowing height and frequency will produce

smaller grass clippings that will decompose more readily

so nutrients go back into your lawn.

The following note will not be part of the actual label copy.

guarantee are optional except where mandated by State fertilizer regulation.

Optional Marketing Statement(s)

For Both Cool Season & Warm Season Grasses

Alternate Statements

¹OTHER*

Ĺ

*Please note, some states may still require the term INERT and/or the full chemical name of the Active Ingredients.

Page 6

Additional marketing and fertilizer hype:

- CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS
- TIME-RELEASED NITROGEN PLUS WEED CONTROL FOR A GREEN, WEED-FREE LAWN
- CONTAINS IRON & SULFUR FOR EXTRA DEEP GREENING
- FOR FALL APPLICATION CONTROLS CLOVER, DANDELIONS & MANY OTHER BROADLEAF WEEDS
- LAWN FOOD PROMOTES QUICK GREENING IN THE SPRING

CONTAINS

MILLENNIUM ULTRATM HERBICIDE

x	x	x
NITROGEN:	PHOSPHORUS:	POTASSIUM:
Controlled Slow	Improves root	Vital for growth
Release for	growth & plant	& improves
growing a dense,	food storage	hardiness &
green lawn	capabilities	drought resistance

START YOUR YEAR ROUND LAWN PROGRAM NOW! See back for details

SATISFACTION GUARANTEED (SEAL)

If for any reason, you are not completely satisfied with the results after applying this product, you are entitled to product replacement or your money back simply by presenting evidence of purchase to your local (Seller Location)

Riverdale Millennium Ultra™ N.Y. Weed and Feed (Non-Commercial)

Early Spring Crabgrass Preventer plus Lawn		Late Spring Lawn Food or Weed & Feed
Food	(Seller's Name)	
	YEAR ROUND LAWN CARE PROGRAM	
Fall Winter Lawn Food or Winter Weed & Feed		Summer Super Fortifier Plus and/or Insect Control

Alternate Spreader Information

́ь

Do not use your hands to spread this product.

Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.

BROADCAST SPREADER SETTINGS (One Pass Application) For best results, apply in a circular path starting on the outside and working toward the center(see Path A). A little overlap of product will ensure complete coverage.

	3.6# Rate	
Earl May Broadcast	6	
Earl May (SB4000P)	6	
Precision Tractor TBS-90	6	(Illustration)
Republic EZ 540/75450	3-4	(IIIUSCIACION)
Scotts EasyGreen	26½ - 27½	
Scotts SpeedyGreen	6	

DROP SPREADER SETTINGS (One Pass Application) First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

> **3.6# Rate** 6½ - 7

7 - 8 6½

5 - 6½

Earl May	Drop
Republic	EZ 530/75350
Scotts A	ccugreen
Scotts P	F4 Metal

Visit our web site at: ww.xxx.com