

228-391

12/17/2008

1/32



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

DEC 17 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Lizabeth Rea
Nufarm Americas, Inc.
150 Harvester Drive
Suite 200
Burr Ridge, IL 60527

Subject: EPA Reg. 228-391 (Patriot) Label Amendment

Dear Ms. Rea:


The labeling referred to above is acceptable provided you make the following changes:

- 1) Add "exists" after "washables" on page 1 of the label.
- 2) On page 7, change "recommended" to "to be used".
- 3) On page 5, 9, 12, 22, 23, and 24 (and anywhere else in the label when talking about rates) change "recommended rate" to "specified rate".
- 4) On page 2, change "as recommended" to "as directed". On page 3 change "recommendations" to "use directions". Delete "Nufarm will not be responsible... such non-recommended use". This type of language belongs in the Warranty section.
- 5) Throughout the label remove "recommended" from the directions for use sites and replace with "is to be used".
- 6) On page 15, replace "label recommendations" and "those recommendations" with "label instructions".
- 7) On page 20, replace "recommended below" with "specified below". On page 21, change "recommended intervals" to "intervals". On page 20, replace "such as" with "including".
- 8) On page 13, specify a product name for 2,4-D under rush skeletonweed and spotted knapweed.

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik

Page 2
EPA Reg. 228-391

Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

DEC 17 2008
Jim Tompkias
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Patriot

ACTIVE INGREDIENT:	By Weight
Metsulfuron methyl: Methyl2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	60.0%
OTHER INGREDIENTS:	40.0%
TOTAL:	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

EPA Reg. No. 228-391

EPA Est. No.

Manufactured for:
Nufarm Americas, Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527


Net Weight:

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

12-17-08

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-391



USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff water may flow, during periods of intense rainfall or to water saturated soils as off-target movement and injury may occur. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely affected from drift and run-off.

IMPORTANT

DO NOT USE ON FOOD OR FEED CROPS EXCEPT AS RECOMMENDED BY THIS LABEL. Injury to or loss of desirable trees or other plants may result if the precautions listed below are not followed.

- Do not apply Patriot (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the product may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas except as recommended by this label.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water.
- Keep from contact with fertilizers, insecticides, fungicides and seeds.

Spraying and mixing equipment used with Patriot must not be used for subsequent applications to food or feed crops with the exception of pastures, rangeland and wheat, as low rates of Patriot can kill or severely injure most food or feed crops.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistance weed biotypes, it may be necessary to change cultural practices within, and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as a part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

Patriot should be used only in accordance with recommendations on this label.

Nufarm will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by Nufarm. User assumes all risks associated with such non-recommended use.

Do not apply more than 4 ounces Patriot per acre per year.

Do not use on food or feed crops except as recommended by this label.

TANK MIXES

Patriot may be tank mixed with other herbicides registered for the use sites described in this label. Use only those tank mix partners which are labeled for the appropriate use site. When tank mixing, use the most restrictive label limitations for each of the products being used in the tank mix.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS DURING THE RESTRICTED-ENTRY INTERVAL (REI) OF 4 HOURS.

PPE required for early entry that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

Non-crop industrial weed control, selective weed control in turf (industrial, unimproved only), and weed control in pastures and rangeland are not within the scope of the Worker Protection Standard.

GENERAL INFORMATION

Patriot is a dispersible granule that is mixed in water and applied as a spray. Patriot controls many annual and perennial weeds and woody plants in non-crop areas, conifer and hardwood plantations. Patriot may also be used on pastures, or CRP as well as selected uncultivated agricultural areas (fence rows,

farmyards, and rights-of-way) directly adjacent to treated pastures or rangeland, where grazing or harvesting for animal feed may occur.

Patriot may be used for general weed and brush control and for the control of certain noxious weeds on noncrop sites, ditchbanks of dry drainage ditches and for selective weed control in certain types of unimproved turfgrass. Do not use on irrigation ditches. Patriot can also be used for controlling and suppressing undesirable weeds and hardwoods in conifer plantations and weeds in hardwood plantations.

Patriot controls weeds and woody plants primarily by postemergent activity. Although Patriot has preemergence activity, best results are generally obtained when Patriot is applied to foliage after emergence or dormancy break. Generally, for the control of annual weeds, Patriot provides best results when applied to young, actively growing weeds. For the control of perennial weeds, applications made at the bud/bloom stage or while the target weeds are in the fall rosette stage may provide the best results. The use rate depends upon the weed species and size at the time of application.

The degree and duration of control may depend on the following:

- Weed spectrum and infestation intensity
- Weed size at application
- Environmental conditions at and following treatment
- Soil pH, soil moisture, and soil organic matter.

Patriot may be applied on conifer and hardwood plantations and noncrop sites that contain areas of temporary surface water caused by the collection of water between planting beds, in equipment ruts, or in other depressions created by management activities. It is permissible to treat intermittently flooded low lying sites when no water is present. It is also permissible to treat marshes, swamps, and bogs after water has receded as well as seasonally dry food deltas. Do not make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams and canals.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

Patriot is absorbed primarily through the foliage of plants, and by the roots to a lesser degree. Plant cell division is generally inhibited in sensitive plants within a few hours following uptake. Two to 4 weeks after application, leaf growth slows followed by discoloration and tissue death. The final effects on annual weeds are evident about 4 to 6 weeks after application. The ultimate effect on perennial weeds or woody plants occurs in the growing season following application.

Warm, moist conditions following treatment promote the activity of Patriot while cold dry conditions may reduce or delay activity. Weeds and brush hardened off by cold weather or drought stress may not be controlled.

One to two inches of rainfall or sprinkler irrigation (enough to wet the top 2-3 inches of soil profile) may be needed to move Patriot into the weed root zone before the next flush of weeds emerge. The amount of

moisture required for sufficient activation increases with crop or weed residue and for finer textured soils. Without sufficient rainfall or sprinkler irrigation to move Patriot into the weed root zone, weeds that germinate after treatment will not be controlled.

Application of Patriot provides the best control in vigorously growing grasses that shade competitive weeds. Weed control in areas of thin grass may not be as satisfactory. However, a grass canopy that is too dense at application can intercept a spray and reduce weed control.

Patriot is safe to grasses under normal conditions. However, grasses that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices may be injured by applications of Patriot. In addition, different species of grass may be sensitive to treatment with Patriot under otherwise normal conditions. Application of Patriot to these species may result in injury.

The use of a surfactant is recommended to enhance the control of susceptible plants, except where noted. Apply at a minimum rate (concentration) of ¼% volume/volume (1 quart per 100 gallons of spray solution) or at the manufacturer's recommended rate. Use only EPA approved surfactants containing at least 80% active ingredient. Certain types of surfactants, such as those incorporating acetic acid (i.e. LI-700), may not be compatible with Patriot and may result in decreased performance. Certain surfactants may not be suitable for use on desirable plants such as turf and conifers, listed on this label. Consult the surfactant manufacturer's label for appropriate uses.

Weed and brush control may be reduced if rainfall, snowfall or sprinkler irrigation occurs within 4 hours following application.

AGRICULTURAL USES

CONIFER PLANTATIONS

Application Information

Patriot is recommended for the control of many species of weeds and deciduous trees on sites where conifers are growing or are to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" and "Brush Species Controlled" for a listing of susceptible species.

Application Timing

Apply Patriot after weeds have emerged or after undesirable hardwoods have broken winter dormancy and have reached the point of full leaf expansion.

Conifer Site Preparation

-- Application Before Transplanting

After consulting the "Weeds Controlled" and "Brush Species Controlled" tables, apply the rates of Patriot recommended for the most difficult to control species on the site.

Southeast- Apply up to 4 ounces per acre for loblolly and slash pines. Transplant the following planting season.

Northeast and Lake states- Apply up to 2 ounces per acre for red pine. Transplant the following planting season. Apply up to 2 ounces per acre for black, white and Norway spruce. Transplant the following spring.

West- Apply up to 2 ounces per acre prior to planting Douglas Fir, Sitka Spruce, Western Red Cedar, Western Hemlock, Ponderosa Pine, and Grand Fir in the Coast Rangeland and western slope of the Cascades in Oregon and Washington. These conifer species listed can be planted anytime after application. Other conifer species can be planted providing the user has prior experience indicating acceptable tolerance to Patriot soil residues.

Without prior experience, it is recommended that other species be planted on a small scale to determine selectivity before large-scale plantings are made as unacceptable injury may occur. Nufarm will not assume responsibility for injury to any conifer species not listed on this label.

TANK MIX COMBINATIONS

For broader spectrum control, the following products are recommended in combination with Patriot.

Razor® Herbicide

Tank mix 1 to 2 ounces of Patriot with 10 to 24 fluid ounces of Razor per acre. Refer to the product container for a list of species controlled.

Nufarm Polaris® AC Herbicide Applicator's Concentrate

Tank mix 1 to 2 ounces of Patriot with 10 to 24 fluid ounces of Nufarm Polaris® AC Applicator's Concentrate per acre. Loblolly and slash pines may be transplanted the planting season following the application. This combination controls, ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, persimmon, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, and red maple.

Razor + Nufarm Polaris® AC Herbicide Applicators Concentrate

Tank mix ½ to 1 ounce of Patriot with 16 to 64 fluid ounces of Razor and 10 to 12 fluid ounces of Nufarm Polaris AC Applicator's Concentrate per acre. Slash and loblolly pines may be transplanted the planting season following application. This combinations controls, cherry, dogwood, elms, oaks (red and water), persimmon, sassafras, sweetgum and suppresses hickory.

VELPAR® L or VELPAR® DF

Tank mix 1 to 2 ounces of Patriot per acre with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. Loblolly and slash pines may be transplanted the planting season following application. Refer to the product container for a list of species controlled.

OUST® EXTRA

Tank mix ½ to 1 ½ ounces of Patriot with 2 to 3 ounces of OUST® EXTRA per acre for herbaceous weed control. Refer to the product container and the "Weeds Controlled" section of this label for a listing of the weeds controlled. Loblolly and slash pines may be transplanted the planting season following application. Tank mix 2 ounces of Patriot with 3 ounces of OUST® EXTRA per acre for herbaceous weed control and early spring suppression of bull thistle and Canada thistle in the Coast Rangeland and western slope of the Cascade Mountains. Douglas fir may be transplanted at least 90 days following application.

RELEASE—HARDWOOD CONTROL AND SUPPRESSION

Patriot is recommended for application over the top of established slash and loblolly pine to control the species listed in "Weeds Controlled" and "Brush Species Controlled" section of this label. Apply 1 to 4 ounces per acre to control the species indicated, including kudzu.

Tank Mix Combinations--

For broader spectrum control the following products are recommended in combination with Patriot.

Nufarm Polaris® AC Herbicide Applicator's Concentrate

Tank mix 1 to 2 ounces of Patriot with 8 to 16 fluid ounces of Nufarm Polaris® AC Herbicide Applicator's Concentrate per acre for application to loblolly pine. Refer to the Nufarm Polaris® AC Herbicide Applicator's Concentrate label regarding the use of surfactants and the appropriate application timing with respect to the age and development stage of the pines. This combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, oaks (red, white and water), sassafras, sweetgum, Vaccinium species and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, persimmon, and red maple.

VELPAR® L or VELPAR® DF

Tank mix 1 to 2 ounces of Patriot with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. This combination may be applied to loblolly and slash pines.

RELEASE—HERBACEOUS WEED CONTROL

Patriot may be applied to transplanted loblolly and slash pine for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when Patriot is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations-

For broader spectrum control the following products are recommended in combination with Patriot.

Nufarm Polaris® AC Herbicide Applicators Concentrate

Tank mix ½ to 1 ounce of Patriot with 4 fluid ounces of Nufarm Polaris® AC Applicators Concentrate per acre. The tank mix may be used on loblolly pine.

Spyder®

Tank mix ½ to 1 ½ ounces of Patriot with 2 to 3 ounces of Spyder per acre. Best results are obtained when Patriot is applied just before weed emergence until shortly after weed emergence. This tank mix may be used on loblolly and slash pine.

VELPAR® L or VELPAR® DF

Tank mix ½ to 1 ounce of Patriot with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. This combination may be applied to loblolly and slash pines.

IMPORTANT PRECAUTIONS

--CONIFER PLANTATIONS ONLY

- Applications of Patriot made to conifers that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock or other stresses may injure or kill the trees.
- Applications of Patriot made for herbaceous release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- Do not apply Patriot to conifers grown as ornamentals.
- Patriot applications may result in damage and mortality to other species of conifers when they are present on sites with those listed in the preceding recommendations for conifer plantations.

HARDWOOD PLANTATIONS

Application Information

Patriot is recommended at rates of up to 2 ounces per acre for the control of many weed species on sites where yellow poplar is growing or is to be planted, and on sites where red alder is to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" section of this label for a listing of susceptible species.

Application Timing

Patriot may be applied as a site preparation treatment prior to planting red alder or yellow poplar. As a prior to planting site preparation treatment for red alder, Patriot may be tank mixed with other herbicides labeled for this use.

Patriot may also be applied over-the-top of planted yellow poplar seedlings after the soil has settled around the root system, but before the seedlings have broken dormancy (prior to bud break).

Release—Herbaceous Weed Control

Patriot may be applied to yellow poplar for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when Patriot is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations-

Tank mix 1/2 ounce of Patriot with 4 to 6 pints of VELPAR® L as recommended on the package label for "RELEASE - HERBACEOUS WEED CONTROL" in pine plantations in the eastern U.S. Follow the VELPAR® L label recommendations regarding altering the application rate by soil texture.

IMPORTANT PRECAUTIONS

-- HARDWOOD PLANTATIONS ONLY

- Application of VELPAR® L and Patriot made to yellow poplar that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock or other stresses may injure or kill the seedlings.
- Applications of Patriot made for release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- The use of surfactant is not recommended for applications made over the tops of trees.
- Careful consideration must be given by an experienced and knowledgeable forester to match the requirements of yellow poplar and/or red alder to conditions of the site. Treatment of yellow poplar and/or red alder planted on a site inadequate to meet its requirements may injure or kill the seedlings.

PASTURE, RANGELAND AND CONSERVATION RESERVE PROGRAM (CRP)

APPLICATION INFORMATION FOR GRASS ESTABLISHMENT IN PASTURE, RANGELAND AND CONSERVATION RESERVE PROGRAM (CRP)

Use Patriot for the suppression or control of broadleaf weeds to aid in the establishment of the following perennial native or improved grasses planted in pasture, rangeland or acres enrolled in the Conservation Reserve Program (CRP):

- Blue grama
- Bluestems - big, little, plains, sand, WW spar
- Buffalograss
- Green sprangletop
- Indiangrass
- Kleingrass
- Lovegrasses - atherstone, sand, weeping, wilman
- Orchardgrass
- Sideoats grama
- Switchgrass - Blackwell
- Wheatgrasses - bluebunch, crested, intermediate, pubescent, Siberian, slender, streambank, tall, thickspike, western
- Wildrye grass - Russian

Consult with the Natural Resources and Conservation Service or other local experts concerning planting techniques and other cultural practices to maximize potential for grass establishment.

Due to the inability of newly planted grass stands to sufficiently compete with weeds, and the severity of weed pressure in new grass stands, performance from Patriot may not always be satisfactory. An additional herbicide application or mowing may be needed.

Use Rates and Application Timing for Grass Establishment in Pasture, Rangeland and CRP

Preplant (prior to planting) or Preemergence (after planting but before grass emergence)

Apply Patriot preplant or preemergence at 1/10 ounce/acre on all labeled grasses except orchardgrass and Russian wildrye grass. Do not apply Patriot preplant or preemergence to orchardgrass and Russian wildrye grass as severe crop injury may result.

Early Postemergence to New Plantings

Apply Patriot at 1/10 ounce/acre, plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution on all labeled grasses anytime after grass emergence. Do not use a spray adjuvant other than non-ionic surfactant.

Because grass species differ in time of emergence, apply only after majority of grasses are in the 3 to 4 leaf stage.

Postemergence to stands with 1-5 leaf grasses planted the previous season

Apply Patriot at 1/10 ounce/acre plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution, on all labeled grasses when the majority of the grasses have one or more leaves. Do not use a spray adjuvant other than non-ionic surfactant.

APPLICATION INFORMATION FOR ESTABLISHED GRASSES IN PASTURE, RANGELAND AND CONSERVATION RESERVE PROGRAM (CRP)

Use Rates for Established Pastures, Rangeland and CRP

Apply 1/10 to 1 ounce of Patriot per acre as a broadcast application to established grasses in pasture rangeland and CRP. For spot application, use 1 ounce per 100 gallons of water. Do not apply more than 1 2/3 ounces of Patriot per acre per year.

Application Timing – Established Pastures, Rangeland and CRP

Patriot may be applied to established native grasses such as bluestems and grama, and on other established pasture grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

Pasture Grass	Minimum time from grass establishment to Patriot application
Bermudagrass	2 months
Bluegrass, bromegrass, and orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Fescue Precautions:

Note that Patriot may temporarily stunt fescue, cause yellowing or seed suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 oz./A of Patriot
- Tank mix Patriot with Weedestroy® AM-40 Amine Salt and other 2,4-D products.
- Use the lowest recommended rate for target weeds
- Use a non-ionic surfactant at ½ to 1 pint per 100 gallons of spray solution (1/16 to 1/8% v/v)
- Make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall
- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use a spray adjuvant other than non-ionic surfactant

The first cutting yields may be reduced due to seedhead suppression resulting from treatment with Patriot.

Timothy Precautions:

Timothy should be at least 6" tall at application and be actively growing. Applications of Patriot to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 oz./A of Patriot
- Tank mix Patriot with Weedestroy AM-40 Amine Salt and other 2,4-D products.
- Use the lowest recommended rate for target weeds
- Use a non-ionic surfactant at ½ pint per 100 gallons

- Make application in the late summer or fall
- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use a spray adjuvant other than non-ionic surfactant

Application of Patriot to Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison's creeping foxtail may cause severe injury to and /or loss of pastures.

Other Pasture and Rangeland Grasses: Varieties and species of forage grasses differ in their tolerance to herbicides. When using Patriot on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated the following season. Broadleaf pasture species such as alfalfa and clover are highly sensitive to Patriot and will be severely stunted or injured by Patriot.

WEEDS AND BRUSH CONTROLLED OR SUPPRESSED IN PASTURES, RANGELAND AND CONSERVATION RESERVE PROGRAM (CRP)

Unless otherwise directed, treat when weeds are less than 4" tall or in diameter and are actively growing.

Before using Patriot, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your pasture, rangeland or CRP acres at the same time.

1/10 OUNCE PER ACRE

Bitter sneezeweed
 Blue/purple mustard*
 Broomweed, common
 Bur buttercup (testiculate)
 Buttercup
 Canada thistle*‡
 Carolina geranium
 Coast fiddleneck (tarweed)
 Common chickweed
 Common mullein
 Common Purslane
 Conical catchfly
 Corn gromwell*‡
 Cowcockle
 Curly dock
 Cutleaf evening primrose*‡
 Dandelion
 False chamomile
 Field pennycress (fanweed)
 Filaree
 Flixweed*
 Groundsel (common)
 Henbit
 Kochia*
 Lambsquarters (common, slimleaf)

Marestail
 Mayweed chamomile
 Miners lettuce
 Pigweed (redroot, smooth, tumble)
 Plains coreopsis
 Plantain
 Prickly lettuce*
 Prostrate knotweed*‡
 Russian thistle*
 Shepherd's purse
 Smallseed falseflax
 Smartweed (green, ladythumb, pale)
 Snow speedwell
 Tansymustard*
 Treacle mustard (Bushy Wallflower)
 Tumble/Jim Hill mustard
 Volunteer sunflower*
 Waterpod
 Wild buckwheat*‡
 Wild garlic*
 Wild mustard
 Wild sunflower*‡
 Woolly croton*

2/10 OUNCE PER ACRE

Annual marshelder
 Blackeyed-Susan
 Buckbrush‡
 Burclover
 Common yarrow
 Dogfennel

Horsemint (beebalm)
 Musk thistle*
 Purple scabious
 Scotch thistle*
 Western snowberry‡
 Wild carrot

3/10 to 1/2 OUNCE PER ACRE

Annual sowthistle
 Aster
 Bittercress
 Chicory
 Clover
 Cocklebur
 Corn cockle
 Crown vetch
 Goldenrod
 Maximillion sunflower
 Multiflora rose*‡
 Pennsylvania smartweed

Pensacola bahiagrass*
 Redstem filaree
 Rough fleabane
 Seaside arrowgrass
 Sericea lespedeza*
 Silky crazywood (locoweed)
 Sweet clover
 Wild lettuce
 Wood sorrel
 Yankeeweed

1/2 to 1 OUNCE PER ACRE

Black henbane
 Blackberry
 Broom snakeweed
 Buckhorn plantain
 Common crupina
 Dewberry
 Dyer's woad
 Gorse
 Halogeton

Honeysuckle
 Multiflora rose and other wild roses*
 Plumeless thistle
 Rosering gaillardia
 Spotted knapweed*
 Teasel
 Wild caraway
 Yucca*‡

1 OUNCE PER ACRE

Bull thistle
 Common tansy
 Field bindweed‡
 Gumweed
 Houndstongue
 Perennial Pepperweed
 Poison hemlock
 Purple loosestrife

Rush skeletonweed*‡
 Salsify
 Scouringrush
 Snowberry
 St. Johnswort
 Western salsify
 Whitetop (hoary cress)

*See the Specific Weed Problems section of this label.

‡Weed suppression is a reduction in weed competition (reduced population and/or vigor) as visually compared to an untreated area. The degree of suppression varies with the rate used, the size of the weeds, and the environmental conditions following treatment.

SPOT APPLICATIONS FOR THE SUPPRESSION‡ OF WEEDS AND BRUSH**APPLICATION INFORMATION FOR SPOT APPLICATIONS**

Patriot is recommended for the suppression of the following undesirable weed and brush species growing in pastures, rangeland or CRP using spot applications. Spot applications may be made by using equipment such as back pack sprayers or hand sprayers. Patriot should be applied as a spray to the foliage and stems. The application volume required will vary with the height and density of the brush and the application equipment used. Regardless of the application volume and equipment used, thorough coverage of the foliage and stems is necessary to optimize results. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage. Add a non-ionic surfactant having at least 80% active ingredient at 2-4 pints per 100 gallons of spray solution.

Use Rates for Spot Application

Mix 1 ounce of Patriot per 100 gallons of water.

Application Timing for Spot Applications

Make a foliar application of the recommended rate of Patriot during the period from full leaf expansion in the spring until the development of full fall coloration.

Weed and Brush Species Suppressed with Spot Applications

Blackberry‡ Dewberry‡
Canada Thistle*‡ Multiflora Rose‡

*See the Specific Weed Problems section.

‡Weed and brush suppression is a reduction in weed and brush competition (reduced population and/or vigor) as visually compared to an untreated area. The degree of suppression varies with the rate used, the size of the weeds, and the environmental conditions following treatment.

SPECIFIC WEED PROBLEMS

Note: Thorough spray coverage of all weed species listed below is very important.

Blue/Purple Mustard, Flixweed, and Tansymustard: For best results, apply Patriot tank mixtures with Weedestroy® AM-40 Amine Salt and other 2,4-D products or MCPA postemergence to mustards, but before bloom.

Broom Snakeweed: For best results, apply Patriot at ½ ounce/acre in the fall. Applications of Patriot in the spring will provide suppression only.

Canada Thistle: For suppression with broadcast applications, apply either Patriot or Patriot plus Weedestroy® AM-40 Amine Salt and other 2,4-D products or MCPA in the spring after the majority of thistles have emerged and are small (rosette stage to 6" elongating stems) and actively growing. The application will inhibit the ability of emerged thistles to compete with grass.

For suppression with spot applications, apply as a foliar spray once plant is fully leaved.

Corn Gromwell, Cutleaf Evening Primrose and Prostrate Knotweed: Apply Patriot when weeds are actively growing, are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing Weedestroy® AM-40 Amine Salt and other 2,4-D products or MCPA with Patriot can improve results.

Kochia, Russian thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use Patriot in a tank mix with Veteran 720 and Weedestroy® AM-40 Amine Salt and other 2,4-D products. Patriot should be applied in the spring when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing.

Multiflora Rose: For control with broadcast applications, apply Patriot at ½ ounce per acre as a broadcast application. For control with foliar applied spot applications, apply Patriot at 1 ounce per 100 gallons of water.

For suppression with broadcast applications, apply Patriot at rates of 3/10 up to ½ ounce per acre. Applications should be made in the spring, soon after multiflora rose is fully leafed and is less than 3 feet tall.

For control with Spotgun Basal Soil Treatment, prepare a spray suspension of Patriot by mixing 1 ounce per gallon water. Mix vigorously until the Patriot is dispersed and agitate periodically while applying the spray suspension. Apply the spray preparation with an exact delivery handgun applicator. Apply at the rate of 4 ml for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of the stem union. When treating large plants and more than one delivery is required, make applications on opposite sides of the plant.

Make applications from early spring to summer.

Musk Thistle, Scotch Thistle: Apply Patriot at 2/10 to ¼ ounce per acre in the spring or early summer prior to flowering or in the fall after newly emerged plants have reached the rosette stage of growth. Certain biotypes of Musk and Scotch Thistles are less sensitive to Patriot and may not be controlled with Patriot rates less than ¼ ounce per acre. Consult with your local Nufarm representative, dealer or

applicator for specific use rate and tank mix recommendations for your area. Fall applications should be made before the soil freezes.

Pensacola bahiagrass control in established Bermudagrass pasture: Apply Patriot at 3/10 ounce per acre after green-up in the spring but before bahiagrass seedhead formation. Application should be made when moisture is sufficient to enhance grass growth.

Patriot is very effective for removal of bahiagrass from bermudagrass pastures. In highly infested pastures, the use of Patriot can clear the areas of useful forage until the bermudagrass has time to cover the area. Therefore, Patriot treatments should be spread out over a period of years. Do not apply to an entire farm or ranch in one year. Fertilization (particularly with nitrogen and potassium) and/or replanting may accelerate the process of reestablishment of bermudagrass.

Under heavy bahiagrass pressure, grazing pressure, or adverse weather conditions (heat and drought), bahiagrass regrowth may occur.

Do not use Patriot for control of common or Argentine bahiagrass. Also, do not use Patriot in liquid fertilizer solutions for Pensacola bahiagrass control as poor control and/or regrowth may occur.

Rush skeletonweed: For best results, apply Patriot at 1 ounce per acre with 8 fluid ounces of dicamba (such as Diablo or Veteran 720) and 16 fluid ounces of 2,4-D.

Sericea lespedeza: For best results, apply Patriot at 4/10 to ½ ounce per acre beginning at flower bud initiation through the full bloom stage of growth. Consult with your local Nufarm representative, dealer or applicator for specific use rate recommendations for your area. Do not make applications if drought conditions exist at intended time of application.

Spotted Knapweed: For best results, apply Patriot at ½ ounce per acre with 8 fluid ounces of dicamba (such as Diablo or Veteran 720) and 16 ounces active ingredient per acre of 2,4-D.

Sunflower (wild or volunteer): Apply either Patriot or Patriot plus Weedestroy® AM-40 Amine Salt and other 2,4-D products or MCPA after the majority of sunflowers have emerged, are 2" to 4" tall and are actively growing. Use spray volumes of at least 3 gallons by air or 10 gallons by ground.

Wild Buckwheat: For best results, apply Patriot plus Weedestroy® AM-40 Amine Salt and other 2,4-D products or MCPA when plants have no more than 3 true leaves (not counting cotyledons). If plants are not actively growing, delay treatment until environmental conditions favor active weed growth.

Wild Garlic: Apply 1/10 to 2/10 ounce per acre of Patriot in the early spring when wild garlic is less than 12" tall with 2" to 4" of new growth.

Woolly Croton: Apply 1/10 to 2/10 ounce per acre of Patriot in the late spring or early summer from cotyledon through 2 true leaf stage.

Yucca: For best results, apply Patriot at ½ to ¾ ounce per acre plus 2,4-D, Diablo, Veteran 720, or Relegate from two weeks before blooming to two weeks after blooming.

SPRAY ADJUVANTS

Unless otherwise directed on this label, Patriot applications must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer can be used unless specifically prohibited by tank mix partner labeling. Consult your local Nufarm representative prior to using other adjuvant systems. If another herbicide is tank mixed with Patriot, select adjuvants authorized for use with both products. Products must contain only EPA exempt ingredients (40 CFR 1001).

Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- MSO adjuvants may be used at 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactants (NIS)

- Apply at 0.25% v/v (1 quart per 100 gallons spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLD) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 pounds/acre of a spray grade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by Nufarm. Consult your local Nufarm representative before using adjuvant types not specified on this label.

Exceptions: (1) On Fescue pastures use ½ to 1-pint non-ionic surfactant per 100 gallons; (2) on Timothy pastures use ½ pint non-ionic surfactant per 100 gallons.

Antifoaming agents may be used if needed.

Do not use low rates of liquid fertilizer as a substitute for surfactant.

GROUND APPLICATION

To obtain optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flood nozzles on 30" spacings, use at least 10 gallons per acre (GPA), flood nozzles no larger than TK10 (or equivalent), and a pressure of at least 30 pounds per square inch (psi). For 40" nozzle spacings, use at least 13 GPA; for 60" spacings, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

With "Raindrop RA" nozzles, use at least 30 GPA and ensure that nozzle spray patterns overlap 100%.

For flat-fan nozzles, use at least 10 GPA for broadcast applications to pasture, rangeland or CRP.

Use 50-mesh screens or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage. Use a minimum of 2 GPA. In Idaho, Oregon and Washington, use a minimum of 3 GPA.

When applying Patriot by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the Spray Drift Management section of this label.

TANK MIXTURES

With Insecticides and Fungicides

Patriot may be tank mixed or used sequentially with insecticides and fungicides registered for use on pastures, rangeland or CRP. However, under certain conditions (drought stress or cold weather), tank mixes or sequential applications of Patriot with organophosphate insecticides (such as parathion) may produce temporary grass yellowing or, in severe cases, grass injury.

The potential for grass injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application.

Test these tank mixtures in a small area before treating large areas. Do not use Patriot plus Malathion, as grass injury will result.

With Herbicides

Patriot may be tank mixed with other suitable registered herbicides to control weeds listed under Weeds Suppressed, weeds resistant to Patriot, or weeds not listed under Weeds Controlled. Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with Patriot.

Herbicide Tank Mixtures for Pastures or Rangeland:

For postemergence control of the following weeds in pastures or rangeland:

Annual marshelder	Common milkweed
Burclover	Common ragweed
Carolina horsenettle	Giant ragweed
Common cocklebur	Western ragweed

Apply Patriot at 1/10 to 1 ounce per acre in a tank mix with one of the following products. Refer to companion herbicide labels to confirm that the product is labeled for control of the weeds listed above and is registered for use in your state.

Product	Rate (ounce product/A)
Trooper P+D	8 to 32
Trooper 22K	4 to 16
Weedmaster	8 to 32
Relegate	8
Amber	0.35*

*For suppression of Western Ragweed in Phenoxy Restricted and Herbicide Regulated counties.

Product	Rate (ounces a.i./A)
2,4-D	8 to 16
Dicamba (such as Diablo or Veteran 720)	2 to 16
2,4-D + Veteran 720	1 + 2.87 to 4 + 11.48

Herbicide Tank Mixtures for CRP:

Preplant

Patriot may be tank mixed with glyphosate as a pre-plant (prior to the planting of CRP grasses) treatment to control broadleaf and grassy weeds. When using a glyphosate tank mix, allow at least 7 days after application before planting grasses. Refer to glyphosate containing product fact sheets and labels for all use instructions, label rates, weed control claims, and precautions.

Postemergence

For best weed control performance in CRP, use Patriot in a tank mix with 2,4-D (ester formulations perform best) or dicamba (such as Diablo or Veteran 720).

Patriot can be tank mixed with 2,4-D at ¼ pound a.i./A for all labeled grasses larger than the 5-leaf stage. For fully tillered stands, up to ½ pound a.i./A of 2,4-D may be used. A spray adjuvant may be added. However, the addition of spray adjuvant may increase the chance of grass injury.

Patriot can also be tank mixed with dicamba (such as Diablo or Veteran 720). Use not more than 1/8 to ¼ pound a.i./A of Diablo plus Patriot after majority of grasses are in the 3-leaf stage. In established grasses (2nd year stands), use not more than ¼ to ½ pound a.i./A Diablo plus Patriot. A spray adjuvant may be added. However, the addition of spray adjuvant may increase the chance of grass injury.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing Patriot in fertilizer solution.

Patriot must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the Patriot is added. Use of this mixture may result in temporary grass yellowing and stunting.

If using low rates of liquid nitrogen fertilizer (between 5% and 50% of the spray solution volume) in the spray solution, the addition of a non-ionic surfactant is necessary. Add surfactant at $\frac{1}{4}$ pint per 100 gallons of spray solution (0.03% v/v).

Do not use a spray adjuvant other than non-ionic surfactant.

When using high rates of liquid nitrogen fertilizer (greater than or equal to 50% of the spray solution volume) in the spray solution, adding spray adjuvant(s) increases the risk of grass injury. Consult your agricultural dealer, consultant, fieldman, or Nufarm representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with Patriot and liquid nitrogen fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Do not add spray adjuvants when using Patriot in tank mix with 2,4-D ester and liquid nitrogen fertilizer solutions greater than 5% of the spray volume.

The use of liquid nitrogen fertilizer solutions greater than 5% of the spray volume with Patriot rates greater than 0.5 ounce/acre may cause grass injury.

Do not use low rates of liquid fertilizer as a substitute for spray adjuvants.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

Rotation Intervals in Pasture, Rangeland or CRP for Overseeding and Renovation

Location	Crop or Grass Species	Maximum Patriot Rate on Pasture (ounce/acre)	Minimum Rotation Interval (months)
AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, ryegrass, tall fescue	1/10 to 3/10	4
	Wheat (except durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
All states not included above	Red clover, white clover and sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, ryegrass	1/10 to 2/10	6
	Tall Fescue	1/10 to 2/10	18
	Wheat (except durum)	1/10 to 2/10	1
All areas with soil pH of 7.5 or less	Russian wildrye	1/10 to 1/2	1
	Green needlegrass, switchgrass, sheep fescue	1/10 to 1	1
	Meadow brome, smooth brome, alta fescue, red fescue, meadow foxtail, orchardgrass, Russian wildrye, timothy	1/10 to 1	2
All areas with soil pH of 7.9 or less	Alkali sacaton, mountain brome, blue grama, thickspike wheatgrass	1/10 to 1	1
	Sideoats grama, switchgrass	1/10 to 1/2	2
	Western wheatgrass	1/10 to 1	2
	Sideoats grama, switchgrass, big bluestem	1/10 to 1	3
AL, AR, FL, GA, KS, KY, LA, MS, MO, NC, OK, SC, TN, TX, VA, WV, with soil pH of 7.0 or less	STS soybeans	1/10 to 2/10	6
	Field corn	1/10 to 2/10	12

CROP ROTATION

Before using Patriot, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your pasture, rangeland or CRP acres at the same time.

Minimum Rotational Intervals

Minimum rotation intervals* are determined by the rate of breakdown of Patriot applied. Patriot in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase Patriot breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow Patriot breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

*The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

Do not apply Patriot on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Patriot could remain in the soil for 34 months

or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of Patriot.

Checking Soil pH

Before using Patriot, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

Bioassay

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table.

To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow the following year in fields previously treated with Patriot. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or Nufarm representative for information detailing the field bioassay procedure.

Grazing/Haying


There are no grazing or haying restrictions for Patriot. Coveralls, shoes plus socks must be worn if cutting within 4 hours of treatment.

NON-AGRICULTURAL USES

WEEDS CONTROLLED

1/3 TO 1/2 OUNCE PER ACRE

Annual sowthistle	Goldenrod
Aster	Lambsquarters
Bahiagrass	Marestail/horseweed ****
Beebalm	Maximillion sunflower
Bittercress	Miners lettuce
Bitter sneezeweed	Pennsylvania smartweed
Blackeyed-susan	Plains coreopsis
Blue mustard	Plantain
Bur buttercup	Redroot pigweed
Chicory	Redstem filaree
Clover	Rough fleabane
Cocklebur	Sheperd's-purse
Common chickweed	Silky crazyweed (locoweed)
Common groundsel	Smallseed falseflax
Common purslane	Smooth pigweed
Common yarrow	Sweet clover
Conical catchfly	Tansymustard
Corn cockle	Treacle mustard
Cow cockle	Tumble mustard
Crown vetch	Wild carrot
Dandelion	Wild garlic
Dogfennel	Wild lettuce
False chamomile	Wild mustard
Fiddleneck tarweed	Wooly croton
Field pennycress	Wood sorrel
Flixweed	Yankeweed



1/2 TO 1 OUNCE PER ACRE

Blackberry	Honeysuckle
Black henbane	Multiflora rose and other
Broom snakeweed*	wild roses
Buckhorn plantain	Musk thistle***
Bull thistle	Oxeye daisy
Common crupina	Plumeless thistle
Common sunflower	Prostrate knotweed
Curly dock	Rosering gaillardia
Dewberry	Seaside arrowgrass
Dyer's woad	Sericea lespedeza
Gorse	Tansy ragwort
Halogeton	Teasel
Henbit	Wild caraway

1 TO 2 OUNCES PER ACRE

Common mullein	Purple loosestrife
Common tansy	Purple scabious
Field bindweed**	Scotch thistle
Greasewood	Scouringrush
Gumweed	Salsify
Houndstongue	Snowberry
Lupine	St. Johnswort
Old world climbing fern (Lygodium)	Sulphur cinquefoil
Perennial pepperwood	Western salsify
Poison hemlock	Whitetop (hoary cress)
	Wild iris

1 1/2 TO 2 OUNCES PER ACRE

Canada thistle**	Tall larkspur
Dalmation toadflax**	Wild parsnip
Duncecap larkspur	Yellow toadflax**
Russian knapweed**	

3 TO 4 OUNCES PER ACRE

Kudzu

* Apply fall through spring

** Suppression, which is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Apply as a full coverage spray for best performance.

*** Certain biotypes of musk thistle are more sensitive to Patriot and may be controlled with rates of 1/4 to 1/2 ounce per acre. Treatments of Patriot may be applied from rosette through bloom stages of development.

**** Certain biotypes of maretail/horsetail are less sensitive to Patriot and may be controlled by tank mixes with herbicides with a different mode of action.

PROBLEM WEED CONTROL

For broader spectrum control and for use on certain biotypes of broadleaf weeds which may be resistant to Patriot and herbicides with the same mode of action, the following tank mixes are recommended.

Diablo + 2,4-D

Weed	Rate of Patriot (ounces per acre)	Rate of Diablo (fl oz/acre)	Rate of 2,4-D (fl oz/acre)
Kochia control	1/2	8	16
Spotted knapweed Control	1/2	8	16
Rush skelentonweed suppression	1	8	16

NONCROP SITES

Application Information

Patriot is recommended for general weed control on private, public and military lands as follows: Uncultivated areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas- noncrop producing (such as farmyards, fuel storage areas, fence rows, soil bank land, barrier strips, etc.); industrial sites- outdoor (such as lumberyards, pipeline and tank farms, etc.). It is also recommended for the control of certain noxious and troublesome weeds.

Consult the "Weeds Controlled" and "Brush Species Controlled" tables to determine the appropriate application rate.

Patriot may be applied in tank mixture with other herbicides labeled for use on non-crop sites. Fully read the labels and follow all the directions and restrictions on each label.


Applications may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the target vegetation with the application equipment being used.

Application Timing

For best results, Patriot should be applied postemergence to young, actively growing weeds. Applications may be made at any time of the year, except when the ground is frozen.

GRASS REPLANT INTERVALS

Following an application of Patriot to non-crop areas, the treated sites may be replanted with various species of grasses at the intervals recommended below.



For soils with a pH of 7.5 or less, observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Brome, Meadow	½ - 1	2
	1-2	3
Brome, Smooth	½ - 1	2
	1-2	4
Fescue, Alta	½ - 1	2
	1-2	4
Fescue, Red	½ - 1	2
	1-2	4
Fescue, Sheep	½ - 1	1
	1-2	4
Foxtail, Meadow	½ - 1	2
	1-2	4
Green Needlegrass	½ - 2	1
Orchardgrass	½ - 1	2
	1-2	4
Russian wildrye	½ - 1	1
	1	2
	2	3
Switchgrass	½ - 1	1
	1-2	3
Timothy	½ - 1	2
	1-2	4
Wheatgrass, Western	½ - 1	2
	1-2	3

For soils with a pH of 7.5 or greater, observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Alkali Sacaton	½ - 1	1
	1-2	3
Bluestem, Big	½ - 2	3
Brome, Mountain	½ - 1	1
	1-2	2
Gramma, Blue	½ - 2	1
Gramma, Sideoats	½	2
	>1/2	>3
Switchgrass	½	2
	>1/2	>3
Wheatgrass, Thickspike	½ - 2	1
Wheatgrass, Western	½ - 1	2
	1 - 2	3

The recommended intervals are for applications made in the Spring to early Summer. Because Patriot degradation is slowed by cold or frozen soils, applications made in the late Summer or Fall should consider the intervals as beginning in the Spring following treatment.

Testing has indicated that there is considerable variation in response among the species of grasses when seeded into areas treated with Patriot. If species other than those listed above are to be planted into

areas treated with Patriot, a field bioassay should be performed, or previous experience may be used, to determine the feasibility of replanting treated sites.

TURF, INDUSTRIAL (UNIMPROVED ONLY)

Application Information

Patriot is recommended for selective weed control in unimproved industrial turf where certain grasses are well established and desired as ground cover. Patriot is also recommended for the control of certain noxious and troublesome weeds in turf.

In addition to conventional spray equipment, Patriot may also be applied with invert emulsion equipment. When using an invert emulsion, mix the prescribed rate of Patriot in the water phase.

Consult the "Weeds Controlled" table to determine which weeds will be controlled by the following recommendations:

Turf type	Rate of Patriot (ounces/acre)
Fescue and Bluegrass	1/4 to 1/2
Crested Wheatgrass and Smooth Brome	1/4 to 1
Bermudagrass	1/4 to 2

Application Timing

Applications may be made at anytime of the year, except when the soil is frozen.

When a spring application is made on fescue or bluegrass, a second application may be made during the summer after full seedhead maturation.

**GROWTH SUPPRESSION AND SEEDHEAD INHIBITION
(Chemical Mowing)**

Application Information

Patriot is recommended for growth suppression and seedhead inhibition in well established fescue and bluegrass turf at the use rate of 1/4 to 1/2 ounce per acre.

Tank Mix Combination

Patriot may be tank mixed with T-Pac E-Pro MEC for improved performance in the regulation of growth and seedhead suppression. Tank mix 1/4 to 1/2 ounce of Patriot with 2 to 4 ounces of T-Pac E-Pro MEC

Application Timing

Application may be made after at least 2 to 3 inches of new growth has emerged until the appearance of the seed stalk.

Fescue Precautions:

Patriot may temporarily stunt tall fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce per acre of Patriot.
- Tank mix Patriot with 2,4-D.
- Use the lowest recommended rate for target weeds.
- Use a non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution.
- Make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall.
- Do not use a surfactant when liquid nitrogen is used as a carrier.
- Do not use a spray adjuvant other than non-ionic surfactant.
- The yields from the first cutting may be reduced due to seedhead suppression resulting from treatment with Patriot.

IMPORTANT PRECAUTIONS
-- INDUSTRIAL TURF ONLY

- An application of Patriot may cause temporary discoloration (chlorosis) of the grasses. Use the lower recommended rates for minimum discoloration.
- With fescue and bluegrass, sequential applications made during the same or consecutive growth periods (i.e. spring and fall) may result in excessive injury to turf.
- Excessive injury may result when Patriot is applied to turf that is under stress from drought, insects, disease, cold temperatures (winter injury) or poor fertility.
- Patriot is not recommended for use on bahiagrass.

NATIVE GRASSES

Patriot is recommended for weed control and suppression in the establishment and maintenance of native grasses. It may be used where blue grama, bluestems (big, little, plains, sand, ww spar) bromegrasses (meadow), buffalograss, green sprangletop, indiagrass, kleingrass, lovegrasses, (atherstone, sand, weeping, wilman), orchardgrass, sideoats grama, switchgrass (blackwell), wheatgrass (bluebunch, intermediate, pubescent, Siberian, slender streamband, tall, thickspike, western), and Russian wildrye are established. It may also be applied over these species in the seedling stage, except for orchardgrass and Russian wildrye.

Application Information

Apply Patriot at the rate of 1/10 ounce per acre for the control and suppression* of bur buttercup (testiculate), common purslane, common sunflower*, cutleaf eveningprimrose*, flixweed*, lambsquarters* (common and slimleaf), marestail*, pigweed (redroot and tumble), snow speedwell, tansymustard* and tumble mustard (Jim Hill mustard).

*Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Degree of suppression will vary with the size of weed and environmental conditions following treatment.

Application Timing

For established grasses, apply when weeds are in the seedling stage.

For grasses in the seedling stage, apply preplant or preemergence where the soil (seedbed) has been cultivated.

IMPORTANT PRECAUTIONS
--NATIVE GRASSES

- Grass species or varieties may differ in their response to various herbicides. Nufarm recommends that you first consult your state experimental station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of Patriot to a small area. Components in a grass seed mixture will vary in tolerance to Patriot, so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after Patriot application, temporary discoloration and/or grass injury may occur. Patriot should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.

BRUSH CONTROL

Application Information

Patriot is recommended for the control of undesirable brush growing in non-crop areas. Applications may be made by air, high volume ground application, low volume and ground applications Except as noted for multiflora rose, Patriot should be applied as a spray to the foliage.

The application volume required will vary with the height and density of the brush and the application equipment used. Generally, for aerial and ground applications use sufficient volumes of water to insure thorough coverage.

Regardless of application volume and equipment used, thorough coverage of the foliage, particularly the terminal growing points, is necessary for optimum results.

BRUSH SPECIES CONTROLLED

Species	High Volume Rate (ounces/75 gallons)	Broadcast Rate (ounces/acre)
Ash	1-2	1-3
Aspen	1-2	1-3
Black Locust	1-2	1-3
Blackberry	1-2	1-3
Camelthorn	1-2	1-3
Cherry	1-2	1-3
Cottonwood	1-2	2-3
Eastern red cedar	1-2	2-3
Elder	1-2	2-3
Elm	1-2	1-3
Firs	3	1-2
Hawthorn	1-2	1-3
Honeysuckle	1-2	1/2-1
Mulberry	1-2	2-3
Multiflora rose	1-2	1-3
Muscadine (wild grape)	1-2	2-3
Oaks	1-2	1-3
Ocean spray (<i>Holodiscus</i>)	1-2	2-3
Osage orange	1-2	2-3
Red maple	1-2	2-3
Salmonberry	1/2-1	1-3
Snowberry	1/2-1	1-3
Spruce (black and white)	3	2-3
Thimbleberry	1/2-1	1-3
Tree of heaven (<i>Ailanthus</i>)	1-2	1-2
Tulip tree	1/2-1	1-3
Wild roses	1/2-1	1-3
Willow	1/2-1	1-3

For low volume and ultra-low volume ground applications, mix 4 to 8 ounces of Patriot per 75 gallons of spray solution.

Application Timing

Make a foliar application of the recommended rate of Patriot during the period of full leaf expansion in the spring until the development of full fall coloration on deciduous species to be controlled. Coniferous species may be treated at anytime during the growing season.

Tank Mix Combinations-

Patriot may be tank mixed with any product labeled for non-crop brush control at the application rates specified on the companion product's label for the species specified on the product's companion label.

Read and follow the label instructions of both products when tank mixing. Follow the most restrictive limitations of the products labels being tank mixed.

Low Rate Applications

Nufarm Polaris® Herbicide

Combine 1 to 2 ounces of Patriot with 1 to 4 pints of Nufarm Polaris Herbicide per acre and apply as a broadcast spray. Aerial applications should use a minimum of 5 gallons per acre spray volume. In addition to species listed above controlled by Patriot, this combination controls black gum, hophornbeam, sassafras, sweetgum, Vaccinium species, dogwood, myrtle dahoon, hickories, and persimmon.

Trooper 22K + Nufarm Polaris Herbicide

Combine 1 to 1 1/2 ounce of Patriot with 2 to 8 fluid ounces of Nufarm Polaris and 1 to 2 pints of Trooper 22K per 75 gallons of water. Apply as a high volume spray. This tank mix controls cherry, elms, box elder, maples, hackberry, redbud, ash, oaks (including shingle oak), black locust and sassafras.

Trooper 22K is a restricted use pesticide

Spotgun Basal Soil Treatment

For control of multiflora rose, prepare a spray suspension of Patriot by mixing 1 ounce per gallon of water. Mix vigorously until the Patriot is dispersed and agitate periodically while applying the spray suspension.

Apply the spray preparation with an exact delivery handgun applicator. Apply at the rate of 4 milliliters for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of stem union. When treating large plants and more than one delivery is required, make applications on opposite sides of the plant.

Applications should be made from early spring to summer.

IMPORTANT PRECAUTIONS

-- NON-CROP BRUSH ONLY

- When using tank mixtures of Patriot with companion herbicides, read and follow all the use instructions, application rates, warnings and precautions appearing on the labels. Follow the most restrictive label instruction for each of the herbicides used.

SPRAY EQUIPMENT

Spraying and mixing equipment used with Patriot must not be used for subsequent application to food or feed crops with the exception of pastures, rangeland and wheat, as low rates of Patriot can kill or severely injure most food or feed crops.

The selected sprayer should be equipped with an agitation system to keep Patriot suspended in the spray tank.

Use a sufficient volume of water to thoroughly cover the foliage of undesirable weeds, generally 30 to 50 gallons per acre. Select a spray volume and delivery system that will deliver a uniform spray pattern. Be sure the sprayer is calibrated before use. Avoid overlapping and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to desired plants.

Refer to the brush control section of this label for information unique to that particular use.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of Patriot.
3. Continue agitation until the Patriot is fully dispersed, at least 5 minutes.
4. Once the Patriot is fully dispersed, maintain agitation and continue filling tank with water. Patriot should be thoroughly mixed with water before adding any other material.

5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Patriot spray preparations are stable if they are pH neutral or alkaline and stored at or below 100°F.
8. If Patriot and a tank mix partner are to be applied in multiple loads, pre-slurry the Patriot in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Patriot.
9. All tank mixes with Patriot should be sprayed and the spray tank emptied at the end of each days work.

SPRAYER CLEANUP

Spray equipment must be cleaned before Patriot is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined below.

At the End of the Day

When multiple loads of Patriot are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gallons of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse tank, boom, and hoses with clean water.
6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

*Equivalent amounts of an alternate-strength ammonia solution or an Nufarm-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult you agricultural dealer, applicator, or Nufarm representative for a listing of approved cleaners.

Notes:

1. **ATTENTION:** Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When Patriot is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to the cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making applications.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150-200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!**

See, **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size- General Techniques

- **Volume-** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rate flows produce larger droplets.
- **Pressure-** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type-** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size- Aircraft

- **Number of Nozzles-** Use the minimum number of nozzles with the highest flow rate that provides uniform coverage.
- **Nozzle Orientation-** Orienting nozzles so that the spray is emitted backwards, parallel to the air stream will produce larger droplets than other orientations.
- **Nozzle Type-** Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length-** The boom length should not exceed $\frac{3}{4}$ of the wing or rotor length- longer booms increase drift potential.
- **Application Height-** Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS. Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from the ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind

conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

USE PRECAUTIONS

- Do not drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the product may be washed or moved into contact with their roots, as injury or loss of desirable trees or other plants may result.
- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Exposure to Patriot may injure or kill most crops. Injury may be more severe when crops are irrigated. Do not apply Patriot when these conditions are identified and powdery, dry soil or light, and sandy soils are known to be prevalent in the area being treated.
- Applications made where runoff water flows onto agricultural land may injure crops. Applications made during periods of intense rainfall, to soils saturated with water, to surfaces paved with materials such as asphalt or concrete, or to soils through which rainfall will not readily penetrate may result in runoff and movement of Patriot. Do not treat frozen soil. Treated soil should be left undisturbed to reduce the potential for Patriot movement by soil erosion due to wind or water.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Do not apply through any type of irrigation system.
- Do not apply to irrigated land where the tailwater will be used to irrigate crops.
- Do not apply to snow-covered ground.
- Spraying and mixing equipment used with Patriot must not be used for subsequent application to food or feed crops with the exception of pastures, rangeland, and wheat, as low rates of Patriot can kill or severely injure most food or feed crops.
- Applications of Patriot to pastures, rangeland or CRP undersown with legumes may cause injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of Patriot.
- When used as directed, there are no grazing or haying restrictions for use rates of 1 2/3 ounces per acre and less. At use rates of 1 2/3 to 3 2/3 ounces per acre, forage grasses may be cut for hay fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos
- Do not use this product in California.
- Do not use on grasses grown for seed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Amber® is a trademark of Syngenta Crop Protection, Inc.

Diablo®, Polaris®, Razor®, Relegate®, Spyder®, Trooper®, Veteran®, and Weeddestroy® are trademarks of Nufarm Americas, Inc.

Weedmaster® is a trademark of BASF Ag Products.

Velpar[®] and Oust[®] are trademarks of E.I. du Pont de Nemours and Company.
Weedmaster[®] is a trademark of Velsicol Chemical Corporation.

oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo
oooo