



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
228-401

Date of Issuance:
APR 24 2003

Term of Issuance:
Conditional

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name of Pesticide Product:
Riverdale Sweet
Sixteen Weed and Feed
With DEAPADA

Name and Address of Registrant (include ZIP Code):

Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60521-0866

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label changes listed below before you release the product for shipment:
 - a. Add the phrase, "EPA Reg. No. 228-401".
 - b. On the "Commercial" and "Non-Commercial" labeling delete "and similar non-crop areas". This phrase implies that this product could be applied to sites other than ornamental turf. On the "Non-Commercial" labeling delete "Golf Courses (Fairways, Aprons, Tees and Roughs)", "Parks" and "Cemeteries". These are commercial turf sites and are not appropriate for a homeowner use label.

Signature of Approving Official:

191

Date:

APR 24 2003

page 2
EPA Reg. No. 228-401

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

4. Submit the results of storage stability (GRN 830.6317) and corrosion characteristics (GRN 830.6320) studies to EPA upon completion.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3/10

(Commercial)

RIVERDALE®

SWEET SIXTEEN
WEED AND FEED WITH DEAPADA™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	1.346%
2,4-Dichlorophenoxyacetic Acid*	0.182%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.359%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.090%

INERT INGREDIENTS:	98.023%
TOTAL	100.000%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	1.076%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.359%
***3,6-Dichloro-o-Anisic Acid	0.090%

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

GUARANTEED ANALYSIS:

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash(K ₂ O)	X%
Primary Plant Foods derived from (List sources).	

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer registrant)

DEAPADA is a Trademark of Riverdale Chemical Company. Patent Pending
Riverdale and The Optical Advantage are Registered Trademarks of Riverdale Chemical Co.

ACCEPTED
with COMMENTS
In EPA Letter Dated:

APR 24 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
028-401

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-

NET WEIGHT 16 LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
BURR RIDGE, ILLINOIS 60527-0866



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH DEAPADA is a stable mixture of three herbicides. It is for use on Ornamental Turf such as, Lawns, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, ~~Common Bermuda, Blue Grass~~, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf, Drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow,

5/10

Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange Hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .8 lb. per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizers, seeds, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL MARKETING STATEMENTS

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration is optional except where mandated by State fertilizer regulation.

7/10

(Non-Commercial)

RIVERDALE®

SWEET SIXTEEN
WEED AND FEED WITH DEAPADA™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	1.346%
2,4-Dichlorophenoxyacetic Acid*	0.182%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.359%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.090%
INERT INGREDIENTS:	98.023%
TOTAL	100.000%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	1.076%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.359%
***3,6-Dichloro-o-Anisic Acid	0.090%

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash(K ₂ O)	X%
Primary Plant Foods derived from (List sources).	
Chlorine (Cl) Maximum	X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

Fertilizer guaranteed by (Name of Fertilizer registrant)

DEAPADA is a Trademark of Riverdale Chemical Company. Patent Pending
Riverdale and The Optical Advantage are Registered Trademarks of Riverdale Chemical Co.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

APR 24 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
EPA REG. NO. 228-40

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 16 LBS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
BURR RIDGE, ILLINOIS 60527-0866

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SWEET SIXTEEN WEED AND FEED WITH DEAPADA™ is a stable mixture of three herbicides. It is for use on Ornamental Turf such as, Lawns, Golf Courses (Fairways, Aprons, Tees and Roughs), Parks, Cemeteries and other similar non-crop areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bidden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf dymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherds-purse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet

9/10

leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .8 lb. per 250 square feet.

SPREADER SETTINGS

ONE PASS APPLICATION:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.)* pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizers, seeds or other pesticides. Do not stack more than two pallets high.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled** - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration is optional except where mandated by State fertilizer regulation.