

228-401

3/19/2010

1 of 14



**U.S. ENVIRONMENTAL PROTECTION AGENCY**  
**Office of Pesticide Programs**  
**Registration Division (7505P)**  
**Ariel Rios Building**  
**1200 Pennsylvania Ave., NW**  
**Washington, D.C. 20460**

EPA Reg. Number:

228-401

Date of Issuance:

MAR 19 2010

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Sweet Sixteen Weed and Feed with Viper

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.  
 150 Harvester Drive, Suite 200  
 Burr Ridge, IL 60527

~~Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA registration number.~~

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make all the following changes to the product label:
  - a. Add the following statement to both labels:  
 "Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/litstatus/wtc/index.htm>"
  - b. On page 2 of the Residential label, change the text to read "The maximum number of broadcast applications per treatment site is 2 per year."
  - c. On page 4 of both labels, change the phrase to read "Possible retreatment in approximately 30 days..."
  - d. On page 4 of both labels, change the sentence to read "Do not exceed recommended specified rates because damage to turf may occur."

Continued on Page 2

Signature of Approving Official:

Kathryn V. Montague  
 Product Manager 23  
 Herbicide Branch  
 Registration Division (7505P)

Date:

MAR 19 2010

Zof 14

- e. On the Residential label, revise the Storage and Disposal section as follows:
- i. Add the statement "Do not contaminate water, food, or feed by storage and disposal." directly below the Storage and Disposal heading.
  - ii. Revise the heading from "Pesticide Container Disposal" to read "Pesticide Disposal and Container Handling"
  - iii. Revise the Pesticide Disposal and Container Handling text to read "Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain."
- f. On both labels, correct the typo to "DELIVERED TO YOU" in the Warranty Disclaimer section.

The basic confidential statement of formula (CSF) dated November 6, 2007 is acceptable.

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after twelve (12) months from the date of this notice or the next printing of the label, whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

Commercial

# SWEET SIXTEEN WEED AND FEED WITH VIPER™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.  
Treats (or covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

**ACTIVE INGREDIENTS:**

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	1.346%
2,4-Dichlorophenoxyacetic Acid*	0.182%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	0.359%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.090%

**OTHER INGREDIENTS:**

TOTAL: 98.023%  
100.000%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	1.076%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.359%
***3,6-Dichloro-o-anisic Acid	0.090%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

**GUARANTEED ANALYSIS:**

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Primary Plant Foods derived from (List sources).	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	
Fertilizer guaranteed by (Name of fertilizer registrant)	

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION - PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-401  
EPA EST. NO. 228-IL-1

MANUFACTURED FOR  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS 16 LBS. 000228-00401.COM 20100111- EPA FPL

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**MAR 19 2010**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-401

40914

Commercial

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION - PRECAUCION**

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

**PERSONAL PROTECTION EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**All loaders, applicators, and other handlers must wear:**

- long-sleeved shirt and long pants,
- shoes plus socks, and
- chemical-resistant gloves

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**FIRST AID**

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistem or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

This product is a stable mixture of four herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

Not for use on turf grown for sale or other commercial use as sod, or commercial seed production, or for research purposes.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

- |                            |                       |                        |                            |
|----------------------------|-----------------------|------------------------|----------------------------|
| Alder                      | Dogbane               | Narrowleaf vetch       | Spotted spurge             |
| Annual yellow sweet clover | Dogfennel             | Nettle                 | Spurge                     |
| Artichoke                  | Elderberry            | Nutgrass               | Spurweed                   |
| Aster                      | English daisy         | Orange hawkweed        | Stinging nettle            |
| Austrian fieldcress        | Fall dandelion        | Oxalis                 | Stinkweed                  |
| Bedstraw                   | False dandelion       | Oxeye daisy            | Stitchwort                 |
| Beggartick                 | False flax            | Parsley-piert          | Strawberry clover          |
| Biden                      | False sunflower       | Parsnip                | Sumac                      |
| Bindweed                   | Fiddleneck            | Pearlwort              | Sunflower                  |
| Bird vetch                 | Field bindweed        | Pennycress             | Sweet clover               |
| Bitterweed                 | Field pansy           | Pennywort              | Tall nettle                |
| Bitter wintercress         | Flea bane             | Peppergrass            | Tall vervain               |
| Black-eyed Susan           | (daisy)               | Pepperweed             | Tansy mustard              |
| Black medic                | Flixweed              | Pigweed                | Tansy ragwort              |
| Black mustard              | Florida betony        | Pineywoods bedstraw    | Tanweed                    |
| Blackseed plantain         | Florida pusley        | Plains coreopsis       | Tarweed                    |
| Blessed thistle            | Frenchweed            | Plantain               | Thistle                    |
| Blue lettuce               | Galinsoga             | Poison hemlock         | Tick trefoil               |
| Blue vervain               | Garlic mustard        | Poison ivy             | Toadflax                   |
| Box elder                  | Goathead              | Poison oak             | Trailing Crownvetch        |
| Bracted plantain           | Goatsbeard            | Pokeweed               | Tumble mustard             |
| Brassbuttons               | Goldenrod             | Poorjoe                | Tumble pigweed             |
| Bristly oxtongue           | Ground ivy            | Povertyweed            | Tumbleweed                 |
| Broadleaf dock             | Gumweed               | Prickly lettuce        | Velvet leaf                |
| Broadleaf plantain         | Hairy bittercress     | Prickly sida           | Venice mallow              |
| Broomweed                  | Hairy fleabane        | Primrose               | Veronica                   |
| Buckhorn                   | Hawkweed              | Prostrate knotweed     | Vervain                    |
| Buckhorn plantain          | Healall               | Prostrate pigweed      | Vetch                      |
| Bulbous buttercup          | Heartleaf drymary     | Prostrate spurge       | Virginia buttonweed        |
| Bull nettle                | Hedge bindweed        | Prostrate vervain      | Virginia creeper           |
| Bull thistle               | Hedge mustard         | Puncture vine          | Virginia pepperweed        |
| Burdock                    | Hemp                  | Purslane               | Wavyleaf bullthistle       |
| Burning nettle             | Henbit                | Ragweed                | Western clematis           |
| Bur ragweed                | Hoary cress           | Red clover             | Western salsify            |
| Burweed                    | Hoary plantain        | Redroot pigweed        | White clover               |
| Buttercup                  | Hoary vervain         | Red sorrel             | White mustard              |
| Canada thistle             | Honeysuckle           | Redstem filaree        | Wild aster                 |
| Carolina geranium          | Hop clover            | Rough cinquefoil       | Wild buckwheat             |
| Carpetweed                 | Horsenettle           | Rough fleabane         | Wild carrot                |
| Catchweed bedstraw         | Horsetail             | Roundleafed marigold   | Wild four-o'clock          |
| Catsear                    | Indiana mallow        | Rush                   | Wild garlic                |
| Catnip                     | Ironweed              | Russian pigweed        | Wild geranium              |
| Chickweed                  | Jewelweed             | Russian thistle        | Wild lettuce               |
| Chicory                    | Jimsonweed            | St. Johnswort          | Wild marigold              |
| Cinquefoil                 | Kochia                | Scarlet pimpernel      | Wild mustard               |
| Clover                     | Knawel                | Scotch thistle         | Wild onion                 |
| Cockle                     | Knotweed              | Sheep sorrel           | Wild parsnip               |
| Cocklebur                  | Lambsquarter          | Shepherdspurse         | Wild radish                |
| Coffeebean                 | Lespedeza             | Slender plantain       | Wild rape                  |
| Coffeeweed                 | Locoweed              | Smallf lower galinsoga | Wild strawberry            |
| Common chickweed           | Lupine                | Smartweed              | Wild sweet potato          |
| Common mullein             | Mallow                | Smooth dock            | Wild vetch                 |
| Common sowthistle          | Marshelder            | Smooth pigweed         | Willow                     |
| Corn Chamomile             | Matchweed             | Sneezeweed             | Witchweed                  |
| Creeping jenny             | Mexicanweed           | Southern wild rose     | Wooly morningglory         |
| Crimson clover             | Milk vetch            | Sowthistle             | Woodsorrel                 |
| Croton                     | Milkweed bloodf lower | Spanishneedle          | Wooly croton               |
| Cudweed                    | Morningglory          | Spatterdock            | Wooly plantain             |
| Curly dock                 | Mousear chickweed     | Speedwell              | Wormseed                   |
| Curly indigo               | Mugwort               | Spiny Amaranth         | Yarrow                     |
| Dandelion                  | Musk thistle          | Spiny cocklebur        | Yellow rocket              |
| Dead nettle                | Mustard               | Spotted catsear        | Yellowflower pepperweed    |
| Dock                       | Narrowleaf plantain   | Spotted knapweed       | and other broadleaf weeds. |

Commercial

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.8 pounds per 250 square feet.

**SPREADER SETTINGS  
ONE PASS APPLICATION**

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.)\* pounds of nitrogen per 1,000 square feet (optional P2O5 and K2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

**Limitations:** The maximum rate per application is 3.2 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 6.4 pounds per 1,000 square feet, excluding spot treatments. Minimum of 30 days between applications.

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

**Environmental Benefits**

(Grasscycling Logo)

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & Other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**For nonrefillable bags of granular or dry formulation**

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

or

**Solid dilutable formulations in nonrefillable plastic containers**

**CONTAINER DISPOSAL:** Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**CONTAINER DISPOSAL:** Nonrefillable containers larger than 5 Gallons. Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV01110A)

80A14

Commercial

**OPTIONAL SPREADER INSTRUCTIONS:**

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

**\*NOTE:** Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.  
Chlorine declaration is



9 of 14

Residential

# SWEET SIXTEEN WEED AND FEED WITH VIPER™

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.  
Treats (or covers) 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

**ACTIVE INGREDIENTS:**

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	1.346%
2,4-Dichlorophenoxyacetic Acid*	0.182%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	0.359%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.090%
<b>OTHER INGREDIENTS:</b>	<b>98.023%</b>
<b>TOTAL:</b>	<b>100.000%</b>

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	1.076%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.359%
***3,6-Dichloro-o-anisic Acid	0.090%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

**GUARANTEED ANALYSIS:**

Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Primary Plant Foods derived from (List sources).	
Chlorine (Cl) Maximum	X%
The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.	
Fertilizer guaranteed by (Name of fertilizer registrant)	

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION - PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-401  
EPA EST. NO. 228-IL-1

MANUFACTURED FOR  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS 16 LBS. 000228-00401.RES 20100111- EPA FPL

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
MAR 19 2010

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-401

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION - PRECAUCION**

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**FIRST AID**

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is a stable mixture of four herbicides. It is for use on residential turf.

Not for use on turf grown for sale or other commercial use as sod, or commercial seed production, or for research purposes.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

The maximum number of broadcast applications per treatment site is 2 per year.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and to control such broadleaf weeds as:

- |                            |                      |                       |                            |
|----------------------------|----------------------|-----------------------|----------------------------|
| Alder                      | Dogbane              | Narrowleaf vetch      | Spotted spurge             |
| Annual yellow sweet clover | Dogfennel            | Nettle                | Spurge                     |
| Artichoke                  | Elderberry           | Nutgrass              | Spurweed                   |
| Aster                      | English daisy        | Orange hawkweed       | Stinging nettle            |
| Austrian fieldcress        | Fall dandelion       | Oxalis                | Stinkweed                  |
| Bedstraw                   | False dandelion      | Oxeye daisy           | Stitchwort                 |
| Beggartick                 | False flax           | Parsley-piert         | Strawberry clover          |
| Biden                      | False sunflower      | Parsnip               | Sumac                      |
| Bindweed                   | Fiddleneck           | Pearlwort             | Sunflower                  |
| Bird vetch                 | Field bindweed       | Pennycress            | Sweet clover               |
| Bitterweed                 | Field pansy          | Pennywort             | Tall nettle                |
| Bitter wintercress         | Flea bane            | Peppergrass           | Tall vervain               |
| Black-eyed Susan           | (daisy)              | Pepperweed            | Tansy mustard              |
| Black medic                | Flixweed             | Pigweed               | Tansy ragwort              |
| Black mustard              | Florida betony       | Pineywoods bedstraw   | Tanweed                    |
| Blackseed plantain         | Florida pusley       | Plains coreopsis      | Tarweed                    |
| Blessed thistle            | Frenchweed           | Plantain              | Thistle                    |
| Blue lettuce               | Galinsoga            | Poison hemlock        | Tick trefoil               |
| Blue vervain               | Garlic mustard       | Poison ivy            | Toadflax                   |
| Box elder                  | Goathead             | Poison oak            | Trailing Crownvetch        |
| Bracted plantain           | Goatsbeard           | Pokeweed              | Tumble mustard             |
| Brassbuttons               | Goldenrod            | Poorjoe               | Tumble pigweed             |
| Bristly oxtongue           | Ground ivy           | Povertyweed           | Tumbleweed                 |
| Broadleaf dock             | Gumweed              | Prickly lettuce       | Velvet leaf                |
| Broadleaf plantain         | Hairy bittercress    | Prickly sida          | Venice mallow              |
| Broomweed                  | Hairy fleabane       | Primrose              | Veronica                   |
| Buckhorn                   | Hawkweed             | Prostrate knotweed    | Vervain                    |
| Buckhorn plantain          | Healall              | Prostrate pigweed     | Vetch                      |
| Bulbous buttercup          | Heartleaf drymary    | Prostrate spurge      | Virginia buttonweed        |
| Bull nettle                | Hedge bindweed       | Prostrate vervain     | Virginia creeper           |
| Bull thistle               | Hedge mustard        | Puncture vine         | Virginia pepperweed        |
| Burdock                    | Hemp                 | Purslane              | Wavyleaf bullthistle       |
| Burning nettle             | Henbit               | Ragweed               | Western clematis           |
| Bur ragweed                | Hoary cress          | Red clover            | Western salsify            |
| Burweed                    | Hoary plantain       | Redroot pigweed       | White clover               |
| Buttercup                  | Hoary vervain        | Red sorrel            | White mustard              |
| Canada thistle             | Honeysuckle          | Redstem filaree       | Wild aster                 |
| Carolina geranium          | Hop clover           | Rough cinquefoil      | Wild buckwheat             |
| Carpetweed                 | Horsenettle          | Rough fleabane        | Wild carrot                |
| Catchweed bedstraw         | Horsetail            | Roundleafed marigold  | Wild four-o'clock          |
| Catsear                    | Indiana mallow       | Rush                  | Wild garlic                |
| Catnip                     | Ironweed             | Russian pigweed       | Wild geranium              |
| Chickweed                  | Jewelweed            | Russian thistle       | Wild lettuce               |
| Chicory                    | Jimsonweed           | St. Johnswort         | Wild marigold              |
| Cinquefoil                 | Kochia               | Scarlet pimpernel     | Wild mustard               |
| Clover                     | Knawel               | Scotch thistle        | Wild onion                 |
| Cockle                     | Knotweed             | Sheep sorrel          | Wild parsnip               |
| Cocklebur                  | Lambsquarter         | Shepherdspurse        | Wild radish                |
| Coffeebean                 | Lespedeza            | Slender plantain      | Wild rape                  |
| Coffeeweed                 | Locoweed             | Smallflower galinsoga | Wild strawberry            |
| Common chickweed           | Lupine               | Smartweed             | Wild sweet potato          |
| Common mullein             | Mallow               | Smooth dock           | Wild vetch                 |
| Common sowthistle          | Marshelder           | Smooth pigweed        | Willow                     |
| Corn Chamomile             | Matchweed            | Sneezeweed            | Witchweed                  |
| Creeping jenny             | Mexicanweed          | Southern wild rose    | Woolly morningglory        |
| Crimson clover             | Milk vetch           | Sowthistle            | Woodsorrel                 |
| Croton                     | Milkweed bloodflower | Spanishneedle         | Wooly croton               |
| Cudweed                    | Morningglory         | Spatterdock           | Wooly plantain             |
| Curly dock                 | Mouseear chickweed   | Speedwell             | Wormseed                   |
| Curly indigo               | Mugwort              | Spiny Amaranth        | Yarrow                     |
| Dandelion                  | Musk thistle         | Spiny cocklebur       | Yellow rocket              |
| Dead nettle                | Mustard              | Spotted catsear       | Yellowflower pepperweed    |
| Dock                       | Narrowleaf plantain  | Spotted knapweed      | and other broadleaf weeds. |

Residential

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide, Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.8 pounds per 250 square feet.

**SPREADER SETTINGS  
ONE PASS APPLICATION**

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.)\* pounds of nitrogen per 1,000 square feet (optional P2O5 and K2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

**Limitations:** The maximum rate per application is 3.2 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 6.4 pounds per 1,000 square feet, excluding spot treatments. Minimum of 30 days between applications.

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

**Environmental Benefits**

(Grass cycling Logo)

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & Other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Store in a cool, dry place. Keep container closed and away from food, feedstuffs, and domestic water supplies.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV011110A)

Viper is a trademark of Nufarm Americas Inc.  
The Optical Advantage is a registered trademark of Nufarm Americas Inc.

**OPTIONAL LANGUAGE:**

Contains Natural Non-Burning, Organic Slow-Release Nitrogen and Iron for Deep Greening  
Controls Over 200 Broadleaf Weeds.  
Milorganite. For Better Results. Naturally.

Controls Weeds

Controls Over 200 Weeds! Including Dandelion, Chickweed, Knotweed  
Lush Green Lawn

140F14

Residential

**OPTIONAL SPREADER INSTRUCTIONS:**

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.  
Or
- (Illustration) For Broadcast Spreaders: Apply to the border of the lawn then work toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

**NOTE:** Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration is